Zeitschrift:	Acta Tropica				
Herausgeber:	Schweizerisches Tropeninstitut (Basel)				
Band:	31 (1974)				
Heft:	4				
Artikel:	Miscellanea : Autoimmunity in trypanosome infections. Part IV, Natural infections in hartebeest, hyaena and lion in the Serengeti National Park,				
	Tanzania				
Autor:	Mackenzie, A.R. / Boreham, P.F.L.				
DOI:	https://doi.org/10.5169/seals-311966				

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 13.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Department of Zoology and Applied Entomology, Imperial College Field Station, Silwood Park, Ascot, Berks., England

Autoimmunity in Trypanosome Infections IV. Natural Infections in Hartebeest, Hyaena and Lion in the Serengeti National Park, Tanzania

A. R. MACKENZIE * and P. F. L. BOREHAM

Abstract

Serum samples from 5 hartebeests, 10 hyaenas and 8 lions were examined for the presence of anti-liver and anti-Wassermann autoantibodies. No evidence of autoantibodies was found in hyaenas and lions. Two hartebeests showed low levels of both autoantibodies and two others of only one of the autoantibodies. Since multiple parasitic infections were found in four of the hartebeests and suspected in the other it is not possible to say which if any of the parasites is involved.

Introduction

Autoantibodies are known to be present in a variety of parasitic diseases including malaria (DAVIS 1944), coccidiosis (ASHERSON & ROSE 1963), leprosy (WAGER 1969) and schistosomiasis (BASSILY et al. 1973). Recent studies in trypanosomiasis have shown the presence of a number of autoantibodies in rabbits, cattle and man. These include an anti-liver autoantibody, anti-Wassermann autoantibody and an autoantibody against a component of the fibrinogen/fibrin system (MACKENZIE et al. 1973; MACKENZIE & BOREHAM 1974; BOREHAM & FACER 1974).

Since a number of serum samples were available from naturally infected wild animals it was decided to examine these for the presence of autoantibodies as little is known about the immune response to trypanosomes in reservoir hosts.

Material and Methods

Collection of samples

Serum samples were collected from 5 hartebeests (Alcelaphus buselaphus), 10 hyaenas (Crocuta crocuta) and 8 lions (Panthera leo) in the Serengeti National Park during the 1971 trypanosomiasis survey organised by Professor Geigy (GEIGY et al. 1973).

Animals were captured as described by GEIGY & KAUFFMANN (1973). Blood samples were removed from the jugular veins of hartebeests and hyaenas and from the vena saphena of the lions. The blood was allowed to clot at ambient temperature, stored on ice until the serum could be separated by centrifugation and the serum maintained at -20 °C until tested.

^{*} Present address: Pathology Department, Allen & Hanburys Research Ltd., Ware, Herts., England.

Anti-liver autoantibody

The complement fixation test (CFT) was used to measure the amount of anti-liver autoantibody present in the serum (MACKENZIE & BOREHAM 1974). Saline extracts of ox, hyaena and cat livers were used as antigens respectively in tests for hartebeest, hyaena and lion autoantibodies.

Anti-Wassermann autoantibody

A CFT was employed using cardiolipin (Wellcome Reagents Ltd.) as antigen for all three species (MACKENZIE & BOREHAM 1974).

Results

The table shows the results of the parasitological studies made by Professor Geigy and his colleagues (GEIGY & KAUFFMANN 1973) and the autoantibody measurements.

All the animals examined were infected with piroplasms, probably *Babesia* sp. in lions and hyaenas and *Theileria* sp. in hartebeests. All the lions and hyaenas were infected with *Hepatozoon* sp. and in addition all the hyaenas and four of the five hartebeests had microfilariae present in their blood. Trypanosomes were also common in these animals: 2 hartebeest were infected with *Trypanosoma brucei* and one with *T. congolense*. *T. brucei* was found in 3 hyaenas, *T. congolense* in two hyaenas and two hyaenas had mixed infections. One hyaena No. 250 had trypanosomes present in the blood but it was not possible to identify the species because of damage during preparation of the slide. Six lions were infected with *T. brucei* and one with *T. congolense*. The uninfected lion No. 280 was a young lion not more than 8 months of age. No investigation of gastrointestinal helminths was made.

There appears to be no rise in either anti-liver or anti-Wassermann autoantibodies in hyaenas or lions except for hyaena 294 which showed a small elevation of anti-Wassermann antibody levels. This hyaena was not infected with trypanosomes.

Two hartebeests (253 and 258) showed slight increases in both anti-liver and anti-Wassermann autoantibodies while 257 had increased anti-liver autoantibody and 259 anti-Wassermann autoantibody only. Hartebeest No. 255 showed no increase in either of these autoantibodies.

Discussion

The results of these studies indicate that in lions and hyaenas the natural parasite infections do not elicit the formation of autoantibodies as is seen in trypanosomiasis of rabbits, cattle and man (MACKENZIE et al. 1973).

In contrast it appears that a proportion of the hartebeests develop autoantibodies but it is not possible to suggest which of the multiple parasite infections is responsible. It is possible that helminth infections were involved but since these animals were only immobilised for the removal of blood samples and not killed, it was not possible to investigate this aspect.

Whether these differences are due to differences in the immune systems of these hosts or to differences in the parasitological infections is not known. It is possible that in reservoir hosts, where trypanosomes appear not to be highly pathogenic these autoantibodies do not develop. In man and rabbits where marked pathological symptoms are usually seen it is believed that trypanosomes exert some kind of adjuvant effect breaking down self-tolerance with subsequent production of autoantibodies. Cattle appear to show an intermediate effect where a single infection with either *T. brucei* or *T. congolense* produces only a small transient rise in anti-liver autoantibody but repeated fly borne challenge and periodic

Host		Infections				CTF Titres			
No.	Species	Tryp	MF	Нер	Piro	a-C ¹	anti- liver	anti- Wass.	
253 255	Hartebeest		-+	_	+ +	2	16 2	32 2	
257	Alcelaphus	b	+		+	2	8	< 2	
258	buselaphus	b	+		+	2	8	32	
259	-	c	+	_	+	< 2	2	16	
239		bc	÷	+	+	4	< 2	< 2	
248		b	+	+	+	4	< 2	< 2	
249	Hyaena		+	+	+	2	< 2	< 2	
250	States - servered - servered	T ?	+	÷	+	2	< 2	< 2	
251		b	+	+	+	4	< 2	< 2	
252	Crocuta	с	+	+	+	2	< 2	< 2	
284	crocuta	bc	+	+	+	4	< 2	< 2	
293		b	+	+	+	8	< 2	< 2	
294			+	+	+	4	< 2	16	
295		с	+	+	+	4	< 2	< 2	
245		b	_	+	+	4	< 2	< 2	
246		с	-	+	+	< 2	< 2	< 2	
247	Lion	b	-	+	+	< 2	< 2	< 2	
272		b		+	+	4	< 2	< 2	
276	Panthera	b	_	+	+	2	< 2	< 2	
277	leo	b	—	+	+	2	< 2	< 2	
278		b		+	+	2	< 2	< 2	
280		-		+	+	2	< 2	< 2	
Tryp = Trypanosomes: b = T. brucei c = T. congolense T? = Unidentified Trypanosoma									
MF Hep Piro a-C ¹ Anti	$= Mid = He_{\mu}$ $= Pire = ant$ $-Wass = ant$	crofilaria <i>patozoon</i> oplasm i-compler i-Wasserr	nentary nann au	titre toantibo	dy titre				

Table. Infections and autoantibody titres in hartebeests, hyaenas and lions captured in the Serengeti National Park, Tanzania

treatment with Diminazene aceturate (Berenil) results in much higher autoantibody levels (MACKENZIE et al., in preparation).

The low levels of autoantibodies recorded here in hartebeest infections may be comparable to the single infections in cattle when the immunogenic stimulus is insufficient to completely break tolerance.

Acknowledgements

We wish to thank Professor Geigy for allowing one of us (ARM) to join his survey in the Serengeti National Park and for all his advice. Our thanks are due to the Director of the East African Trypanosomiasis Research Organisation for providing laboratory space for part of this work. We are extremely grateful to the Overseas Development Administration of the Foreign and Commonwealth Office and the Medical Research Council for financial support. This work was undertaken while one of us (ARM) held a Medical Research Council Scholarship for training in research methods.

References

- ASHERSON, G. L. & ROSE, M. E. (1963). Autoantibody production in rabbits. III. The effect of infection with *Eimeria stiedae* and its relation to natural antibody. – Immunology 6, 207–216.
- BASSILY, S., MOHRAN, Y. Z., SAID, M. I., MOUSSA, M., ISKANDER, F. & MOORO, H. (1973). Autoantibodies in schistosomal liver fibrosis. J. trop. Med. Hyg. 76, 153–155.
- BOREHAM, P. F. L. & FACER, C. A. (1974). Autoimmunity in trypanosome infections. II. Anti-fibrin/fibrinogen (anti-F) autoantibody in *Trypanosoma (Trypanozoon) brucei* infections of the rabbit. – Int. J. Parasit. 4, 601–607.
- DAVIS, B. D. (1944). Biologic false positive serologic tests for syphilis. Medicine (Baltimore) 23, 359–414.
- GEIGY, R. & KAUFFMANN, M. (1973). Sleeping sickness survey in the Serengeti Area (Tanzania) 1971. I. Examination of large mammals for trypanosomes. Acta trop. 30, 12–23.
- GEIGY, R., KAUFFMANN, M., ROGERS, D., BOREHAM, P. F. L. & BERTRAM, B. C. R. (1973). Sleeping sickness survey in the Serengeti Area (Tanzania) 1971. Acta trop. 30, 12–48.
- MACKENZIE, A. R. & BOREHAM, P. F. L. (1974). Autoimmunity in trypanosome infections. I. Tissue autoantibodies in *Trypanosoma (Trypanozoon) brucei* infections of the rabbit. Immunology 26, 1225–1238.
- MACKENZIE, A. R., BOREHAM, P. F. L. & FACER, C. A. (1973). Autoantibodies in African trypanosomiasis. Trans. roy. Soc. trop. Med. Hyg. 67, 268.
- WAGER, O. (1969). Immunological aspects of leprosy with special reference to autoimmune disease. Bull. Wld Hlth Org. 41, 793–804.