

Miscellanea : Isoelectric focusing in the taxonomy of bulinid snails

Autor(en): **Saladin, B. / Degrémont, A. / Weiss, N.**

Objekttyp: **Article**

Zeitschrift: **Acta Tropica**

Band (Jahr): **33 (1976)**

Heft 4

PDF erstellt am: **05.06.2024**

Persistenter Link: <https://doi.org/10.5169/seals-312240>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

Miscellanea

Swiss Tropical Institute, Socinstrasse 57, CH-4051 Basel, Switzerland

Isoelectric Focusing in the Taxonomy of Bulinid Snails*

B. SALADIN, A. DEGRÉMONT and N. WEISS

The identification of Bulinid snails is still difficult and their taxonomy is not always well established (SOUTHGATE & KNOWLES, 1975). Field workers who are not familiar with morphological taxonomy are frequently confronted with this problem. This was the case in Madagascar, where it was very difficult to distinguish *Bulinus obtusispira*, the intermediate host of *Schistosoma haematobium*, from *B. liratus* (DEGRÉMONT, 1973).

Several attempts to develop serological and biochemical taxonomy have been made (WRIGHT, 1966) using different methods such as electrophoresis on cellulose acetate membranes (WRIGHT & ROSS, 1967), enzyme electrophoresis (COLES, 1970), gel diffusion (BURCH & LINDSAY, 1970) and disc electrophoresis (DAVIS & LINDSAY, 1967). We have already studied the abilities of all the methods mentioned above to distinguish between Bulinid snails from Madagascar (WEISS et al., 1974). Recently, the method of isoelectric focusing was used in bacterial (MATTHEW & HARRIS, 1975) and bird taxonomy (FRÉLIN et al., 1973).

The concept of isoelectric focusing in gels (IEF) was developed by SVENSSON (1961). We used the method described by RIGHETTI & DRYSDALE (1974) employing the Multiphor (LKB 2117) apparatus and Ampholine^R polyacrylamide gel plates with the pH-range from 3.5 to 9.5 (PAG plate Nr. 1804-101 by LKB). The d.c. power supply was a PS 10A multistab capable of delivering 1000 V.

The snails studied were: *Bulinus obtusispira* and *Bulinus liratus* (Tanandava, Madagascar). We compared their patterns with those of snails of three Bulinid groups, namely: *Bulinus tropicus*¹ (Zambia), *Bulinus cernicus*¹ (Mauritius) and *Bulinus nasutus* (Ifakara, Tanzania). For the experiment an egg-clutch was simply crushed on a filter-paper, which was laid upon the PAG plate. The number of eggs per clutch should be between three and ten to get the main bands and between ten and twenty to get the minor bands. Up to 48 samples could be examined on one plate. The egg clutches can be preserved in tap water at 4°C, in glycerol or in physiological salt solution at room temperature to inhibit the development of the snail embryos.

The patterns obtained are presented in figures 1 and 2. With the IEF-method we found at least twice the number of separated proteins as with disc-electrophoresis. With the procedure described above the reproducibility was very good. Although the general pattern is very complex, some of the most characteristic bands are easily identified. In agreement with the systematics of the Bulinid group *B. liratus* and *B. tropicus* are showing very similar patterns. The pattern of *B. obtusispira* is completely different from that of *B. (Physopsis) nasutus*. The distinction between *B. obtusispira* and *B. liratus* can be made accurately with only one egg-clutch. *B. cernicus* shows the pattern of a species belonging to the *forskali* group.

* Supported by a grant of the Schweizerischer Nationalfonds (3.764.72)

¹ Kindly supplied by G. Mandahl-Barth (Danish Bilharziasis Laboratory, Charlottenlund, Denmark).




Fig. 1. IEF patterns of egg proteins from different Bulinid snails.




Fig. 2. Schematic representation of IEF shown in fig. 1.

With eggs stored for two weeks the results were not modified as long as the embryogenesis had been blocked.

In conclusion, isoelectric focusing appears to be an accurate and reliable technique for snail taxonomy. It has the advantage that a large number of specimens can be tested simultaneously and in a short period of time. Although this method is rather sophisticated, the facts that commercial plates are available and that eggs or living snails can be stored or posted, make it especially useful for taxonomical research and epidemiological work at large scale.

References

- BURCH, J. B. & LINDSAY, G. K. (1970). An immuno-cytological study of *Bulinus* species. – *Malacol. Rev.* 3, 1–18.
- COLES, C. G. (1970). Enzyme electrophoresis and speciation of *Schistosoma* intermediate hosts. – *J. Parasit.* 61, 19–25.
- DAVIS, G. M. & LINDSAY, G. K. (1967). Disc electrophoretic analysis of molluscan individuals and populations. – *Malacologia* 5, 311–334.
- DEGRÉVONT, A. A. (1973). Mangoky Project, 273 pp. – Basle: Swiss Tropical Institute.
- FRÉLIN, C., GASQUEZ, J. & DORST, J. (1973). Taxinomie. Essai d'utilisation de la focalisation iso-electric en taxinomie animale et végétale. – *C. R. Acad. Sci. Paris*, 277, 17.
- MATTHEW, M. & HARRIS, A. M. (1976). Identification of β -lactamases by isoelectric focusing: Correlation with bacterial taxonomy. – *J. gen. Microbiol.* 94, 55–67.
- RIGHETTI, P. G. & DRYSDALE, J. W. (1974). Isoelectric focusing in gels. – *J. Chromatogr.* 98, 271–321.
- SOUTHGATE, V. R. & KNOWLES, R. J. (1975). The intermediate hosts of *Schistosoma bovis* in western Kenya. – *Trans. roy. Soc. trop. Med. Hyg.* 69, 356–357.
- SVENSSON, H. (1961). Isoelectric fractionation, analysis and characterization of ampholytes in natural pH gradients. I. The differential equation of solute concentrations at a steady state and its solution for simple cases. – *Acta chem. scand.* 15, 325.
- WEISS, N., DEGRÉVONT, A. A., TANNER, K. & ZUMSTEIN, A. (1974). Susceptibility to *Schistosoma haematobium* and comparative antigenic analysis of two closely related *Bulinus* species from Madagascar. – ICOPA, München.
- WRIGHT, C. A. (1966). Experimental taxonomy: A review of some techniques and their applications. – *Int. Rev. gen. exp. Zool.* 2, 1–42.
- WRIGHT, C. A. & ROSS, G. C. (1967). Serological studies on the taxonomy of planorbid snails. – *J. Zool. (London)* 151, 489–495.