

Zeitschrift: Acta Tropica
Herausgeber: Schweizerisches Tropeninstitut (Basel)
Band: 40 (1983)
Heft: 1

Artikel: "Trypanosoma cruzi" : variation in susceptibility of inbred strains of rats
Autor: Rivera-Vanderpas, M.T. / Rodriguez, A.M. / Afchain, D.
DOI: <https://doi.org/10.5169/seals-313106>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Centre d'Immunologie et de Biologie Parasitaire, Institut Pasteur, B.P. 3415, 59019 Lille Cedex, France

***Trypanosoma cruzi*: variation in susceptibility of inbred strains of rats**

M. T. RIVERA-VANDERPAS, A. M. RODRIGUEZ, D. AFCHAIN,
H. BAZIN, A. CAPRON

Summary

Eight strains of male rats (AUG, BN, LEW, LIS, WAG, F 344, LOU/M, DA) and 3 strains of female rats (LEW, F 344, KGH) were challenged with 15×10^4 trypomastigotes of the Tehuantepec strain of *T. cruzi*. Parasitemia and mortality were observed for 60 days. Varying degrees of susceptibility were demonstrated between strains: complete resistance (R), no parasitemia; low resistance (LR), mild parasitemia; and no resistance (NR), high parasitemia. The differences in susceptibility to *T. cruzi*, using inbred strains of rats, were unrelated to Rt-1 haplotype. However, the level of parasitemia and host survival are not necessarily related; both male and female F 344 hosts are susceptible (NR) to *T. cruzi* but only females survive.

Key words: *Trypanosoma cruzi*; Chagas' disease; rats, inbred strains; major histocompatibility complex, MHC; genetic locus, Rt-1 haplotype; genes for immune response, Ir; susceptibility, host survival.

Introduction

Susceptibility to infection with *Trypanosoma cruzi* varies depending on the species and strain of host (Goble, 1970; Pizzi et al., 1954; Trischmann et al., 1978; Wrightsman et al., 1982). The immunological system plays a role in natural resistance against *T. cruzi* infection, but other factors are probably involved. A major histocompatibility complex (MHC) has been demonstrated in the majority of vertebrates (Vaiman, 1978) and two systems have been studied in depth: the H-2 system of mice (Colombani, 1975) and the human HLA

Correspondence: Doctor D. Afchain, Centre d'Immunologie et de Biologie Parasitaire, Institut Pasteur, 15 rue Camille Guérin, BP. 3415, F-59019 Lille Cedex, France

system (Dausset, 1976). Although the functions of these systems are similar for these two species, the organisation of loci within the MHC is different. The MHC of rats, known as Rt-1 (Report of the first international ..., 1978), seems to resemble the human MHC more closely than the MHC of mice (Gill et al., 1978). It also represents a polygenic control (Gill and Kunz, 1971) and plays an important role in the defence mechanism of the animal, with regard to different biological processes. It is currently thought to contain two regions, A and B (Gill et al., 1978).

Most studies on *T. cruzi* infection have been carried out on mice (Hauschka, 1947; Pizzi et al., 1954; Goble, 1951), a highly susceptible host. On the other hand, rats are relatively resistant to this infection and have the advantage of developing a chronic phase of Chagas' disease, which permits the analysis of a certain number of immunological phenomena that appear during the course of the infection (Pizzi et al., 1953).

It has been observed that the resistance generally exhibited by rats does not always occur and that strains of male F 344 rats are particularly susceptible to infection (Rodriguez et al., 1981). Consequently, the susceptibility of different inbred strains of rats to *T. cruzi* infection has been investigated.

Materials and Methods

Animals. Eight different inbred strains of male rats which differed in their Rt-1 haplotype were used for the investigation: August (AUG), Brown Norway (BN), Lewis (LEW), Hooded Lister (LIS), Wistar Ag (WAG) obtained from Service de Génétique, CNRS, Orléans, France; Fischer (F 344) obtained from Iffa Credo, L'Arbresle, France; Louvain (LOU/M) and Dark Agouti (DA) obtained from Institut Pasteur, Lille, France. Each group of animals to be infected was carefully matched for weight (220–300 g) and age (12–16 weeks).

In the second part of the study, three strains of female rats with very close genetic characteristics on their MHC were chosen: Lewis (LEW) obtained from CNRS, Orléans, France; Fischer (F 344) obtained from Iffa Credo, France; and KGH obtained from Institut Pasteur, Lille France. These also were matched for weight (150–200 g) and age (12–16 weeks).

Parasites. The Tehuantepec strain of *T. cruzi* (isolated from *Triatoma* from Mexico by Brumpt) was maintained by serial passage in male F 344 rats by the method described for mice by Pizzi and Prager (1952). For infection of the animals, blood was drawn by cardiac puncture from male F 344 rats, mixed with heparin and diluted in Alsever solution. Rats were inoculated intraperitoneally (i.p.) with 1 ml of this diluted blood containing 15×10^4 trypomastigotes/ml.

Parasitemia: In order to determine the course of parasitemia, tail blood (5 μ l) was taken with a micropipette and smears prepared. 60 microscope fields were examined and the number of trypanosomes per milliliter was estimated by the method of Pizzi and Prager (1952) and Philipps (1960).

Results

Variation in susceptibility of inbred male rat strains

The course of parasitemia obtained in the different inbred male rat strains studied is shown in Fig. 1. Susceptibility was classified into three groups: the first group (AUG, LOU/M and DA) showed resistance and did not develop a




Fig. 1




Fig. 2

Fig. 1. Course of *Trypanosoma cruzi* infection in different male rat strains. Male rats aged 3–4 months, given 15×10^4 trypanosomes i.p.; (○—○) NR = no resistance: male F 344 rats; (-----) LR = low resistance: no statistical difference (variance analyses) was found between strains with low parasitemia (LEW, WAG, BN, LIS), LEW was chosen as representative for the four strains; (---) R = resistant: male AUG, LOU/M and DA rats. Parasitemia expressed as geometric mean. † Death in 34 ± 4 days after infection.

Fig. 2. Course of *Trypanosoma cruzi* in F 344 rats. F 344 rats aged 3–4 months, given 15×10^4 trypanosomes i.p. Male rats (—), female rats (---). Parasitemia expressed as geometric mean. † Death in 34 ± 4 days after infection.

parasitemia: it was considered as “resistant (R)”. The second (LIS, LEW, WAG, BN) showed low resistance and developed a mild parasitemia between the 11th and 42nd day: it was considered as “low resistant (LR)”. In contrast, the third group (F 344) was exceptional, showing no resistance; it presented a high level of susceptibility with high parasitemia: it was considered as “no resistance (NR)”.

All the inbred male rat strains survived except for the F 344 strain all of which died between the 34th and 42nd day of the infection. The mortality and susceptibility is shown in Table 1. Strains with the same MHC (Rt-1^c: AUG and LIS, or Rt-1^u: WAG and LOU/M) differed in the outcome of infection.

Variation in susceptibility of inbred female rat strains

Following the preliminary results obtained from the different inbred male rat strains, 3 female strains were used. As shown in Table 2, all had identical MHC B regions and different A regions (A region of the F 344 strain is a variant of the Rt-1^l haplotype of the LEW strain and the KGH strain shows a Rt-1^g haplotype) (Gill and Kunz, 1978).

Table 1. Male rat strains aged 3–4 months given 15×10^4 trypanosomes of *T. cruzi* i.p.

	Strains	AUG	LIS	LEW	F 344	WAG	LOU/M	DA	BN
	Rt-1	C	C	l	lv1	U	U	a	n
Outcome of infection		R	LR	LR	NR	LR	R	R	LR
Mortality		0/6	0/6	0/6	6/6	0/6	0/6	0/6	0/6

Six rats per group; R = resistant; LR = low resistance; NR = no resistance.

Table 2. Female rats aged 3–4 months given 15×10^4 trypanosomes *T. cruzi* i.p.

	Strains	F 344	LEW	KGH
	Rt-1	lv1	l	g
	Region A	lv1	l	g
	Region B	l	l	l
Outcome of infection		NR	LR	LR
Mortality		0/7	0/7	0/7

Seven rats per group; LR = low resistance; NR = no resistance.

The analysis of the infection of these strains is shown in Table 2. The LEW strain and KGH strain demonstrated a low level of parasitemia (low resistance: LR), whereas the F 344 strain showed a high parasitemia (no resistance: NR). All female F 344 rats survived the infection and showed the highest parasitemia between the 34th and 42nd day; subsequently the parasite numbers diminished progressively until they totally disappeared (Fig. 2).

Discussion

Biological aspects

Most of the inbred strains were either resistant (AUG, LOU/M and DA) or showed a low level of parasitemia (LIS, LEW, WAG, BN). This confirms former observations with regard to the resistance of rats to *T. cruzi* infection (Kolodny, 1940; Pizzi et al., 1953). Experiments carried out with the male F 344 rats indicate that this strain develops an acute phase of the disease and dies during this phase. However, the other strains may exhibit a chronic phase. This should be investigated by physiological and histopathological examination of their hearts.

Genetic aspects

Strains with the same MHC responded differently (LOU/M and WAG, AUG and LIS). In order to show the role of the MHC haplotypes more precise-

ly, a further experiment was carried out, in which 3 strains with genetically close MHC haplotypes were infected. According to Table 2, the hypothesis was that if the B region played an important role in the control of *T. cruzi* infection, the F 344 and LEW strains should show the same susceptibility; this was not the case. The other possibility was that the A region played an important role, which would mean that the LEW and KGH strains should show different susceptibilities; and this was not the case.

In conclusion, it may be stated that the MHC does not seem to play an important role in *T. cruzi* susceptibility. These results agreed with those of Trischmann et al. (1978), using inbred strains of mice, which showed that the principal genetic determinant of resistance was not linked with an H-2 haplotype. But another paper (Wrightsmann et al., 1982) shows that the level of parasitemia was not linked with the H-2 region and H-2-linked gene(s) was involved in survival of the infection. We have clearly demonstrated that the level of parasitemia and host survival are not necessarily related. The course of *T. cruzi* infection in female and male inbred rats of strain F 344 was not the same (see Fig. 2). Both showed a high degree of parasitemia, but all the males died between the 34th and 42nd day of the infection, whereas all the females survived the infection, with total disappearance of the parasitemia.

Both male and female F 344 hosts are highly susceptible to *T. cruzi* but the host survival are different. This distinction between susceptibility and host survival corroborate data on *T. cruzi* infected mice in a recent paper of Wrightsmann et al. (1982); the MHC is probably not important for susceptibility but the MHC could still be of value in survival in mice.

The difference of host survival in F 344 rats does seem to depend on sex (a hormonal factor or a sex-linked genetic factor). Finally, complementary experiments (the use of F1 and back crosses) might lead to the discovery of the mechanisms that control infection with *T. cruzi*.

Acknowledgments

The work reported here was supported by grants from INSERM (U 167) and CNRS (ERA 422). We wish to thank Christine Mazingue for assistance in preparing this manuscript. The encouragement and interest of Ferruccio Santoro, the technical collaboration of Marie-Ange Hospied, Marc Loyens and the staff of the animal unit are highly appreciated.

Colombani J.: Immunogénétique du complexe majeur d'histocompatibilité (H-2) chez la souris. Bull. Inst. Pasteur 73, 305–356 (1975).

Dausset J.: Le complexe HLA. I. Immunogénétique du système HLA. Les séries alléliques HLA-A, B et C. Nouv. Presse méd. 5, 1301–1304 (1976).

Gill III T. J., Kunz H. W.: Genetic and cellular factors in the immune response. I. Evidence for the polygenic control of the antibodies response from further breeding studies and from pedigree analyses. J. Immunol. 106, 980–992 (1971).

Gill III T. J., Kunz H. W.: Recombinants strains: rat. In: Inbred and genetically defined strains of

- laboratory animals. Part I: mouse and rat, ed. by L. A. Phillip, p. 260. Fed. Amer. Soc. exp. Biol. Maryland (1978).
- Gill III T. J., Cramer D., Kunz H. W.: The major histocompatibility complex comparison in the mouse, man and rat. *Amer. J. Path.* 90, 751–777 (1978).
- Goble F. C.: Studies on experimental Chagas'disease in mice in relation to chemotherapeutic testing. *J. Parasit.* 37, 408–414 (1951).
- Goble F. C.: South American trypanosomiasis. In: Immunity to parasitic animals, ed. by G. J. Jackson, R. Herman and I. Springer, p. 597–689. Appleton, New York 1970.
- Hauschka T. S.: Sex of host as a factor in Chagas'disease. *J. Parasit.* 33, 399–404 (1947).
- Kolodny M.: Studies on age resistance against *Trypanosoma* infections. VII. The influence of age upon the immunological response of rats to infection with *Trypanosoma cruzi*. *Amer. J. Hyg.* 31C, 1–8 (1940).
- Phillips N. R.: Experimental studies on the quantitative transmission of *Trypanosoma cruzi*: considerations regarding the standardization of materials. *Ann. trop. Med. Parasit.* 54, 60–70 (1960).
- Pizzi T., Prager R.: Estabilizacion de la virulencia de una cepa de *Trypanosoma cruzi* por pasage seriado en ratones de constitucion genetica uniforme: analisis cuantitativo del curso de la infeccion. *Biologica* 16/17, 3–12 (1952).
- Pizzi T., Rubio M., Knierim F.: Contribucion al conocimiento de los mecanismos inmunitarios de la enfermedad de Chagas experimental de la rata. *Bol. Inf. parasit. chil.* 8, 66–72 (1953).
- Pizzi T., Rubio M., Knierim F.: Immunology of Chagas'disease. *Revista de Parasit.* 4, 578–592 (1954).
- Report of the first International Workshop on alloantigenic systems in the rat. *Transplant. Proc.* 10, 271–285 (1978).
- Rodriguez A. M., Santoro F., Afchain D., Bazin H., Capron A.: *Trypanosoma cruzi* infection in B-cell-deficient rats. *Infect. Immun.* 31, 524–529 (1981).
- Trischmann T., Ranowitz H., Wittner M., Bloom B.: *Trypanosoma cruzi*: Role of the immune response in the natural resistance of inbred strains of mice. *Exp. Parasit.* 45, 160–168 (1978).
- Vaiman M.: Le complexe majeur d'histocompatibilité chez les animaux (homme et souris exceptés). Dans le complexe principal d'histocompatibilité de l'homme (HLA). Cours supérieur d'histocompatibilité. Imprimerie du Signe à Gergy Villago (1978).
- Wrightsmann R., Krassner S., Watson J.: Genetic control of responses to *Trypanosoma cruzi* in mice: multiple genes influencing parasitemia and survival. *Infect. Immun.* 36, 637–644 (1982).