

Orphicarbonate rocks : metamorphic reactions and possible origin

Autor(en): **Trommsdorff, Volkmar / Evans, Bernard W. / Pfeifer, H.-R.**

Objekttyp: **Article**

Zeitschrift: **Archives des sciences [1948-1980]**

Band (Jahr): **33 (1980)**

Heft 1-3

PDF erstellt am: **23.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-739502>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

OPHICARBONATE ROCKS: METAMORPHIC REACTIONS AND POSSIBLE ORIGIN

BY

Volkmar TROMMSDORFF¹, Bernard W. EVANS² and H.-R. PFEIFER³

In a first approximation, ophicarbonate rocks may be regarded as forming part of a multisystem with the components CaO-MgO-SiO₂-CO₂-H₂O. The prograde reactions in this system can be described for the minerals antigorite, diopside, forsterite, talc, calcite, dolomite and magnesite by 30 equilibria only four of which are independent (Trommsdorff and Evans, 1977a). Adding chrysotile to the system requires another equilibrium. Calculation of these equilibria using experimentally calibrated data yields within the limits of experimental error several possible topologies in P-T-X_{CO₂} space for the ophicarbonate phase diagram (Trommsdorff and Evans, 1977b). Only one of these topologies is confirmed by field data (Fig. 1). Ophicarbonate rocks have a limited stability in T-X_{CO₂} space, their equilibrium fluid being always very rich in H₂O.

At surface conditions the pH of fluids emanating from serpentinite and ophicarbonate rocks is considerably higher than in most other rocks (pH 9-12, Pfeifer, 1977). For metamorphic conditions, a similar behavior, i.e. pH 6-7 at 300° C, 0.5-1 kb, in contrast to pH 3-5 in felsic rocks, has been estimated, using the model of fluid-rock interaction proposed by Helgeson (1970) and Helgeson *et al.* (1970), see Table 1. The formation of carbonate minerals in ophicarbonate rocks is conceived to be a precipitation reaction, which can be driven by two processes: (1) Mixing of an alkaline fluid from serpentinites with more acid fluids from felsic or mafic rocks which leads to supersaturation with carbonate minerals, or (2), sudden drop of pressure in a hydrothermal vein system, followed by a separation of the vapour phase (flashing, cf. Arnorsson 1977).

¹ Inst. für Kristallographie und Petrographie, ETH-Zentrum, CH-8092 Zürich

² Dept. of Geological Sciences, University of Washington, Seattle, Wash. 98195, U.S.A.

³ Inst. de Minéralogie, Palais Rumine, CH-1005 Lausanne




TABLE 1. — *Example of a fluid composition in equilibrium with an Antigorite-Tremolite-Diopside rock at 300° C, 500 bar. It has been calculated by simulating an irreversible dissolution of a peridotite, which ends when equilibrium with a serpentinite paragenesis is reached (Pfeifer 1977, 1979).*

A. Total concentrations		B. Distributed among species		
	mole/kg H ₂ O	ppm		
	mole/kg H ₂ O	Activity		
Na	0.5	11 169	Al ³⁺	—
Ca	1.56 × 10 ⁻³	61	Al(OH) ₂ ⁺	—
Mg	8.33 × 10 ⁻⁶	0.19	Al(OH) ₄ ⁻	—
SiO ₂	4.67 × 10 ⁻⁴	27	Al(SO ₄) ⁺	—
HS	7.70 × 10 ⁻⁸	0.003	K ⁺	—
SO ₄	9.90 × 10 ⁻⁶	0.92	KCl ^o	—
CO ₃	0.0	0.0	KSO ₄ ⁻	—
Cl	0.5	17 322	Na ⁺	0.47
OH	1.1 × 10 ⁻⁴	1.80	NaCl ^o	0.03
I.st.	0.474	—	Ca ²⁺	1.54 × 10 ⁻³
pH	6.61	—	Ca(OH) ⁺	9.35 × 10 ⁻⁶
			Mg ²⁺	2.87 × 10 ⁻⁶
			Mg(OH) ⁺	5.46 × 10 ⁻⁶
			SiO ₂ aq.	4.68 × 10 ⁻⁴
			HS ⁻	1.47 × 10 ⁻⁸
			SO ₄ ²⁻	9.60 × 10 ⁻⁶
			HSO ₄ ⁻	2.94 × 10 ⁻⁷
			S ²⁻	3.41 × 10 ⁻⁹
			H ₂ S ^o	5.99 × 10 ⁻⁸
			CO ₃ ²⁻	—
			HCO ₃ ⁻	—
			H ₂ CO ₃	—
			Cl ⁻	0.47
			HCl ^o	3.36 × 10 ⁻⁷
			OH ⁻	9.40 × 10 ⁻⁵
			H ₂ O	5.55 × 10 ¹
			H ⁺	3.33 × 10 ⁻⁷

This proposed origin for ophicarbonate rocks excludes of course not other geologically possible origins i.e. sedimentary or tectonic mixing of serpentine plus carbonate minerals.

ACKNOWLEDGEMENTS

The authors acknowledge the support of Schweizerischer Nationalfonds No. 2.115-0.78 and U.S. National Science Foundation (Grant EAR-7803350).

FIG. 1. — Progressive metamorphism of antigorite-ophicarbonate rocks, Val Ventina, Malenco, Italy in comparison to the calculated stable $T\text{-}X_{\text{CO}_2}$ topology for the ophicarbonate system ($\text{CaO}\text{-MgO}\text{-SiO}_2\text{-CO}_2\text{-H}_2\text{O}$) at three kilobars total fluid pressure. Vertical streaks indicate occurrence of assemblages (see also Trommsdorff and Evans, 1977b). Abbreviations: *A* = antigorite, *Cc* = calcite, *Di* = diopside, *Do* = dolomite, *Fo* = forsterite, *M* = magnesite, *Tc* = talc, *Tr* = tremolite.

REFERENCES

- ARNORSSON, S. (1977). Precipitation of calcite from flashed geothermal waters in Iceland. Proc. 2nd Internat. Symp. Water-rock interaction, Strasbourg, III, 152-169.
- HELGESON, H. C. (1970). A chemical and thermodynamic model of ore deposition in hydrothermal systems. *Min. soc. Amer. spec. Pap.*, 3, 155-186.
- HELGESON, H. C., I. BROWN, A. NIGRINI and T. JONES (1970). Calculation of mass transfer in geochemical processes involving aqueous solutions. *Geochim. Cosmochim. Acta*, 34, 569-592.
- PFEIFER, H.-R. (1977). A model for fluids in metamorphosed ultramafic rocks. I. Observations at surface and subsurface conditions (high pH spring waters). *Schweiz. Mineral. Petrogr. Mitt.*, 57, 361-396.
- (1979). Fluid-Gesteins-Interaktion in metamorphen Ultramafititen der Zentralalpen. Ph. Diss. No. 6379, ETH-Zürich, 200 p.
- TROMMSDORFF, V. and B. W. EVANS (1977a). Antigorite-Ophicarbonates: Phase relations in a portion of the system CaO-MgO-SiO₂-H₂O-CO₂. *Contrib. Mineral. Petrol.*, 60, 39-56.
- (1977b). Antigorite-ophicarbonates: contact metamorphism in Valmalenco, Italy. *Contrib. Mineral. Petrol.*, 62, 301-312.