

On the occurrence of *Acacia raddiana* Savi and *A. albida* Delile in Jericho

Autor(en): **Karschon, R.**

Objekttyp: **Article**

Zeitschrift: **Boissiera : mémoires de botanique systématique**

Band (Jahr): **24 (1975-1976)**

Heft 1

PDF erstellt am: **27.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-895521>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

On the occurrence of *Acacia raddiana* Savi and *A. albida* Delile in Jericho¹

R. KARSCHON

SUMMARY

The possibility is discussed that the Lower Jordan Valley may constitute the only area in the Middle East where the ranges of *A. raddiana* and *A. albida* overlap.

RÉSUMÉ

La probabilité est envisagée que la vallée inférieure du Jourdain soit la seule zone où les aires de l'*Acacia raddiana* et de l'*Acacia albida* se chevauchent.

The occurrence of *Acacia albida* Del. in Jericho was reported by Hart (1891) in the following terms: "An acacia near Ain es Sultan was, I believe, *A. albida* Del. gathered previously at Gaza. It was a stunted bush...".

At present, the only acacia found near Ain es Sultan is *A. raddiana* Savi which was already recorded by Dinsmore in Post & Dinsmore (1932; under the name of *A. tortilis* Hayne). The single tree growing in a shallow depression between the SSW slope of the ancient mound of Tel es Sultan and the recently surfaced Jericho-Ramallah road displays a short asymmetrical bole with severe lean; its height of 11.5 m and girth at breast height of 222 cm makes it the largest tree of this species in Palestine (Plate I).

In order to determine if this tree could have been confused by Hart (1891) with *A. albida*, two increment cores were taken at about 1 m above ground, on opposite sides of the stem (for technical reasons it was not possible to extract complete cores). Since growth rings are indistinct, counts were made (by Mr. K. Tischler) of the broad concentric bands of aliform-confluent parenchyma (cf. Fahn, 1959) and compared with those from a 17-year-old tree planted in the Ilanot Arboretum in the central Coastal Plain. The average number of wood parenchyma bands formed per year at Ilanot was found to be 2.35. The application of this ratio to the Tel es Sultan material suggests an age of 108 years for a 22.7-cm-long core with 253 parenchyma bands, and an age of 87 years for a 14.4-cm-long core with 205 bands. There is, therefore,

¹ Contribution from the Agricultural Research Organization, Bet Dagan, Israel. 1974 Series, No. 115-E.


reason to believe that the tree at Tel es Sultan is much older than 100 years, as is also suggested by its size; thus, it cannot have grown from the "stunted bush" recorded by Hart in 1891.

The site of Tel es Sultan (210 m below sea level), at 31°52'N, represents the northernmost point of the general distribution of *A. raddiana* (Halevy & Orshan, 1972). There is little, if any evidence to show that in the recent past the species extended farther northward along the Jordan Rift Valley. This view is supported by the toponymy of the area, i.e., the complete absence of place names which can be associated with *A. raddiana* (Conder & Kitchener, 1883). There is, however, proof that even at the end of the 19th century, the tree was more frequent in the Jericho area than it is today. For instance, a woodland of *A. raddiana*, *Balanites aegyptiaca* (L.) Del. and *Ziziphus spina-christi* (L.) Willd. extended from Ain es Sultan to Jericho along the perennial stream discharging into Wadi Kelt (Conder & Kitchener, 1883); at present, the spring flow irrigates the intensively cultivated oasis.

No evidence is available so far on the actual occurrence of *A. albida* in Jericho. However, its presence in the Lower Jordan Valley, approximately 70 km south of several well-documented occurrences near Bet She'an (32°30'N), cannot be excluded since the species was apparently collected near St. John's Convent, about 8 km ESE of Jericho (coll. Kushnir, 1942), and in the environs of the Dead Sea (coll. Naftulski, 1924). These two collections were quoted by Halevy (1971) and also inspected by the present author at the Herbarium of the Department of Botany, Hebrew University, Jerusalem. If the determination of the herbarium sheets is indeed correct, these sites would represent the southernmost occurrences of *A. albida* in the Jordan Rift Valley. The Jericho area would then be the only area in the Middle East where the ranges of *A. raddiana* and *A. albida* overlap.

LITERATURE CITED

- Conder, C. R. & H. H. Kitchener (1883) *The survey of Western Palestine. Memoirs of the topography, orography, hydrography and archeology* 3. Palestine Exploration Fund, London.
- Fahn, A. (1959) Xylem structure and annual rhythm of development of trees and shrubs of the desert. II. *Acacia tortilis* and *A. raddiana*. *Bull. Res. Council Israel Sect. D* 7: 23-28.
- Halevy, G. (1971) A study of *Acacia albida* in Israel. *La-Yaarot* 21: 52-63 (Hebrew), 89-97 (English).
- & G. Orshan (1972) Ecological studies on *Acacia* species in the Negev and Sinai. I. Distribution of *Acacia raddiana*, *A. tortilis* and *A. gerrardii* spp. *negevensis* as related to environmental factors. *Israel J. Bot.* 21: 197-208.
- Hart, H. C. (1891) *Some account of the fauna and flora of Sinai, Petra and Wâdy 'Arabah*. London.
- Post, G. E. & J. E. Dinsmore (1932) *Flora of Syria, Palestine and Sinai* 1. 2nd ed. Beirut.



Acacia raddiana Savi. Tel es Sultan, Jericho, August 1967 (Photo Y. Zohar).