

Noronhia edentata

Objekttyp: **Chapter**

Zeitschrift: **Boissiera : mémoires de botanique systématique**

Band (Jahr): **70 (2016)**

PDF erstellt am: **06.06.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

27. *Noronhia edentata* (H. Perrier) Hong-Wa, **comb. & stat. nov.** (Fig. 18A).

≡ *Noronhia emarginata* var. *edentata* H. Perrier in Mém. Inst. Sci. Madagascar, Sér. B, Biol. Vég. 2: 296. 1949.

Lectotypus (designated here): **MADAGASCAR. Prov. Toamasina:** Forêt d'Analamazaotra, [18°55'48"S 48°25'48"E], 800 m, s.d., *Perrier de la Bâthie* 8809 (P [P00791237]!); isolecto-: K [K000233192, K000233193] image seen; P [P03558849, P03558850]!).

Syntypi: **MADAGASCAR. Prov. Toamasina:** Analamazaotra RS, 2.IV.1905, *Thouvenot* 58 (K [K000233191] image seen, P [P00701239, P00701240]!); *ibid. loc.*, comm. 3.XII.1934, *Ursch* 82 (P [P03558851]!).

Description

Trees to 25 m tall, trunk to 35 cm diameter; young twigs cylindrical, 1-6 mm diameter, glabrous; bark medium to light gray, smooth to rugose. *Leaves* opposite, persistent; bud scales persistent; blades dark green above, medium green below, elliptic to obovate, 5-17 × 2-7.5 cm, coriaceous, glabrous, domatia absent, base acute to attenuate, margin slightly revolute, apex rounded to slightly acuminate, the acumen 0-5(-15) mm long, midrib sunken above, raised below, secondary veins conspicuous, 7-17 per side, 7-25 mm apart, looping 1.5-7 mm from the margin; petiole medium gray, 6-38 × 1.3-4 mm, entirely woody, glabrous. *Thyrse*s fasciculate, pauciflorous, compact; peduncle 8-9 mm long, moderately pubescent; pedicel 1.5-8 mm long, moderately pubescent; calyx moderately pubescent outside, glabrous inside, lobes triangular, 1-1.5 × 0.8-2 mm; corolla white sometimes tinged greenish, urceolate, 3-5.5 mm long, glabrous on both sides, the tube 1.2-3 mm long, lobes widely ovate, apex obtuse; corona present, 1.4-2.2 mm long, undivided; stamens 1.8-2.5 mm long, anthers widely oblong, 1.1-1.9 mm long; pistil 1.4-2.5 mm long, stigma capitate. *Fruiting* pedicel 2-9 × 1.8-3.6 mm; young fruits green, reddish black when mature, ovoid to subglobose, 12-31 × 8.5-26 mm, surface smooth, sometimes covered with a white pellicle, apex flat to bluntly pointed or apiculate; dry pericarp 1.3-4.2 mm thick; endocarp woody; seed 11.5-21.5 × 7-18 mm.

Distribution, ecology and phenology

Noronhia edentata occurs in low- to high-elevation humid forests in the east, from Sambava to Ambondrombe (Fig. 19). It produces flowers and fruits from September to May.

Conservation status

The assessment was based on 27 collections representing 19 localities, resulting in an EOO of 38,642 km², an AOO of 68 km², and 13 subpopulations representing 12 locations, of which six occur within protected areas (Analamazaotra, Betampona, Makira, Makirovana, Masoala, and Zahamena). Being widespread and present in many protected areas, *Noronhia edentata* is assigned a preliminary status of "Least Concern".




Fig. 18. Photographs of *Noronhia* Stadtm. ex Thouars. **A.** *Noronhia edentata* (H. Perrier) Hong-Wa [Razanatzima 266]; **B.** *Noronhia crassiramosa* (H. Perrier) Hong-Wa [A. Randrianasolo s.n.].

Photos: F. Rakotoarivony




Fig. 19. Distribution maps of species of *Noronhia* Stadtm. ex Thouars. *N. edentata* (H. Perrier) Hong-Wa to *N. insularis* (Labat, M. Pignal & O. Pascal) Hong-Wa & Besnard.

Notes *Noronhia edentata* can be recognized by its leaves, which have broadly elliptic blades and long, woody petioles, its white greenish flowers and its large fruits, with thick pericarp. There is a slight leaf heteroblasty in this species. In particular, the leaf base is narrower and the acumen longer (up to 15 mm long) in seedlings and saplings than in adult plants. This taxon was previously recognized as a variety of *N. emarginata*, from which it differs by its habit, leaf shape, and flower color and shape. Moreover, these two entities belong to separate clades (HONG-WA & BESNARD, 2013, 2014).

PERRIER DE LA BÂTHIE (1949) described *N. emarginata* var. *edentata* based on the following syntypes: *Perrier de la Bâthie* 8809, *Thouvenot* 58 and *Ursch* 820; the latter is a typographical mistake for *Ursch* 82. The overall quality and completeness of *Perrier de la Bâthie* 8809 justify the selection of one of the specimens at P as the lectotype.

Additional specimens examined

MADAGASCAR. **Prov. Antsiranana:** Ambatobiribiry, [14°11'S 50°05'E], 50-345 m, 29-30.XI.1950, *Humbert & Capuron* 24423 (G, MO, P). **Prov. Toamasina:** Maroantsetra, Ambinanitelo, Ankirindro forest, 15°17'35"S 49°23'59"E, 664 m, 16.I.2003, *Antilahimena et al.* 1741 (G, MO, P); Manambolo, Amparihibe forest, 15°02'26"S 49°34'59"E, 800 m, 22.II.2003, *Antilahimena* 1919 (MO); Moramanga, Sahaevo, 18°50'38"S 48°16'54"E, 1006 m, 11.XI.2006, *Antilahimena & Edmond* 4944 (G, MO, P, TAN, TEF); Ambatovy, 18°51'43"S 48°17'39"E, 1049 m, 9.XII.2006, *Antilahimena & Edmond* 5092 (MO, P, TAN, TEF); *ibid. loc.*, 18°50'28"S 48°18'16"E, 1085 m, 13.V.2010, *Bernard* 1566 (MO, P, TAN); Analamazaotra RS, 18°56'12"S 48°25'09"E, 953 m, 2.V.2010, *Hong-Wa & Ortiz* 650 (MO, P, TAN); Betampona RNI, 17°51'S 49°12'E, 275-650 m, 30.IX.1993, *Lewis* 711 (K, MO, P, TAN); Analamazaotra RS, [18°56'S 48°26'E], 25.VII, *Louvel* 88 (P); Andasibe, 18°55'50"S 48°25'04"E, 963 m, 5.XII.2008, *Rakotondrafara et al.* 932 (MO, P, TAN); Moango, Zahamena RNI, 17°33'32"S 48°53'55"E, 800 m, 4.V.2003, *Rakotondrajaona et al.* 274 (CNARP, G, MO, P, TEF); Lakato, Ambodigavo, forêt d'Analanjahana, 19°07'57"S 48°24'00"E, 794 m, 31.V.2007, *Razanatsima et al.* 266 (G, MO, P, TAN); 32 km E of Moramanga, 18°57'S 48°27'E, 870 m, 29.I.1993, *Schatz* 3452 (MO, P, TAN); Maroantsetra, Farankaraina, [15°25'S 49°52'E], 20-25 m, 16.IX.1952, *Service Forestier* 6151 (P, TEF); Périnet, [18°56'S 48°26'E], 1000 m, 1.VI.1954, *Service Forestier* 10374 (P, TEF); Masoala Peninsula, Ambanizana, 15°40'24"S 49°57'51"E, 110-260 m, 28.X.1994, *Vasey & Velo* 83 (MO, P, TAN).