

# **Peptidoglycan synthesis in eucaryotic cells and its possible roles**

Autor(en): **Roten, Claude-Alain H. / Karamata, Dimitri**

Objekttyp: **Article**

Zeitschrift: **Bulletin de la Société Vaudoise des Sciences Naturelles**

Band (Jahr): **82 (1992-1993)**

Heft 1

PDF erstellt am: **25.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-280168>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, [www.library.ethz.ch](http://www.library.ethz.ch)

## Peptidoglycan synthesis in eucaryotic cells and its possible roles

CLAUDE-ALAIN H. ROTEN<sup>1,2</sup> and DIMITRI KARAMATA<sup>1</sup>

*Résumé.*—ROTEN C.-A. H. et KARAMATA D., 1992. De la synthèse du peptidoglycane dans la cellule eukaryote et de ses rôles éventuels. *Bull. Soc. vaud. Sc. nat.* 82.1: 87-89.

Des évidences de présence du peptidoglycane, considéré comme un marqueur bactérien spécifique, ont été découvertes chez les plantes et chez les animaux. Nous discutons de la possible synthèse endogène du peptidoglycane qui expliquerait la présence de composés du peptidoglycane dans le cerveau, organe protégé par une barrière efficace contre le peptidoglycane exogène. De plus, la quantité de ces composés produite par la flore bactérienne de l'intestin grêle est insuffisante pour expliquer la quantité de ceux présents dans l'urine. Des fonctions possibles de ces composés sont discutées.

*Summary.*—ROTEN C.-A. H. et KARAMATA D., 1992. Peptidoglycan synthesis in eucaryotic cells and its possible roles. *Bull. Soc. vaud. Sc. nat.* 82.1: 87-89.

Evidence of peptidoglycan, believed to be a specific bacterial marker, is found in plants and animals. We discuss a possible endogenous eucaryotic peptidoglycan synthesis which would explain peptidoglycan components found in brain, where an efficient barrier against exogenous peptidoglycan exists. Moreover, the amounts of such compounds produced by bacterial flora of the small bowel are insufficient to account for those present in urine. Possible functions of these components are discussed.

**Key words:** diaminopimelate, muramate, muramyl-peptide, peptidoglycan in eucaryotes, synthesis of peptidoglycan, sleep muropeptide, endosymbiosis.

---

<sup>1</sup>Institut de Génétique et de Biologie Microbiennes, rue César-Roux 19, CH-1005 Lausanne, Suisse.

<sup>2</sup>From October 1992, at the Boston Biomedical Research Institute, 20, Staniford Street, Boston, Massachusetts 02114, USA.

A fundamental taxonomic criterion defining the bacterial kingdom is the presence of peptidoglycan (PG), the cell wall component and a specific target for antibiotics. Frequently, the basic unit of this polymer is N-acetyl-glucosaminyl-N-acetyl-muramyl-L-alanyl-D-glutamyl-L,D-diaminopimelyl-D-alanine, and thus muramate, as well as L,D-diaminopimelate (DAP) are bacteria-specific taxonomic markers. Although mitochondria and chloroplasts are of bacterial origin (GRAY *et al.* 1989), PG and its components are not believed to be synthesized by eucaryotic cells.

However, several observations on higher organisms call this belief into question. First, DAP containing compounds are excreted in constant daily amounts in human, bovine and swine urine (10, 30 and 70  $\mu\text{g} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$ , respectively) (KRYSCIAK 1980). Second, muramate was reproducibly found in rat brain (50-100 picomole  $\cdot \text{g}^{-1}$ ), and liver (100-150 picomole  $\cdot \text{g}^{-1}$ ) (SEN and KARNOVSKY 1984). Third, there is a dramatic effect of 1 picomole of urine sleep factor (FSu), the anhydro PG unit. This PG degradation product, extracted from animal urine or brain, induces a slow-wave sleep in rabbits when infused cerebro-intraventricularly (KRUEGER *et al.* 1984). The presence of these compounds in animal cells and urine was hitherto attributed to degradation products of the bacterial intestinal flora or to bacteria engulfed by macrophages (KRYSCIAK 1980, PAPPENHEIMER 1983, SEN and KARNOVSKY 1984).

These explanations are unsatisfactory, however, since the total amount of DAP containing bacteria in the small bowel ( $10^3$ - $10^7$  bacteria/ml) (GOLDIN 1986), where amino acid absorption occurs, is insufficient to account for the amount of DAP daily excreted in human urine. The latter being equivalent to the PG content of  $6 \cdot 10^{11}$  *E. coli* cells.

Quantitative analysis of muramate contained in rat hepatocytes strongly suggests that muramate is synthesized in a pulse, whose amount is equal to that present in the PG of *E. coli* cells whose total volume is equivalent to the volume of mitochondria present in dividing hepatocytes only. Thus, PG derived compounds, mitogenic for lymphocytes (DZIARSKI 1989), could act as division signals. Since cerebrospinal fluid is protected by a highly specific barrier, presence of exogenous PG compounds in the brain is most unlikely. The very strong effects of FSu (somnogenic and pyrogenic like interleukine 1) suggest that this molecule is part of a biological clock; like in septating bacteria, PG would be synthesized as a pulse (LLEO *et al.* 1990).

In conclusion, we postulate that eucaryotic cells are capable of endogenous PG synthesis; the latter being a trigger for cell division or, in mammals, sleep.

Presence of the PG biosynthetic pathway or parts of it in eucaryotic cells is not surprising. For instance, unlike certain fungi, green plants synthesize lysine in chloroplasts, via a specific bacterial DAP pathway (VOGEL 1965, WALLSGROVE *et al.* 1983), while the cyanelle of *Cyanophora paradoxa*, an organelle related to the chloroplast (URBACH. *et al.* 1992), is surrounded by a PG containing cell wall (KIES and KREMER 1990).

Appearance of the PG biosynthetic pathway in higher organisms could be accounted for by the serial endosymbiosis theory (MARGULIS 1981).

A more complete discussion of a possible endogenous eucaryotic PG will be published in its entirety in a manuscript in review (ROten and KARAMATA).

## REFERENCES

- DZIARSKI R., 1989. Correlation between ribosylation of pertussis toxin substrates and inhibition of peptidoglycan-, muramyl dipeptide- and lipopolysaccharide-induced mitogenic stimulation in B lymphocytes. *Eur. J. Immunol.* 19: 125-130.
- GOLDIN B. R., 1986. *In situ* bacterial metabolism and colon mutagens. *Ann. Rev. Microbiol.* 40: 367-393.
- GRAY M. W., CEDERGREN R., ABEL Y. and SANKOFF D., 1989. On the evolutionary origin of the plant mitochondrion and its genome. *Proc. Natl. Acad. Sci. USA* 86: 2267-2271.
- KIES L. and KREMER B. P., 1990. Phylum Glauco-cystophyta. In: L. MARGULIS, J. O CORLISS, M. MELKONIAN and D. J. CHAPMAN (eds) *Handbook of Protocista*, p. 152-166. Jones and Barlett Publishers, Boston.
- KRUEGER J. M., KARNOVSKY M. L., MARTIN S. A., PAPPENHEIMER J. R., WALTER J. and BIEMANN K., 1984. Peptidoglycans as promoters of slow-wave sleep. II Somnogenic and pyrogenic activities of some naturally occurring muramyl-peptides; correlations with mass spectrometric structure determination. *J. Biol. Chem.* 259: 12659-12662.
- KRYSCIAK J., 1980. Diaminopimelate in mammalian urine. *Folia Biol. (Kraków)* 28: 47-51.
- LLEO M. M., CANEPARI P. and SATTA G., 1990. Bacterial cell shape regulation: testing of additional predictions unique to the two-competing-sites model for peptidoglycan assembly and isolation of conditional rod-shaped mutants from some wild-type cocci. *J. Bact.* 172: 3758-3771.
- MARGULIS L., 1981. *Symbiosis in Cell Evolution*. Freeman, W. H. and Company, San Francisco.
- PAPPENHEIMER J. R., 1983. Induction of sleep by muramyl peptides. *J. Physiol.* 336: 1-11.
- SEN Z. and KARNOVSKY M. L., 1984. Qualitative detection of muramic acid in normal mammalian tissues. *Infect. Immun.* 43: 937-941.
- URBACH E., ROBERTSON D. L. and CHISHOLM S. W., 1992. Multiple evolutionary origins of prochlorophytes within the cyanobacterial radiation. *Nature* 355: 267-270.
- VOGEL H. J., 1965. Lysine biosynthesis and evolution. In: V. BRYSON and H. J. VOGEL (eds), *Evolving Genes and Proteins*, p. 25-40. Academic Press Inc., New York and London.
- WALLSGROVE R. M., LEA P. J. and MIFLIN B. J., 1983. Intracellular localization of aspartate kinase and the enzymes of threonine and methionine biosynthesis in green leaves. *Plant Physiol.* 71: 780-784.

*Manuscrit reçu le 5 juin 1992*

