

On mappings into group-like spaces.

Autor(en): **Whitehead, George W.**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **28 (1954)**

PDF erstellt am: **27.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-22627>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

On mappings into group-like spaces

by GEORGE W. WHITEHEAD

Dedicated to H. Hopf on his 60th birthday

1. Introduction. Let G be a space with continuous multiplication and inversion, which satisfies the group axioms up to homotopy. Then the homotopy classes of maps of any space X into G form a group $\pi(X; G)$. We shall first show that, under reasonable hypotheses on X , G , the group $\pi(X; G)$ is nilpotent; an upper bound for the class of nilpotency is $c - 1$, where c is the Lusternik-Schnirelmann category of X .

In particular, if X is the product of k spheres, $\pi(X; G)$ has class $\leq k$; an explicit central chain for $\pi(X; G)$ can be constructed in this case; the successive factor groups are direct products of homotopy groups of G . In particular, $\pi(S^p \times S^q; G)$ is a central extension of $\pi_{p+q}(G)$ by $\pi_p(G) \times \pi_q(G)$, which is semi-split in the sense that each of the groups $\pi_p(G)$, $\pi_q(G)$ can be lifted to a subgroup of $\pi(S^p \times S^q; G)$. The group extension is then completely described by the commutators of elements of $\pi_p(G)$ with elements of $\pi_q(G)$. These provide a bilinear map $(\alpha, \beta) \rightarrow \langle \alpha, \beta \rangle$ of $\pi_p(G) \times \pi_q(G)$ into $\pi_{p+q}(G)$. This map has been used by Samelson [3].

In the group $\pi(S^p \times S^q \times S^r; G)$ the iterated commutators provide a trilinear map $(\alpha, \beta, \gamma) \rightarrow \langle \alpha, \langle \beta, \gamma \rangle \rangle$ of the product

$$\pi_p(G) \times \pi_q(G) \times \pi_r(G)$$

into $\pi_{p+q+r}(G)$. Now if Γ is a group, the iterated commutators satisfy a "Jacobi congruence" modulo the fourth member of the descending central series of Γ ; since $\pi(S^p \times S^q \times S^r; G)$ is nilpotent of class ≤ 3 , this congruence reduces to an identity. From this fact we deduce a Jacobi identity for the operation $\langle \alpha, \beta \rangle$.

Suppose that G is the space of loops in a space X ; then

$$\pi_n(G) \approx \pi_{n+1}(X) .$$

By this isomorphism the commutator mapping is transformed, except

for sign, into the Whitehead product. Thus the above results provide another proof that the Whitehead products satisfy a Jacoby identity¹⁾.

2. Nilpotency of $\pi(X; G)$. Let X be a separable metric locally contractible space. A subset A of X is said to be *categorical* if and only if there is an open set $U \supset A$ such that U is contractible in X . A covering of X by sets A_α is said to be categorical if and only if each of the sets A_α is categorical. The *category* $\text{cat } X$ of X is the least of the cardinal numbers of the categorical coverings of X .

Let G be a space, $e \in G$, and let $\mu: (G \times G, (e, e)) \rightarrow (G, e)$, $i: (G, e) \rightarrow (G, e)$ be maps. We say that G is an H^* -space with respect to e, μ, i if and only if the following conditions are satisfied:

2.1) The maps $(x, y, z) \rightarrow \mu(x, \mu(y, z))$ and $(x, y, z) \rightarrow \mu(\mu(x, y), z)$ of $G \times G \times G$ into G are homotopic rel. (e, e, e) ;

2.2) The maps $x \rightarrow \mu(x, e)$ and $x \rightarrow \mu(e, x)$ of G into G are homotopic rel. e to the identity map;

2.3) The maps $x \rightarrow \mu(x, i(x))$ and $x \rightarrow \mu(i(x), x)$ of G into G are homotopic rel. e to the constant map.

Hereafter we abbreviate $\mu(x, y)$ to $x \cdot y$ and $i(x)$ to x^{-1} .

Let $\Phi: G \times G \rightarrow G$ be the commutator map, defined by

$$\Phi(x, y) = (xy)(x^{-1}y^{-1}),$$

and let $G \vee G$ be the subset $G \times e \cup e \times G$ of $G \times G$. Then from 2.1 to 2.3 we conclude easily:

2.4) The map $\Phi|G \vee G$ is homotopic rel. (e, e) to the constant map.

Let X be a space, $A \subset X$, and let G be an H^* -space. If $f, g: (X, A) \rightarrow (G, e)$ are maps, we define $f \cdot g: (X, A) \rightarrow (G, e)$ by

$$(f \cdot g)(x) = f(x) \cdot g(x).$$

This operation preserves the relation of homotopy, and therefore induces an operation in the set $\pi(X, A; G)$ of homotopy classes of maps of (X, A) into (G, e) . It is easily verified that $\pi(X, A; G)$ is a group under this operation. Moreover, if $h: (Y, B) \rightarrow (X, A)$ is a map, then composition with h induces a homomorphism

$$h: \pi(X, A; G) \rightarrow \pi(Y, B; G);$$

¹⁾ Proofs of the Jacobi identity for the Whitehead product have been obtained recently by Hilton and by Massey-Uehara (to appear in Ann. of Math.). The author believes the present proof to be more elementary. The author is informed by Uehara that proofs have also been found by Nakaoka and by Toda.

the homomorphism induced by a composite map $k \circ h$ is the composition $h \circ k$ in the opposite order.

We further have :

2.5) If S is an n -sphere, $x \in S$, then $\pi(S, x; G) = \pi_n(G)$.

2.6) Let (X', a) be the pair obtained from (X, A) by collapsing A to a point, $j: (X, A) \rightarrow (X', a)$ the identification map. Then $j: \pi(X', a; G) \approx \pi(X, A; G)$.

The proof of 2.5 is similar to the proof in the case where G is a topological group [1]; 2.6 follows from standard properties of identification spaces.

Lemma 2.7. Let $(X; A_1, \dots, A_n)$ be an

$$(n+1) - \text{ad} \ , \quad A = A_1 \cup \dots \cup A_n \ .$$

Suppose that there are retractions $\varrho_i: X \rightarrow A_i$ ($i = 1, \dots, n$) such that $\varrho_i(A_j) \subset A_j$ for all i, j . Suppose further that either $(X; A_1, \dots, A_n)$ is triangulable or X is separable metric and G is an ANR^2 . Then, if $j: X \subset (X, A)$, the homomorphism $j: \pi(X, A; G) \rightarrow \pi(X; G)$ is an isomorphism into.

Proof. We must show that, if $f_0, f_1: X \rightarrow G$ are homotopic maps with $f_0(A) = f_1(A) = e$, then f_0 and f_1 are homotopic rel. A . Let $B_r = A_1 \cup \dots \cup A_r$; then it suffices to show that $f_0 \simeq f_1$ rel. B_r implies $f_0 \simeq f_1$ rel. B_{r+1} . Let $F: X \times I \rightarrow G$ be a homotopy of f_0 to f_1 rel. B_r , and define $F_1: X \times I \rightarrow G$ by

$$F_1(x, t) = F(x, t) \cdot F[\varrho_{r+1}(x), t]^{-1} \ .$$

Then $x \in B_r$ implies $\varrho_{r+1}(x) \in B_r$, and therefore $F_1(x, t) = e \cdot e^{-1} = e$; and $x \in A_{r+1}$ implies $\varrho_{r+1}(x) = x$. Thus $x \in B_{r+1}$ implies that $F_1(x, t) = F(x, t) \cdot F(x, t)^{-1}$. Furthermore, if $t = 0$ or 1 , $F_1(x, t) = f_t(x) \cdot e$.

It follows from 2.3 that $F_1|B_{r+1} \times I$ is homotopic rel $B_{r+1} \times \dot{I}$ to the constant map. It follows from 2.2 that $F_1|X \times \dot{I}$ is homotopic rel $A \times \dot{I}$ to the map $(x, t) \rightarrow f_t(x)$. Let $C = X \times \dot{I} \cup B_{r+1} \times I$; then $F_1|C$ is homotopic to the map $F_2: C \rightarrow G$ such that

$$\begin{aligned} F_2(x, t) &= f_t(x) & (x \in X, t \in \dot{I}) ; \\ F_2(x, t) &= e & (x \in B_{r+1}) \ . \end{aligned}$$

²) In the weak sense.

By the homotopy extension theorem, F_2 has an extension $F' : X \times I \rightarrow G$; F' is a homotopy of f_0 to f_1 rel. B_{r+1} .

Let Γ be a group. We recall that the *descending central series* of Γ is the sequence of subgroups defined inductively by

$$Z_0 = \Gamma; \quad Z_i = [\Gamma, Z_{i-1}] \quad \text{for } i > 0,$$

and that Γ is *nilpotent* if and only if there is a c such that $Z_c = \{1\}$; the least such c is called the *class* of Γ . A *central chain* of length k of Γ is a sequence

$$\Gamma = \Gamma_0 \supset \dots \supset \Gamma_k = \{1\}$$

of subgroups such that $[\Gamma, \Gamma_i] \subset \Gamma_{i+1}$ ($i = 0, \dots, k-1$). Γ has a central chain of length k if and only if Γ is nilpotent of class $\leq k$.

We further recall [6, pp. 60—63]:

$$2.8) \quad [a, bc] \equiv [a, b] \cdot [a, c] \pmod{Z_2},$$

$$2.9) \quad [a, [b, c]] \cdot [b, [c, a]] \cdot [c, [a, b]] \equiv 1 \pmod{Z_3}.$$

Theorem 2.10. *Let X be a separable metric locally contractible space, G a 0-connected H^* -space. Suppose that X has finite category c , and that either a) X is triangulable or b) G is an ANR. Then the group $\pi(X; G)$ is nilpotent of class $\leq c-1$.*

To prove Theorem 2.10, we recall from [2, p. 336] that X has a closed categorical covering A_1, \dots, A_c . If X is triangulable we may assume that the sets A_i are all subcomplexes of a certain simplicial decomposition of X . Theorem 2.10 will then follow from

Lemma 2.11. *Let Γ'_i be the set of all homotopy classes of maps $f : X \rightarrow G$ such that $f|_{A_1 \cup \dots \cup A_{i+1}}$ is inessential ($i = 0, \dots, c-1$). Then the Γ'_i form a central chain for $\pi(X; G)$.*

Proof. Clearly the Γ'_i form a decreasing sequence of subgroups of $\Gamma = \pi(X; G)$; $\Gamma'_0 = \Gamma$ since A_1 is contractible in X , and $\Gamma'_{c-1} = \{1\}$ since $A_1 \cup \dots \cup A_c = X$. It remains to prove that $[\Gamma, \Gamma'_{i-1}] \subset \Gamma'_i$.

Let $f \in \alpha \in \Gamma$, $g \in \beta \in \Gamma'_{i-1}$; then $f|_{A_{i+1}}$ is inessential since A_{i+1} is contractible in X , and $g|_{A_1 \cup \dots \cup A_i}$ is inessential by hypothesis. Hence we may assume $f(A_{i+1}) = g(A_1 \cup \dots \cup A_i) = e$. Define $h : X \rightarrow G$ by $h = \Phi \circ (f \times g)$, i. e.,

$$h(x) = \Phi(f(x), g(x));$$

then $h \in [\alpha, \beta]$. Now if $x \in A_1 \cup \dots \cup A_{i+1}$, then $(f(x), g(x)) \in G \vee G$; it follows from 2.4 that $h|_{A_1 \cup \dots \cup A_{i+1}}$ is inessential, i. e., $[\alpha, \beta] \in \Gamma'_i$.

Corollary 2.12. *If X has finite dimension n , then $\pi(X; G)$ is nilpotent of class $\leq n$.*

For $\text{cat } X \leq 1 + \dim X$ [2, (5.4)].

Corollary 2.13. *If X is the product of k spheres, then $\pi(X; G)$ is nilpotent of class $\leq k$.*

For $\text{cat } X = k + 1$ [2, p. 350].

The following variant of Lemma 2.11 will be more appropriate in what follows³).

Lemma 2.14. *Let X be a CW-complex [5] and let*

$$A_0 \subset \cdots \subset A_k = X$$

be subcomplexes such that, if E is any cell of A_i , then $\dot{E} \subset A_{i-1}$. Let Γ_i be the set of all homotopy classes of maps $f: X \rightarrow G$ such that $f|A_i$ is inessential. Then $\Gamma_0, \dots, \Gamma_k$ is a central chain for $\pi(X; G)$.

Proof. Let $f \in \alpha \in \Gamma = \pi(X; G)$, $g \in \beta \in \Gamma_{i-1}$. For each cell E_λ of A_i , let x_λ be an interior point of E_λ . Then we can find closed cells $F_\lambda \subset \text{Int } E_\lambda$ such that $x_\lambda \in \text{Int } F_\lambda$ and \dot{E}_λ is a deformation retract of $\overline{E_\lambda - F_\lambda}$; it follows that A_{i-1} is a deformation retract of

$$Q = \bigcup_\lambda \overline{E_\lambda - F_\lambda} \cup A_{i-1}$$

and the set $P_0 = \{x_\lambda\}$ is a deformation retract of $P = \bigcup_\lambda F_\lambda$. Since G is 0-connected, $f|P_0$, and therefore also $f|P$, is inessential. Since $g|A_{i-1}$ is inessential, it follows that $g|Q$ is inessential. Hence we may assume $f(P) = g(Q) = e$. The remainder of the proof follows the pattern of Lemma 2.11.

3. The case $X = S^{n_1} \times \cdots \times S^{n_k}$. We now examine in more detail the case $X = S^{n_1} \times \cdots \times S^{n_k}$. We first make some conventions about orientation. Let I^p be the unit cube in Euclidean p -space; we orient I^p by choosing the generator ω_p of $H_p(I^p, \dot{I}^p)$ represented by the identity map of I^p (we are using cubical singular homology theory as in [4]). Let x_p be a fixed point of S^p and choose a fixed map $\psi_p: (I^p, \dot{I}^p) \rightarrow (S^p, x_p)$

³) The hypotheses of Lemma 2.14 imply that $\text{cat } X \leq k + 1$, so that $\pi(X; G)$ is nilpotent of class $\leq k$. The conclusion follows formally from Lemma 2.11 and the following statement, which is easily proved by induction on k : *there are closed categorical subsets B_0, \dots, B_k of X such A_i is a deformation retract of $B_0 \cup \dots \cup B_i$ ($i = 0, \dots, k$).* However, it seemed simpler to prove Lemma 2.14 directly. Actually, only Lemma 2.14 is used in what follows; we state Theorem 2.10 because it seems to give the most natural upper bound for the class of nilpotency.

which maps the interior of I^p topologically onto $S^p - x_p$; we then orient S^p by requiring that ψ_p have degree 1. Next we make the natural identification of $I^{n_1} \times \dots \times I^{n_k}$ with $I^{n_1 + \dots + n_k}$ and then orient $S^{n_1} \times \dots \times S^{n_k}$ by requiring that $\psi = \psi_{n_1} \times \dots \times \psi_{n_k}$ have degree 1.

Let $X = S^{n_1} \times \dots \times S^{n_k}$. We take $S_i = S^{n_i}$ with the CW -decomposition consisting of the 0-cell $e_i = x_{n_i}$ and the n_i -cell $S_i - e_i$. The product of these CW -decompositions gives a CW -decomposition of X . Let A_i be the set of all $x \in X$ such that $x_j \neq e_j$ for at most i values of j ; thus

$$\begin{aligned} A_0 &= e_1 \times \dots \times e_k, \\ A_1 &= S_1 \times e_2 \times \dots \times e_k \cup \dots \cup e_1 \times \dots \times e_{k-1} \times S_k, \\ &\dots \\ A_k &= S_1 \times \dots \times S_k = X. \end{aligned}$$

The A_i clearly satisfy the hypotheses of Lemma 2.14.

For each subset α of $\{1, \dots, k\}$, let $S_\alpha = \{x \in X \mid x_i = e_i \text{ for } i \notin \alpha\}$, $|\alpha|$ = the cardinal of α , $n(\alpha) = \sum_{i \in \alpha} n_i$. S_α is homeomorphic with $\prod_{i \in \alpha} S_i$, and we orient S_α by requiring that the natural homeo-

morphism (the one preserving the order of the coordinates) be orientation-preserving. Finally, let $p_\alpha: (S_\alpha, S_\alpha \cap A_{|\alpha|-1}) \rightarrow (S^{n(\alpha)}, e_{n(\alpha)})$ be an orientation-preserving map which is topological on $S_\alpha - A_{|\alpha|-1}$.

Let $q_\alpha: X \rightarrow S_\alpha$ be the natural retraction: $q_\alpha(x)$ is the point y such that $y_i = x_i$ if $i \in \alpha$ and $y_i = e_i$ if $i \notin \alpha$. Then $\beta \neq \alpha$ implies $q_\beta(S_\alpha) \subset S_\alpha$. We may therefore apply Lemma 2.7 to obtain:

Lemma 3.1. *Let $j: X \subset (X, A_i)$. Then $j: \pi(X, A_i; G) \approx \Gamma_i$.*

Theorem 3.2. *The group Γ_{i-1}/Γ_i is isomorphic with the direct product*

$$\prod_{\alpha} \pi_{n(\alpha)}(G),$$

where α ranges over all i -element subsets of $\{1, \dots, k\}$.

Proof. Let $j_\alpha: (S_\alpha, S_\alpha \cap A_{i-1}) \subset (X, A_{i-1})$. Then

$$j: \pi(X, A_{i-1}; G) \approx \Gamma_{i-1}, j_\alpha: \pi(X, A_{i-1}; G) \rightarrow \pi(S_\alpha, S_\alpha \cap A_{i-1}; G),$$

and $p_\alpha: \pi_{n(\alpha)}(G) \approx \pi(S_\alpha, S_\alpha \cap A_{i-1}; G)$; hence we may define $\eta_\alpha: \Gamma_{i-1} \rightarrow \pi_{n(\alpha)}(G)$ by $\eta_\alpha = p_\alpha^{-1} j_\alpha j^{-1}$. The homomorphisms η_α then define a homomorphism $\eta: \Gamma_{i-1} \rightarrow \Delta = \prod_{\alpha} \pi_{n(\alpha)}(G)$. Clearly $\Gamma_i \subset$

Kernel η . Conversely, if $f \in \gamma \in \text{Kernel } \eta$, then, for each α , $f|S_\alpha$ is

inessential rel. $S_\alpha \cap A_{i-1}$; hence $f|A_i$ is inessential. Therefore η induces an isomorphism η of Γ_{i-1}/Γ_i into Δ .

To show that η is onto, define $i_\alpha: X \rightarrow S^{n(\alpha)}$ by $i_\alpha = p_\alpha \circ q_\alpha$. Since q_α can be factored through (X, A_{i-1}) , it follows that $i_\alpha: \pi_{n(\alpha)}(G) \rightarrow \Gamma_{i-1}$. Now

$$\eta_\alpha \circ i_\alpha = p_\alpha^{-1} \circ j_\alpha \circ j^{-1} \circ q_\alpha \circ p_\alpha ;$$

since q_α is a retraction, $j_\alpha \circ j^{-1} \circ q_\alpha$ is the identity. Hence $\eta_\alpha \circ i_\alpha$ is the identity. If $\alpha \neq \beta$, then $q_\alpha(S_\beta) \subset S_\alpha \cap S_\beta \subset A_{i-1}$ and therefore $j_\beta \circ j^{-1} \circ q_\alpha = 0$; thus

$$\eta_\beta \circ i_\alpha = p_\beta^{-1} \circ j_\beta \circ j^{-1} \circ q_\alpha \circ p_\alpha = 0 .$$

It follows that η is onto, and that the i_α are isomorphisms into.

4. Commutator relations. In this section we write π_{n_1, \dots, n_k} for $\pi(S^{n_1} \times \dots \times S^{n_k}; G)$. Let $i_1: \pi_p \rightarrow \pi_{p,q}$, $i_2: \pi_q \rightarrow \pi_{p,q}$, $i_{1,2}: \pi_{p+q} \rightarrow \pi_{p,q}$ be the isomorphisms into of § 3. Then $i_{1,2}$ maps π_{p+q} isomorphically onto Γ_1 , and if $\alpha \in \pi_p$, $\beta \in \pi_q$, we have $[i_1(\alpha), i_2(\beta)] \in \Gamma_1$. Hence we may define

$$\langle \alpha, \beta \rangle = i_{1,2}^{-1}[i_1(\alpha), i_2(\beta)] \in \pi_{p+q} .$$

Lemma 4.1. (a) $\langle \alpha, \beta \rangle = (-1)^{p^q-1} \langle \beta, \alpha \rangle$; (b) $\langle \alpha, \beta \rangle$ depends bilinearly on α, β .

Proof. Let $t: S^p \times S^q \rightarrow S^q \times S^p$ be the map which interchanges the coordinates. In virtue of our conventions about orientation, t has degree $(-1)^{p^q}$. Furthermore, t induces a map $t': S^{p+q} \rightarrow S^{p+q}$ of the same degree. We verify easily that $i_1 \circ t = t_2$, $i_2 \circ t = i_1$, and $i_{1,2} \circ t = t' \circ i_{1,2}$. Furthermore $t'(u) = (-1)^{p^q} u$ for $u \in \pi_{p+q}$. Hence

$$\begin{aligned} \langle \beta, \alpha \rangle &= i_{1,2}^{-1}[i_1(\beta), i_2(\alpha)] = t'^{-1} i_{1,2}^{-1} t[i_1(\beta), i_2(\alpha)] \\ &= (-1)^{p^q} i_{1,2}^{-1}[ti_1(\beta), ti_2(\alpha)] \\ &= (-1)^{p^q} i_{1,2}^{-1}[i_2(\beta), i_1(\alpha)] \\ &= (-1)^{p^q-1} i_{1,2}^{-1}[i_1(\alpha), i_2(\beta)] \\ &= (-1)^{p^q-1} \langle \alpha, \beta \rangle . \end{aligned}$$

Now if $\alpha \in \pi_p$, $\beta, \gamma \in \pi_q$, we have, by 2.8,

$$\begin{aligned} [i_1(\alpha), i_2(\beta \cdot \gamma)] &= [i_1(\alpha), i_2(\beta) i_2(\gamma)] \\ &\equiv [i_1(\alpha), i_2(\beta)] \cdot [i_1(\alpha), i_2(\gamma)] \pmod{\Gamma_2} . \end{aligned}$$

But $\Gamma_2 = \{1\}$ and $i_{1,2}$ is a homomorphism. Hence right linearity holds. Left linearity follows from right linearity and part (a).

Consider now the group $\pi_{p,q,r}$ and the isomorphisms into

$i_1: \pi_p \rightarrow \pi_{p,q,r}$, $i_2: \pi_q \rightarrow \pi_{p,q,r}$, $i_3: \pi_r \rightarrow \pi_{p,q,r}$, and $i_{1,2,3}: \pi_{p+q+r} \rightarrow \pi_{p,q,r}$. The latter is an isomorphism onto Γ_2 . If $\alpha \in \pi_p$, $\beta \in \pi_q$, $\gamma \in \pi_r$, then $[i_1(\alpha), [i_2(\beta), i_3(\gamma)]] \in \Gamma_2$, and we have:

Lemma 4.2. $\langle \alpha, \langle \beta, \gamma \rangle \rangle = i_{1,2,3}^{-1} [i_1(\alpha), [i_2(\beta), i_3(\gamma)]]$.

Proof. We have, by definition,

$$\langle \alpha, \langle \beta, \gamma \rangle \rangle = i_{1,2}^{-1} [i_1(\alpha), i_2 i_{1,2}^{-1} [i_1(\beta), i_2(\gamma)]] .$$

Let $f: S^p \times S^q \times S^r \rightarrow S^p \times S^{q+r}$ be the map given by

$$f(x, y, z) = (x, p_{1,2}(y, z)) .$$

The maps $p_{1,2} \circ f$ and $p_{1,2,3}$ of $(S^p \times S^q \times S^r, A_2)$ into (S^{p+q+r}, x_{p+q+r}) both preserve orientation. Hence

$$f \circ i_{1,2} = f \circ p_{1,2} = p_{1,2,3} = i_{1,2,3} .$$

On the other hand, $i_2 \circ f = i_{2,3}$ and $i_1 \circ f = i_1$. Let h be the projection of $S^p \times S^q \times S^r$ on $S^q \times S^r$. Then $i_{1,2} \circ h = i_{2,3}$, $i_1 \circ h = i_2$, and $i_2 \circ h = i_3$. Thus

$$\begin{aligned} i_{1,2,3} \langle \alpha, \langle \beta, \gamma \rangle \rangle &= f i_{1,2} \langle \alpha, \langle \beta, \gamma \rangle \rangle \\ &= f [i_1(\alpha), i_2 i_{1,2}^{-1} [i_1(\beta), i_2(\gamma)]] \\ &= [f i_1(\alpha), f i_2 i_{1,2}^{-1} [i_1(\beta), i_2(\gamma)]] \\ &= [i_1(\alpha), i_{2,3} i_{1,2}^{-1} [i_1(\beta), i_2(\gamma)]] \\ &= [i_1(\alpha), h [i_1(\beta), i_2(\gamma)]] \\ &= [i_1(\alpha), [h i_1(\beta), h i_2(\gamma)]] \\ &= [i_1(\alpha), [i_2(\beta), i_3(\gamma)]] . \end{aligned}$$

Lemma 4.3. If $\alpha \in \pi_p$, $\beta \in \pi_q$, $\gamma \in \pi_r$, then

$$\begin{aligned} i_{1,2,3}^{-1} [i_2(\beta), [i_3(\gamma), i_1(\alpha)]] &= (-1)^{p(q+r)} \langle \beta, \langle \gamma, \alpha \rangle \rangle ; \\ i_{1,2,3}^{-1} [i_3(\gamma), [i_1(\alpha), i_2(\beta)]] &= (-1)^{r(p+q)} \langle \gamma, \langle \alpha, \beta \rangle \rangle . \end{aligned}$$

Proof. Let $g: S^p \times S^q \times S^r \rightarrow S^q \times S^r \times S^p$ be the map given by $g(x, y, z) = (y, z, x)$. Then g induces $g': S^{p+q+r} \rightarrow S^{p+q+r}$; g and g' have the same degree $(-1)^{p(q+r)}$, and $i_{1,2,3} \circ g = g' \circ i_{1,2,3}$, $i_1 \circ g = i_2$, $i_2 \circ g = i_3$, $i_3 \circ g = i_1$. Hence

$$\begin{aligned} i_{1,2,3} \{(-1)^{p(q+r)} \langle \beta, \langle \gamma, \alpha \rangle \rangle\} &= g i_{1,2,3} \langle \beta, \langle \gamma, \alpha \rangle \rangle \\ &= g [i_1(\beta), [i_2(\gamma), i_3(\alpha)]] \\ &= [g i_1(\beta), [g i_2(\gamma), g i_3(\alpha)]] \\ &= [i_2(\beta), [i_3(\gamma), i_1(\alpha)]] . \end{aligned}$$

The other statement is proved similarly.

Theorem 4.4. *If $\alpha \in \pi_p$, $\beta \in \pi_q$, $\gamma \in \pi_r$, then*

$$\langle \alpha, \langle \beta, \gamma \rangle \rangle + (-1)^{p(q+r)} \langle \beta, \langle \gamma, \alpha \rangle \rangle + (-1)^{r(p+q)} \langle \gamma, \langle \alpha, \beta \rangle \rangle = 0 .$$

Proof. Let $\alpha' = i_1(\alpha)$, $\beta' = i_2(\beta)$, $\gamma' = i_3(\gamma)$. Since $\Gamma_3 = \{1\}$, we have, by 2.9,

$$1 = [\alpha', [\beta', \gamma']] \cdot [\beta', [\gamma', \alpha']] \cdot [\gamma', [\alpha', \beta']]$$

and therefore

$$\begin{aligned} 0 &= i_{1,2,3}^{-1}[\alpha', [\beta', \gamma']] \cdot [\beta', [\gamma', \alpha']] \cdot [\gamma', [\alpha', \beta']] \\ &= \langle \alpha, \langle \beta, \gamma \rangle \rangle + (-1)^{p(q+r)} \langle \beta, \langle \gamma, \alpha \rangle \rangle + (-1)^{r(p+q)} \langle \gamma, \langle \alpha, \beta \rangle \rangle . \end{aligned}$$

Now suppose that X is a space, $x \in X$, and that G is the path-component of the constant map in the space of loops in X based at x . Then we have a natural isomorphism $T: \pi_{n+1}(X) \approx \pi_n(G)$ for each $n > 0$. Furthermore, if $\alpha \in \pi_{p+1}(X)$, $\beta \in \pi_{q+1}(X)$, and if $[\alpha, \beta] \in \pi_{p+q+1}(X)$ is their Whitehead product, we have [3]:

$$T([\alpha, \beta]) = (-1)^p \langle T(\alpha), T(\beta) \rangle . \quad (4.5)$$

From 4.5 and Theorem 4.4 we then obtain:

Theorem 4.6. *If $\alpha \in \pi_{p+1}(X)$, $\beta \in \pi_{q+1}(X)$, $\gamma \in \pi_{r+1}(X)$, then*

$$(-1)^{r(p+1)} [\alpha, [\beta, \gamma]] + (-1)^{p(q+1)} [\beta, [\gamma, \alpha]] + (-1)^{q(r+1)} [\gamma, [\alpha, \beta]] = 0 .$$

BIBLIOGRAPHY

- [1] *B. Eckmann*, Über die Homotopiegruppen von Gruppenräumen, *Comment. Math. Helv.* 14 (1941) 234–256.
- [2] *R. H. Fox*, On the Lusternik-Schnirelmann category, *Ann. Math.* 42 (1941) 333–370.
- [3] *H. Samelson*, A connection between the Whitehead and the Pontryagin product, *Amer. Journ. Math.* 75 (1953) 744–52.
- [4] *J.-P. Serre*, Homologie singulière des espaces fibrés, *Ann Math.* 54 (1951) 425–505.
- [5] *J. H. C. Whitehead*, Combinatorial homotopy I, *Bull. Amer. Math. Soc.* 55 (1949) 213–45.
- [6] *H. Zassenhaus*, The theory of groups, Chelsea 1949.

(Received February 18, 1954.)