Zeitschrift:	Commentarii Mathematici Helvetici
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	41 (1966-1967)
Artikel:	Actions of Rn on manifolds.
Autor:	Rosenberg, Harold
DOI:	https://doi.org/10.5169/seals-31377

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 03.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Actions of *Rⁿ* on manifolds

by HAROLD ROSENBERG

We shall be concerned with smooth manifolds V^n , compact and without boundary, and actions of R^{n-1} on V all of whose orbits are n-1 dimensional. The rank of V is the largest k such that there is an action of R^k on V with k dimensional orbits; this is the same as the maximal number of linearly independent vector fields on V which pairwise commute. Elon Lima has proved the rank of S^3 is one [1], and the author proved the rank of $S^2 \times S^1$ is one [4]. One of our results is a generalization of Lima's theorem: the rank of a simply connected closed n manifold is less than n-1. Unfortunately, the author knows of no n-dimensional sphere whose rank is greater than one.

We also consider $M \times S^1$ where M is a closed two-dimensional manifold of genus greater than one. Our results are not complete; we do not know the rank of this space. We do prove, however, that if there is a locally free action of R^2 on $M \times S^1$, then it must have a torus orbit, embedded in a nontrivial way.¹)

Definitions and Notation

An action Φ of a Lie Group G on V is a differentiable map $\Phi: G \times V \to V$ such that (i) $\Phi(gh, x) = \Phi(g, \Phi(h, x))$ for all g, $h \in G$ and $x \in V$, and (ii) $\Phi(e, x) = x$ for $x \in V$, e the identity of G. Given $x \in V$, the isotropy subgroup of x is $H_x = \{g \in G/\Phi_g(x) = x\}$, it is a closed subgroup of V. The orbit or leaf of x is $\{\Phi_g(x)/g \in G\}$. The action Φ induces a 1-1 continuous map of G/H_x onto L_x , the orbit of x.

If $X_1, ..., X_k$ are vector fields on V, we say they pairwise *commute* if $[X_i, X_j] \equiv 0$ for all i and j. Let V be a closed manifold and $\xi^1, ..., \xi^k$ the integral curves of $X_1, ..., X_k$ respectively. We know $[X_i, X_j] \equiv 0$ is equivalent to $\xi_s^i \xi_t^j = \xi_t^j \xi_s^i$ for all real numbers s and t.

When $G = R^{K}$, an action of G on V is equivalent to K commuting vector field (we assume V is closed); the relation is

$$\Phi(t,x) = (\xi_{t_1}^1 \circ \xi_{t_2}^2 \circ \cdots \circ \xi_{t_k}^k)(x), \quad t = (t_1, \dots, t_k) \in \mathbb{R}^k.$$

We call Φ a *locally free* action if all the orbits are K-dimensional.

Suppose n=3 and k=2. The orbits of x are classified by their isotropy subgroups H_x and we have the following possibilities. If the dimension of H_x is two, then $H_x = R^2$ and $L_x = X$. When H_x has dimension one we have $H_x = L + nv$, L a line through the origin and $v \in R^2$, $n=0, \pm 1, \pm 2, ..., L_x$ is then a line or circle (i.e., 1-1 continuous

¹⁾ Conversations with Elon Lima and André Haefliger were very useful in the preparation of this paper.

image of) depending on the direction of v. The case dimension $H_x=0$ gives three possible orbits. When $H_x=Z_u$, Z the group of integers, $u \in R^2$, we have $L_x=R^2$ or a cylinder depending on whether u=0 or $u\neq 0$. If $H_x=Zu+Zv$ with u and v independent, then H_x is a torus.

1. The Existence of Compact Leaves

THEOREM 1.1. (Reeb [2]). Let V be a closed Riemannian manifold and ω a closed one form on V satisfying $\|\omega\| = 1$. Let F be the foliation of V defined by $\omega = 0$. Then the leaves of F are homeomorphic and if L is one leaf, there is a covering map $p: R \times L \rightarrow V$.

Proof. Since $\|\omega\| = 1$, the foliation is oriented, and we may choose a unit vector field on V orthogonal to the foliation.

The orthogonal trajectories to a leaf F are geodesics [3]. Let $\Psi_s(x)$ be a parametrization by arc length of the orthogonal trajectory through x. For each x, there is a neighborhood U of x, where we may define a smooth function s(y) by s(y) = the distance of the point y from the leaf containing x. Our assumptions imply $\omega = ds$ locally.

If L is a leaf of F and s a real number, ω vanishes on $\Psi_s(L)$. Thus Ψ_s carries leaves into leaves. The set $\{\Psi_s(L)|s \in R\}$ is open and closed in V, hence all of V. This proves the first assertion.

Let $x_0 \in V$, and H be the subgroup of $\pi_1(V, x_0)$ of homotopy classes representable by closed curves h at x_0 such that

$$\int_{h} \omega = 0$$

Here we use the hypothesis $d\omega = 0$.

Let W be the connected covering space of V over H. On W we have the one form $\omega^* = p^* \omega$ and a foliation F_0 defined by $\omega^* = 0$. W inherits a Riemannian metric such that $\|\omega^*\| = 1$; ω^* is never zero, and $d\omega^* = 0$.

Let a be a closed curve in W based at some point in $p^{-1}(x_0)$. Since

$$\int_{a} \omega^* = \int_{pa} \omega$$

and pa represents an element of H, we have

$$\int_a \omega^* = 0.$$

It follows easily that the integral of ω^* about any closed curve in W is zero. Thus $\omega^* = df$ for some smooth function f on W. The level surfaces of f are precisely the leaves of F_0 . Each orthogonal trajectory to F_0 is an embedding of R in W and each

leaf meets an orthogonal trajectory in precisely one point. Hence W is homeomorphic to $R \times L_0$, where $L_0 \in F_0$, and for each t, $t \times L_0$ corresponds to a leaf of F_0 .

We observe that L_0 is homeomorphic to $p(L_0) = L$, a leaf of F. There is a map $L \rightarrow L_0$ defined as follows: fix $x_0 \in L$ and $\bar{x} \in L_0$ such that $px = x_0$. For $x \in L$, let h be a path in L from x_0 to x. Lift h to a path a in L_0 starting at \bar{x} . We map $L \rightarrow L_0$ by sending x to a(1), the endpoint of a. This map does not depend on the path h, since closed paths in L lift to closed paths in L_0 . Thus V may be covered by $R \times L$.

THEOREM 1.2 (Sacksteder [5]). Let Φ be a locally free action of \mathbb{R}^{n-1} on a closed n manifold V, such that no orbit is compact. There is a Riemannian metric on V and a closed non-vanishing one form ω of norm one, such that the foliation defined by $\omega = 0$ is the same as the foliation defined by Φ . This foliation admits a simple closed curve as an orthogonal trajectory.

COROLLARY 1.3: Let V be a closed n manifold with non-Abelian fundamental group. Then each locally free action of \mathbb{R}^{n-1} on V has a non-simply connected leaf.

Proof. Suppose the orbits of Φ are simply connected. Then theorems 1.1 and 1.2 imply V is covered by \mathbb{R}^n and $H = \{[a] \in \pi_1(V) | \int_a \omega = 0\}$ is isomorphic to $\pi_1(\mathbb{R}^n)$ hence trivial. But H contains the commutator subgroup of V, hence $\pi_1(V)$ is abelian.

COROLLARY 1.4: Let Φ be a locally free action of \mathbb{R}^2 on $M \times S^1$ where M is a closed 2-dimensional manifold of genus greater than one. Then Φ has a compact orbit (a torus).

Proof. Since $\pi_1(M \times S^1)$ is not abelian we know all the orbits of Φ cannot be R^2 . If Φ has no compact orbit, all of the orbits are the one to one continuous image of $R \times S^1$, and each orbit is dense in $M \times S^1$. Let X and Y be linearly independent commuting vector fields on $M \times S^1$ such that X and Y span the orbits of Φ . Let $x_0 \in V = M \times S^1$. The isotropy subgroup of R^2 at x_0 is a discrete group on one generator; hence, we may find real numbers a, b, c, d such that the vector fields X' = aX + bY, Y' = cX + dY are linearly independent and the X' orbit through x_0 is a simple closed curve y. Let ξt and η_t be the integral curves of X' and Y'. Because X' and Y' commute, we have $\xi_t \eta_\tau = \eta_\tau \xi_t$ for all t and τ . Thus $\eta_\tau(\gamma)$ is also a simple closed curve for all τ . Since the Φ orbit of x_0 is dense in V, it follows from continuity that all the integral curves of X' are simple closed curves. Moreover, the foliation of V induced by Φ may be assumed oriented which implies the integral curves of X' have the same period. Consider the quotient space Y of V obtained by identifying each integral curve of X'to a point. Y is a closed two-dimensional orientable manifold. By choosing a nonzero normal vector field to the orbits of Φ we obtain a non-zero vector field on Y; hence Y must be a two-dimensional torus. But this means $M \times S^1$ is a circle bundle over a two torus which is easily seen to be a contradiction. Simply consider the homotopy exact sequence of this fibre bundle. Thus some orbit of Φ is compact.

THEOREM 1.4. Let Φ be a locally free action of \mathbb{R}^{n-1} on a closed n manifold V and assume Φ has no compact orbits. There is a covering map $p: \mathbb{R}^{n-1} \times S^1 \to V$.

Proof. We may apply 1.2 to obtain a metric on V and closed non-vanishing one form ω of norm one which defines the foliation induced by Φ . Let $j: I \rightarrow V$ be a parametrization by arc length of the closed orthogonal trajectory through x_0 ; i.e., $j(0)=j(1)=x_0, j(t_1)\neq j(t_2)$ if $t_1\neq t_2, 0 < t_1, t_2 < 1$ and j(I) is orthogonal to Φ . It is no loss of generality to assume this orbit has length one.

Let L be the Φ orbit of x_0 . By 1.1 we know V is covered by $R \times L$. If L is not simply connected, then $L = R^{n-i} \times T^{i-1}$ where T^{i-1} is the i-1 dimensional torus and i>1. In this case $R \times L$ is covered by $R^{n-1} \times S^1$. So we may assume L is the one to one continuous image of R^{n-1} which implies each orbit of Φ is of the same type. We state in [4] that these assumptions imply V is covered by $R^{n-1} \times S^1$. Since this was stated without proof, we give the proof here.

Let *H* be the subgroup of $\pi_1(V, x_0)$ generated by the homotopy class of *j*. Let *W* be the connected covering space of *V* over *H* with covering map *p*. We will prove *W* is homeomorphic to $\mathbb{R}^{n-1} \times S^1$.

We may think of W as the quotient space of the space of paths $h: I \to V$ starting at x_0 where h_1 is identified with h_2 if $h_1(1) = h_2(1)$ and $h_1 h_2^{-1}$ represents an element of H. Parametrize j by arc length so that the distance of j(t) to x_0 is t.

Define a path $h(\tau)$ at x_0 by $h(\tau)(t) = j(t\tau)$, $0 \le \tau \le 1$. Let $U(\tau) = (h(\tau)) =$ equivalence class of $h(\tau)$ in W. We have U(0) = U(1) since h(1) = j, $h(0) = C_{x_0} =$ constant path at x_0 , and $h(1)h(0)^{-1} = j$ represents an element of H. Also $U(\tau_1) \ne U(\tau_2)$ for $\tau_1 \ne \tau_2$, $0 < \tau_2, \tau_2 < 1$, since $h(\tau_1) \ne h(\tau_2)$. Hence U is a simple closed curve in W such that p U = j.

Let Φ_0 be a lifting of the action Φ to an action on W; that is, $p\Phi_0 = \Phi(1 \times p)$, 1 = the identity map of \mathbb{R}^{n-1} . The orbits of Φ_0 cover the orbits of Φ hence they are also the one to one continuous image of \mathbb{R}^{n-1} . To complete the proof we will show each orbit of Φ_0 intersects the image of U is pre precisely one point.

Suppose some orbits A of Φ_0 meets U in two points $(h(\tau_1))$ and $(h(\tau_2))$. Let $\mu: I \to A$ be a path joining $(h(\tau_1))$ to $(h(\tau_2))$; $p \mu = \beta$ is a path from $j(\tau_1)$ to $j(\tau_2)$ contained in the orbit pA.

For $0 \le \tau \le 1$, define $\eta(\tau): I \to V$ by

$$\eta(\tau)(t) = \begin{cases} j(2t\tau_1), t \le \frac{1}{2} \\ \beta(\tau(2t-1)), t \ge \frac{1}{2} \end{cases}$$

Then $\eta(0) = h(\tau_1) \circ C_{j(\tau_1)}$, $\eta(1) = h(\tau_1) \circ \beta$ so that $\eta h(\tau_1)^{-1}$ is homotopic to C_{x_0} . Let f be the path in W, $f(\tau) = (\eta(\tau))$. We have $pf(\tau) = \eta(\tau)(1) = \beta(\tau)$ and $f(0) = (\eta(0)) = (h(\tau_1))$. Since $p \mu = \beta$ and $\mu(0) = (h(\tau_1))$, we have $\mu = f$; in particular $\mu(1) = f(1)$, $(h(\tau_2)) = (\eta(1)) = (h(\tau_1)\beta)$ so that $h(\tau_1)\beta h(\tau_2)^{-1}$ represents an element of H. Hence

$$\int W h(\tau_1)\beta h(\tau_2)^{-1}$$

is an integer multiple of $\int_i w$. However,

$$\int_{h(\tau_1)\beta} w = \int_{h(\tau_1)} w - \int_{h(\tau_2)} w + \int_{\beta} w = \tau_1 - \tau_2$$

i.e., $\int_{\beta} w = 0$ since β lies in one leaf. Consequently, $\tau_1 = \tau_2$ or $\tau_1 = 1$, $\tau_2 = 0$. In any case $(h(\tau_1)) = (h(\tau_2))$ and A meets U in at most one point.

Now we will show A meets U in at least one point. Let (h) be a point of A. We shall construct a map $G: I \times I \to V$ satisfying: G(1, t) = h(t), G(0, t) = h(a)(t) for some real number a, $G(s, 0) = x_0$ and G(s, 1) is in the orbit through h(1) for $0 \le s \le 1$. The map $s \to (G(s,))$ is then a path in A joining (h) to (h(a)); where G(s,) means the map G(s,)(t) = G(s, t). Since (ha) is a point of V this will complete the proof. Observe that a curve h in V is homotopic to a curve consisting of segments such that each segment is an arc of an orthogonal trajectory or is entirely contained in one leaf. Therefore we may assume there exists numbers $0 = t_0 \le t_1 \le \cdots < t_k = 1$ such that for each i, the arc $h[t_i, t_{i+1}]$ is either a segment of an orthogonal trajectory or is contained in one leaf.

Let L be a leaf of Φ and $x \in L$; C(t) a curve in L starting at x. The orthogonal trajectories are infinitely extendable, hence for any positive number s_0 , the orthogonal trajectories of length s_0 along C define a map $F: I \times [0, s_0] \rightarrow V$ such that for fixed t, F(t, s) is an orthogonal trajectory with F(t, 0) = C(t), and F(t, s) is the point a distance s from C(t) along the orthogonal trajectory through C(t). Moreover, the metric on V guarantees the points F(t, s), for fixed s, are contained in the leaf through F(0, s).

Now G is defined as follows. We may assume $h[t_0, t_1]$ is contained in the leaf L through x_0 , and $h[t_1, t_2]$ is an orthogonal arc. Let C be the path $h[t_0, t_1]$ and s_0 the length of $h[t_1, t_2]$. Apply the last paragraph to obtain a map $F_1: I \times [0, s_0] \rightarrow V$ such that $F_1(0, s) = j(s)$, $F_1(1, s) = h(t_1 + s)$ and $F_1(t, s_0)$ is in the orbit through $j(s_0)$ for $0 \le t \le 1$. Repeat this construction with C the curve $F_1(t, s_0)$ followed by $h[t_2, t_3]$. Induction on k yields the desired map G. This completes the proof of 1.4.

COROLLARY 1.5. Let V be a closed n manifold which cannot be covered by $R^{n-1} \times S^1$. Then a locally free action of R^{n-1} on V has a compact orbit.

LEMMA 1.6. Let $D = \{(x_1, x_2, 0, ..., 0) \in \mathbb{R}^n | x_1^2 + x_2^2 \le 1\}, \{e_1, ..., e_{n-1}\}$ the n-1 frame on ∂D defined as follows: $e_1(x_1, x_2, 0, ..., 0) = (-x_2, x_1, 0, ..., 0), e_2 = (0, 0, 1, 0, ..., 0), ..., e_{n-1} = (0, 0, ..., 0, 1)$. Then $\{e_1, ..., e_{n-1}\}$ does not extend to an n-1 frame on D.

The frame $\{e_1, ..., e_{n-1}\}$ represents the nonzero element of $\pi_1(SO(n))$. This is proved in Chevalley's book on Lie Groups.

THEOREM 1.7. Let V be a simply connected closed n manifold. The rank of V is less than n-1.

Proof. The case n=3 has been proved by Lima [1], and n=4 is trivial since a simply connected 4 manifold does not admit a foliation of codimension one; it does not admit a nonzero vector field. So we assume $n \ge 5$.

Let Φ be a locally free action of \mathbb{R}^{n-1} on V. According to 1.5, Φ has a torus orbit T. Since V is simply connected, $i:T \subset V$, induces the zero homomorphism. Thus there is a simple closed curve C on T which bounds an embedded two-dimensional disk D in V such that D is transverse to T, (here we use $n \ge 5$). But this contradicts 1.6, (cf. [1]).

2. Locally Free Actions of R^2 on $M \times S^1$

(2.1) Let D be a two-dimensional disk with k contours in the interior of D. Let $V = D \times I$ and S be an embedded sphere in V. Then S bounds an embedded ball.

Proof. For k=0 this is Schoenflies Theorem. We consider the case k=1. Let C be an embedding of [0,1] in D with one endpoint on ∂D , the other on the contour, and interior $C \subset \text{interior } D$. If $S \cap A \neq \Phi$, $A = C \times I$, then we may cut V along A to obtain a 3 ball; this is the case k=0. Assume then, that $S \cap A \neq \Phi$ and the intersection is transverse. This is no loss of generality since S may be approximated by an embedded sphere which is transverse to A and then there is a diffeomorphism of V sending one sphere onto the other. Let $a_1 \dots, a_k$ be the simple closed curves in $S \cap A$. Choose a_j so that a_j bounds a disk E on S and E contains no a_i in its interior. A is homeomorphic to $I \times I$ so a_j bounds a disk F on A. Consider the sphere $E \cup F$. For our purposes this sphere is disjoint from A, i.e., $E \cup F$ bounds a ball B in V. Now by an isotopy of B across A we obtain a sphere S_0 which intersects A in the curves $a_1 \cup \dots \cup \hat{a}_j \cup \dots \cup a_k$ (cf. [4] for details). Continuing we see S is isotopic to a sphere which does not intersect A, hence bounds a ball. The general case is just as easy.

Suppose there are k contours with k > 1. Let C be an embedding of I in D with both endpoints on distinct contours and interior $C \subset \text{interior } D$. If $S \cap A = \Phi$, $A = C \times I$, then by cutting V along A we reduce the problem to k-1 contours. Otherwise we take the intersection to be transverse and displace S off A as above.

(2.2) Let M be a closed two-dimensional orientable manifold of connectivity h>1. Let S be a sphere embedded in $M \times S^1$. Then S bounds an embedded ball in $M \times S^1$.

Proof. Let $a_1, ..., a_k$, k = (h+1)/2 be simple closed curves on M as indicated in figure 1.



Denote by $A_i = a_i \times S^1$, and $A = A_1 \cup \cdots \cup A_k$. A separates V into two connected components E_1 and E_2 ; $E_1 = W_1 \times S_1$, $E_2 = W_2 \times S_1$, where W_1 , W_2 are the connected components of $M - (a_1 \cup \cdots \cup a_k)$. W_1 and W_2 are disks with k-1 contours. We may think of $M \times S^1$ as the quotient space of $M \times I$ where (x, 0) is identified with (x, 1), and we identify M with $M \times 0 \in M \times S^1$.

Suppose S is embedded in V so that S is disjoint from M. If S is also disjoint from A then S is contained in E_1 or E_2 . Assume $S \subset E_1$. We have $E_1 = W_1 \times I$ where $W_1 \times 0$ is identified with $W_1 \times 1$. Since $S \cap M = \Phi$, S is really contained in a subspace of V homeomorphic to $W_1 \times I$ and by (2.1), S bounds a ball in this subspace, hence in V. Otherwise we may assume S meets A transversally. Let b be a simple closed curve in $S \cap A$ such that b bounds a disk E on S whose interior is disjoint from A. Since $S \cap M = \Phi$, b bounds a disk F contained in $a_i \times I$ for some i. Then $F \cup E$ is a spheric contained in $W_1 \times I$ or $W_2 \times I$ hence $F \cup E$ bounds a ball. Now by displacing E across this ball we see that S is isotopic to a sphere having one less circle of intersection with $A \cap M$ is void hence this sphere bounds a ball and S also bounds a ball.

It remains to consider the case $S \cap M \neq \Phi$. Let S meet M transversally, and b be a simple closed curve in $S \cap M$ which bounds a disk E on S whose interior is disjoint from M. Since the inclusion of M in V induces a monomorphism of $\pi_1(M)$ into $\pi_1(V)$, b must be null homotopic on M hence b bounds a disk F on M. The sphere $E \cup F$ is (for all practical purposes) disjoint from M hence bounds a ball in V. Then S may be displaced in V to a sphere having one less intersection curve with M and iterating the process removes S from M entirely. This completes the proof of 2.2.

(2.3) Let T be a torus embedded in the interior of $M \times I$ where M is a closed orientable two-dimensional manifold of genus greater than one. Then T separates $M \times I$ into two connected components. Moreover $M \times 0$ and $M \times I$ are contained in the same connected component.

Proof. Let *i* be the inclusion map of *T* into $M \times I$. The map $i_*: H_2(T) \rightarrow H_2(M \times I)$ is zero since $M \times I$ may be retracted onto $M \times 0 = M$, and *M* has genus greater than one so any map of *T* to *M* has degree zero. We must compute $H_0(M \times I - T)$ (all homology and cohomology groups are with Z_2 coefficients). By Lefshetz Duality $H_0(M \times I - T)$ is isomorphic to $H^3(M \times I; T)$. Consider the exact sequence in cohomology:

$$H^{2}(M \times I) \rightarrow H^{2}(T) \rightarrow H^{3}(M \times I; T) \rightarrow H^{3}(M \times I) \rightarrow H^{3}(T)$$

The first map is zero since it is the transpose of i_* and the last group is zero. The second and fourth groups are Z_2 , hence $H^3(M \times I;T) = Z_2 + Z_2$. This proves the first part of 2.3.

Now we will prove $M \times 0$ and $M \times 1$ are in the same component. Let a_1 and a_2 be simple closed curves on $M \times 0$, as in 2.2.

Let T intersect $a_1 \times I$ and $a_2 \times I$ transversally. If T is disjoint from $a_1 \times I$ or $a_2 \times I$ then we may find a curve from $M \times 0$ to $M \times 1$ not meeting T. Assume then that $T \cap (a_1 \times I) = b_1 \cup \ldots \cup b_k$, $T \cap (a_2 \times I) = c_1 \cup \ldots \cup c_l$, where the b_i 's and c_j 's are pairwise disjoint simple closed curves.

If each b_i , or each c_j , is null homotopic in $M \times I$, then we can join $a_1 \times 0$ to $a_1 \times 1$ (or $a_2 \times 0$ to $a_2 \times 1$) by arcs in $a_1 \times I - T$ (or $a_2 \times I - T$). So we may suppose there is a b_i and c_j such that b_i and c_j are not homotopically trivial. Clearly b_i is homotopic to a_1 and c_j to a_2 . Now b_i and c_j are disjoint simple closed curves on the torus T and both represent generators of $\pi_1(T)$, hence b_i and c_j are the boundary circles of a cylinder on T. This implies a_1 is homotopic to a_2 in M which is a contradiction. Thus $M \times 0$ and $M \times 1$ are in the same connected component of $M \times I - T$.

(2.4) Let T be a torus embedded in $M \times S^1$ where M is a closed orientable two manifold of genus greater than one. If $T \cap (M \times x_0) = \Phi$ for some $x_0 \in S^1$, then T separates $M \times S^1$ into two connected components A and B. If h and g are the inclusion maps of T into A and B respectively, then $h_*:\pi_1(T) \to \pi_1(A)$ or $g_*:\pi_1(T) \to \pi_1(B)$ has a nonzero kernel.

It remains to establish the latter assertion of 2.4. First we need an algebraic fact whose proof may be found in Kurosh, volume two.

(2.5) Let G_1 , G_2 and H be groups such that there are subgroups H_1 and H_2 of G_1 , G_2 respectively each isomorphic to H. Denote by $G_1^*HG_2$ the free product of G_1 and G_2 with H amalgamated. Every element of $G_1^*HG_2$ can be written uniquely in the form

$$h \, \bar{a}_1 \, \bar{a}_2 \dots \bar{a}_n$$

where $h \in H$, $n \ge 0$, a_i is a coset representative, other than the unit element, of a right coset of H_i in G_i , i=1,2, and adjacent representatives $\overline{a_i}$, $\overline{a_{i+1}}$, i=1,...,n-1, lie in distinct G_i 's.

From this it follows easily that the center of $G_1 * HG_2$ is contained in H.

Proof of 2.4. Suppose that h_* and g_* are both monomorphisms. Let $G_1 = \pi_1(A)$, $H_1 = h_* \pi_1(T)$, $G_2 = \pi_1(B)$, $H_2 = g_* \pi_1(T)$ and $H = \pi_1(T)$. According to Van Kampen's Theorem and (2.3) we have

$$\pi_1(M\times S^1)=G_1^*HG_2.$$

Since $\pi_1(S^1)$ is contained in the center of $\pi_1(M \times S^1)$ and the center of $G_1 * HG_2$ is contained in H, we have $\pi_1(S^1)$ contained in $\pi_1(T)$. But T is disjoint from $M \times x_0$ for some $x_0 \in S^1$, hence no curve on T can represent a generator of $\pi_1(S^1)$. Thus h_* or g_* is not a monomorphism.

(3.1) Let Φ be a locally free action of \mathbb{R}^2 on $M \times S^1$ with M a closed two manifold

of genus greater than one. Then Φ has a compact orbit, and each compact orbit of Φ intersects M.

This follows immediately from 1.4, 2.4 and [1].

(3.2) If T is a compact orbit of Φ , then $T \cap M$ contains a curve which is a generator of $\pi_1(T)$.

Proof. Assume T is transverse to M and each curve in $T \cap M$ is trivial in $\pi_1(T)$. Let b be such a curve. Then b bounds a disk E on T, hence also bounds a disk F on M and the sphere $E \cup F$ bounds a ball in $M \times S^1$ by 2.2. Thus the intersection curve b may be removed from m by an isotopy of $M \times S^1$ and all intersection curves may be so removed. This gives rise to a new action which is locally free and has a compact orbit disjoint from M. But 3.1. contradicts this.

REFERENCES

[1] E. LIMA, Commuting Vector Fields on S³, Annals Math. 81 (1965), 70-81.

[2] G. REEB, Sur certaines proprietes topologiques des varietes feuilletes, (Paris 1952).

[3] B. REINHART, Foliated manifolds with bundle-like-metrics, Annals Math. 69 (1959), 119-131.

[4] H. ROSENBERG, The rank of $S^2 \times S^1$, American J. of Math. 87 (1965), 11–24.

[5] R. SACKSTEDER, Foliations and pseudogroups, American J. of Math. 87 (1965), 79-102.

Received December 18, 1965