# Every Odd Dimensional Homotopy Sphere has a Foliation of Codimension One

Autor(en): Tamura, Itiro

Objekttyp: Article

Zeitschrift: Commentarii Mathematici Helvetici

Band (Jahr): 47 (1972)

PDF erstellt am: 27.05.2024

Persistenter Link: https://doi.org/10.5169/seals-36356

#### Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

#### Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek* ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

# http://www.e-periodica.ch

# **Every Odd Dimensional Homotopy Sphere has a Foliaton of Codimension One**

by Itiro Tamura

It is well-known that Reeb constructed a foliation of codimension one on  $S^3$  (Reeb [4]). But, after that, nothing was known of codimension-one foliations of higher dimensional spheres for twenty years. In the circumstances Lawson's recent work is significant. He exhibited foliations of codimension one on each of the  $(2^k+3)$ -sphere for k=1, 2, ... (Lawson [2]).

In this paper we shall prove the following.

THEOREM. Every odd dimensional homotopy sphere has a foliation of codimension one.

#### 1. Fiberings over a Circle

Let  $\tilde{S}^{2m+1}$  be a (2m+1)-dimensional homotopy sphere  $(m \ge 3)$  and let  $F^{2m}$  be a compact 2m-dimensional differentiable manifold imbedded in  $\tilde{S}^{2m+1}$  which has the homotopy type of the bouquet of r copies of m-sphere  $S^m$ :

$$F^{2m} \simeq \underbrace{S^m \vee S^m \vee \cdots \vee S^m}_{\mathbf{r}}.$$

Since the normal bundle of the (2m-1)-dimensional differentiable manifold  $\partial F$ , the boundary of  $F^{2m}$ , is trivial, the tubular neighborhood of  $\partial F$  is  $\partial F \times D^2$ . Thus  $\tilde{S}^{2m+1} - (\partial F \times \operatorname{Int} D^2)$  is a (2m+1)-dimensional differentiable manifold with boundary  $\partial F \times S^1$ . In the following the intersection  $F^{2m} \cap (\tilde{S}^{2m+1} - (\partial F \times \operatorname{Int} D^2))$  is simply denoted as  $F^{2m}$ , because they are naturally diffeomorphic.

Let A be the compact (2m+1)-dimensional differentiable manifold (with corner) obtained by splitting  $\tilde{S}^{2m+1} - (\partial F \times \operatorname{Int} D^2)$  at  $F^{2m}$ . Then  $\partial A = F^+ \cup F^- \cup (\partial F \times I)$ , where  $F^+$  and  $F^-$  are copies of  $F^{2m}$ . A has the same homotopy type as  $\tilde{S}^{2m+1} - F^{2m}$ . It is easy to see that A is simply connected and that, by the Alexander duality, homology groups of A are as follows:

$$H_q(A) = \begin{cases} Z & q = 0, \\ Z + Z + \dots + Z & q = m, \\ \hline r & 0 & \text{otherwise.} \end{cases}$$

(Homology groups  $H_*()$  mean homology groups with integral coefficient group  $H_*(; Z)$ .)

Let  $\alpha_1, \alpha_2, ..., \alpha_r$  be a system of generators of  $H_m(F) \cong Z + Z + \cdots + Z$  such that each  $\alpha_i$  is represented by an imbedded *m*-sphere  $S_i(i=1, 2, ..., r)$ . Let  $\alpha_i^+$  (resp.  $\alpha_i^-$ ) denote the element of  $H_m(F^+)$  (resp.  $H_m(F^-)$ ) corresponding to  $\alpha_i \in H_m(F)$ . Let  $\alpha'_i$ denote the element of  $H_{2m}(A)$  corresponding to  $\alpha_i \in H_{2m}(F)$  by the Alexander duality (i=1, 2, ..., r). Then  $\alpha'_1, \alpha'_2, ..., \alpha'_r$  form a system of generators of  $H_m(A)$ .

Let  $\iota^+: F^+ \to A$  and  $\iota^-: F^- \to A$  be the inclusion maps, and let

$$\iota_*^+(\alpha_i^+) = \sum_j a_{ij}^+ \alpha_j', \quad \iota_*^-(\alpha_i^-) = \sum_j a_{ij}^- \alpha_j' \quad (i = 1, 2, ..., r)$$

Then  $a_{ij}^+$  and  $\bar{a}_{ij}$  are expressed by linking numbers as follows.

Denote by  $S_i^+$  (resp.  $S_i^-$ ) a displacement of  $S_i$  in  $\tilde{S}^{2m+1}$  towards the normal direction of  $F^+$  (resp.  $F^-$ ). Then it is easy to see that

$$a_{ij}^{+} = Lk(S_i^{+}, S_j), \quad a_{ij}^{-} = Lk(S_i^{-}, S_j).$$

Furthermore it follows from

$$Lk(S_{i}^{+}, S_{j}) = Lk(S_{i}, S_{j}^{-}) = (-1)^{m+1}Lk(S_{j}^{-}, S_{i}),$$

that

 $a_{ij}^+ = (-1)^{m+1} a_{ji}^-$ 

Denote by L(F) the following  $(r \times r)$  matrix:

$$L(F) = \begin{pmatrix} Lk(S_{1}^{+}, S_{1}) \dots Lk(S_{1}^{+}, S_{r}) \\ \vdots & \ddots & \vdots \\ Lk(S_{r}^{+}, S_{1}) \dots Lk(S_{r}^{+}, S_{r}) \end{pmatrix}$$

Suppose now that L(F) is unimodular. Then the homomorphisms

$$\iota_*^+: H_m(F^+) \to H_m(A), \iota_*^-: H_m(F^-) \to H_m(A)$$

are isomorphisms. This shows, since  $F^+$ ,  $F^-$  and A are simply connected, that  $\iota^+$  and  $\iota^-$  are homotopy equivalence. Thus, according to the relative *h*-cobordism theorem (Smale [5], Corollary 3.2), the following holds:

 $(A, \partial F \times I) = (F^{2m}, \partial F) \times I.$ 

This implies the following proposition which is a differential topological version of so-called Milnor fibering. (See also Tamura [6].)

**PROPOSITION 1.** If the matrix L(F) is unimodular, then there exists a fibering  $\tilde{S}^{2m+1} - (\partial F \times \operatorname{Int} D^2) \rightarrow S^1$  having F as a fibre.

### 2. Construction of Fiberings

Let X, Y denote the following matrices

$$X = \begin{pmatrix} 2 & 1 & & 0 \\ 2 & 1 & & 0 \\ 1 & 2 & 1 & & \\ 1 & 2 & 1 & & \\ 1 & 1 & 2 & 1 & & \\ & 1 & 2 & 1 & & \\ & & 1 & 2 & 1 & & \\ 0 & & & 1 & 2 & 1 & \\ & & & 1 & 2 & 1 & \\ 0 & & & & 1 & 2 & 1 \\ & & & & 1 & 2 & 1 \\ 0 & & & & 1 & 2 & 1 \\ 0 & & & & 1 & 2 & 1 \\ 0 & & & & 1 & 2 & 1 \\ \end{pmatrix}$$

As is well known, X is positive definite and unimodular. The rank of  $(9 \times 9)$  matrix Y is 8 and its elementary divisor is (1, 1, 1, 1, 1, 1, 1, 1).

Let  $\Delta$  denote the diagonal of  $S^{2n} \times S^{2n}$   $(n \ge 2)$  and let N be a tubular neighborhood of  $\Delta$  in  $S^{2n} \times S^{2n}$ . Then N has the homotopy type of  $S^{2n}$  and the self-intersection number of a generator of  $H_{2n}(N) \cong Z$  is 2. Let W(X) be the parallelizable compact oriented 4n-dimensional differentiable manifold formed from  $N_1, N_2, ..., N_8$  (8 copies of N), by plumbing  $N_i$  and  $N_{i+1}$  (i=2, 3, ..., 7), and  $N_1$  and  $N_4$ . Then W(X) has the homotype types of  $\underbrace{S^{2n} \vee S^{2n} \vee \cdots \vee S^{2n}}_{8}$ . The orientation of W(X) is chosen so that



Fig. 1.

the matrix of intersection numbers of  $H_{2n}(W(X))$  is X. Similarly parallelizable compact oriented 4*n*-dimensional differentiable manifolds W(-X) and W(Y) both of which have the homotopy type of bouquets of 2*n*-spheres, are defined. The matrix of intersection numbers of  $H_{2n}(W(-X))$  (resp.  $H_{2n}(W(Y))$ ) is -X (resp. Y).

Let  $W = W(-X) \models W(Y)$  be the boundary connected sum of W(-X) and W(Y). W is a parallelizable compact oriented 4n-dimensional differentiable manifold. Let us imbed W into  $\tilde{S}^{4n+1}$  as indicated in the Fig. 1, by 17 copies of naturally imbedded  $S^{2n} \times S^{2n}$  which osculate consecutively, so that unnecessary linking numbers do not occur in the matrix L(W) (cf. Tamura [6], section 2).

Then it is easy to see that the matrix L(W) of linking numbers is given by

$$L(W) = \begin{pmatrix} P & \\ & Q \end{pmatrix},$$

where

Thus, by Proposition 1, the following holds.

PROPOSITION 2. There exists a fibering  $\tilde{S}^{4n+1} - (\partial W \times \operatorname{Int} D^2) \to S^1$  having W as a fibre  $(n \ge 2)$ .

Let  $\hat{\Delta}$  denote the diagonal of  $S^{2n-1} \times S^{2n-1}$   $(n \ge 2)$  and let  $\hat{N}$  be a tubular neighborhood of  $\hat{\Delta}$  in  $S^{2n-1} \times S^{2n-1}$ . Let us imbed  $\hat{N}$  into  $\tilde{S}^{4n-1}$  by imbedding  $S^{2n-1} \times S^{2n-1}$ into  $\tilde{S}^{4n-1}$  naturally. Then the matrix  $L(\hat{N})$  of linking numbers is given by  $L(\hat{N}) = (1)$ . Thus, by Proposition 1, the following holds.

PROPOSITION 3. There exists a fibering  $\tilde{S}^{4n-1} - (\partial \hat{N} \times \text{Int} D^2) \rightarrow S^1$  having  $\hat{N}$  as a fibre  $(n \ge 2)$ .

This fibering corresponds to the Milnor fibering of  $z_0^2 + z_1^2 + \dots + z_{2n-1}^2 = 0$ .

#### 3. Boundary of the Fibre W

Let *M* denote the boundary of the fibre *W* in Proposition 2. Then *M* is an orientable closed (4n-1)-dimensional differentiable manifold. It follows by the PoincaréLefschetz duality that

 $H_q(W, M) \cong H^{4n-q}(W),$ 

and that the natural homomorphism

 $H_{2n}(W) \rightarrow H_{2n}(W, M) \cong \operatorname{Hom}(H_{2n}(W), Z)$ 

is determined by  $\begin{pmatrix} -X \\ Y \end{pmatrix}$ , the matrix of intersection numbers of  $H_{2n}(W)$ . Thus the following is a direct consequence of the homology exact sequence of (W, M):

$$H_q(M) = \begin{cases} Z & q = 0, 2n - 1, 2n, 4n - 1, \\ 0 & \text{otherwise.} \end{cases}$$

Obviously W is obtained from  $W(-X) \models W(X)$  by attaching a handle  $D^{2n} \times D^{2n}$ :

$$W = (W(-X) \natural W(X)) \bigcup_{g} (D^{2n} \times D^{2n}),$$

where  $g:\partial D^{2n} \times D^{2n} \to \partial (W(-X) \models W(X))$  is an attaching map. The boundary  $\partial (W(-X) \models W(X))$  is the natural (4n-1)-sphere (Kervaire-Milnor [1]), and the following decomposition holds:

$$W = W(-X) \natural W(X) \natural \left( D^{4n} \bigcup_{g} \left( D^{2n} \times D^{2n} \right) \right).$$

According to the *h*-cobordism theorem (Smale [5]),  $B = D^{4n} \bigcup_g (D^{2n} \times D^{2n})$  is the total space of a 2*n*-disk bundle  $\xi$  over  $S^{2n}$ , and its differentiable structure is compatible with the bundle structure. Thus  $M = \partial W = \partial B$  is the total space of an  $S^{2n-1}$ -bundle over  $S^{2n}$  associated with  $\xi$ . Let  $\alpha \in \pi_{2n-1}(SO(2n))$  be the characteristic map of  $\xi$ . Since B is parallelizable,  $\xi$  is stably trivial and, thus,  $\alpha$  belongs to the kernel of  $\pi_{2n-1}(SO(2n)) \to \pi_{2n-1}(SO(2n+1))$ . Let us consider the diagram

consisting of the homotopy exact sequence of the fibering  $SO(2n+1) \rightarrow SO(2n+1)/SO(2n) = S^{2n}$  and the homomorphism induced by the projection  $p: SO(2n) \rightarrow SO(2n)/SO(2n-1) = S^{2n-1}$ . Let  $\iota_{2n}$ ,  $\iota_{2n-1}$  be generators of  $\pi_{2n}(S^{2n})$ ,  $\pi_{2n-1}(S^{2n-1})$  respectively. Since  $\alpha \in \partial(\pi_{2n}(S^{2n}))$ ,  $\alpha = \partial(c\iota_{2n})$  for an integer c. If  $c \neq 0$ ,  $p_*\partial(c\iota_{2n}) = \pm 2c\iota_{2n-1} \neq 0$ and, thus, the Euler class of  $\xi$  is not zero. This implies, by using the Thom-Gysin exact sequence, that  $H_{2n-1}(M) = H_{2n-1}(\partial B) \not\cong Z$ , which is a contradiction. Thus c=0and  $\xi$  is a trivial bundle. This yields the following. LEMMA 1. The boundary of W is  $S^{2n-1} \times S^{2n}$ .

## 4. Proof of Theorem

Let E be a compact connected (2m+1)-dimensional differentiable manifold such that E is a total space of a fibering over  $S^1$  and  $\partial E$  is connected. Then it is well known that there exists a foliation of codimension one on E having  $\partial E$  as the only compact leaf (cf. Lawson [2]).

LEMMA 2. Suppose that  $S^{2n+1}$  has a foliation of codimension one  $(n \ge 2)$ , then the following holds:

(i) Any (4n+1)-dimensional homotopy sphere  $\tilde{S}^{4n+1}$  has a foliation of codimension one.

(ii) Any (4n-1)-dimensional homotopy sphere  $\tilde{S}^{4n-1}$  has a foliation of codimension one.

*Proof.* Let  $\gamma$  be a closed smooth curve in  $S^{2n+1}$  which is transversal to the leaves. The existence of such  $\gamma$  is a classical fact. The tubular neighborhood of  $\gamma$  is  $S^1 \times D^{2n}$ . The foliation on  $S^{2n+1}$  can be modified so that its restriction on  $S^{2n+1} - (S^1 \times \text{Int} D^{2n})$  $= S^{2n-1} \times D^2$  is a foliation having the boundary as a compact leaf.

Now, by Proposition 2,  $\tilde{S}^{4n+1} - (\partial W \times \operatorname{Int} D^2)$  has a foliation of codimension one such that  $\partial W \times S^1$  is the only compact leaf. On the other hand, since  $\partial W = S^{2n-1} \times S^{2n}$  by Lemma 1,  $\partial W \times D^2$  has a foliation of codimension one which is induced by the projection  $\partial W \times D^2 \to S^{2n-1} \times D^2$  from the foliation of  $S^{2n-1} \times D^2$ . This completes the proof of (i).

Making use of Proposition 3 and the projection  $\partial \hat{N} \times D^2 \rightarrow \hat{\Delta} \times D^2 = S^{2n-1} \times D^2$ , the proof of (ii) is completely analogous to that of (i).

Remark. Lemma 2, (ii) is (a slightly generalized form of) a result of Lawson [2].

Let  $\tilde{S}^{2m+1}$  be a (2m+1)-dimensional homotopy sphere. In case m=1, 2, the existence of a foliation of codimension one is proved by Novikov [3] and Lawson [2] respectively. Suppose that, for  $2 \leq m < q$ ,  $\tilde{S}^{2m+1}$  has a foliation of codimension one. Then, if q is even (resp. odd), the existence of a foliation of codimension one of  $\tilde{S}^{2q+1}$  is assured by Lemma 2 (i) (resp. (ii)). This completes the proof of the theorem by induction.

#### REFERENCES

- [1] KERVAIRE M. and MILNOR J., Groups of homotopy spheres I, Ann. Math. 77 (1963) 504-537.
- [2] LAWSON H. B., Codimension-one foliations of spheres, Bull. Amer. Math. Soc. 77 (1971) 437-438.
- [3] NOVIKOV S. P., The topology of foliations, Trudy Moscov Mat. Obšč., 14 (1965) 248-278.

#### ITIRO TAMURA

- [4] REEB G., Sur certains propriétés topologiques des variétés feuilletées, Actualités Sci. Indust., No. 1183, Hermann, Paris, 1952.
- [5] SMALE S., On the structure of manifolds, Amer. J. Math., 84 (1962) 387-399.
- [6] TAMURA I., Fixed point sets of differentiable periodic transformations on spheres, J. Fac. Sci. Univ. Tokyo, Sect. I, 16 (1969) 101-114.

Department of Mathematics University of Tokyo.

Received November 19, 1971.

### 170