

# Explicit Quasiconformal Extensions for some Classes of Univalent Functions

Autor(en): **Fait, Maria / Krzyz, J.G. / Zygmunt, Jadwiga**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **51 (1976)**

PDF erstellt am: **28.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-39443>

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# Explicit Quasiconformal Extensions for some Classes of Univalent Functions

MARIA FAIT, JAN G. KRZYŻ AND JADWIGA ZYGMUNT

## 1. Introduction. Notations

Let  $S$  be the class of functions analytic and univalent in  $\Delta = \{z : |z| < 1\}$  for which  $f(0) = f'(0) - 1 = 0$ .

We say that  $f \in S_k$ ,  $0 \leq k < 1$ , if  $f \in S$  and  $f$  has a quasiconformal extension on the whole plane  $\mathbb{C}$  with complex dilatation  $\mu_f = f_{\bar{z}}/f_z$  that satisfies  $|\mu_f(z)| \leq k$  almost everywhere in  $\mathbb{C}$ . The symbols  $f_z$ ,  $f_{\bar{z}}$  denote formal derivatives of  $f$ .

Let  $S^*(\alpha)$ ,  $0 \leq \alpha < 1$ , denote the subclass of  $S$  consisting of strongly starlike functions of order  $\alpha$ , cf. [1], [5], i.e. of functions  $f$  that satisfy:

$$\left| \arg \frac{zf'(z)}{f(z)} \right| \leq \alpha \frac{\pi}{2}, \quad z \in \Delta. \quad (1)$$

As shown in [1],  $f(\Delta)$  is a Jordan domain for any  $f \in S^*(\alpha)$ .

In this paper we find an explicit quasiconformal extension for an arbitrary function  $f \in S^*(\alpha)$ . We show that  $S^*(\alpha) \subset S_k$ , where  $k \leq \sin \alpha\pi/2$ .

We construct this extension by means of an auxiliary mapping which may be called a reflection with respect to a starlike Jordan curve (Lemma 1). In what follows we call a  $k$ -circle a Jordan curve that is a homeomorphic image of the unit circumference under a quasiconformal mapping  $F$  of the extended plane  $\mathbb{C}$  onto itself whose complex dilatation  $\mu_F$  satisfies  $|\mu_F(z)| \leq k < 1$  a.e.

We obtain explicit quasiconformal extensions for bounded convex functions and for functions with bounded boundary rotation (Theorems 3,4). In particular we show that any convex Jordan curve contained in an annulus  $\{w : r \leq |w| \leq R\}$  is a  $k$ -circle with  $k \leq \sqrt{1 - (r/R)^2}$ .

Similarly, any strongly starlike curve of order  $\alpha$  is a  $k$ -circle with  $k \leq \sin \alpha\pi/2$ .

## 2. Quasiconformal Extension for the Class $S^*(\alpha)$

In this section we shall prove

**THEOREM 1.** *If  $f \in S^*(\alpha)$ ,  $0 \leq \alpha < 1$ , then the mapping  $F$  defined by the formula*

$$F(z) = \begin{cases} f(z) & \text{for } |z| \leq 1 \\ |f(\xi)|^2 / \overline{f\left(\frac{1}{\bar{z}}\right)}, & \text{for } |z| \geq 1, \end{cases} \quad (2)$$

where  $\xi$  satisfies the conditions:  $|\xi| = 1$ ,  $\arg f(\xi) = \arg f(1/\bar{z})$ , belongs to the class  $S_k$  and  $|\mu_F(z)| \leq k = \sin \alpha\pi/2$  a.e.

We first prove

**LEMMA 1.** *Suppose that  $G$  is a domain bounded by a Jordan curve  $\Gamma$  starlike with respect to the origin. Suppose, moreover, that*

$$w = R(\varphi)e^{i\varphi}, \quad 0 \leq \varphi \leq 2\pi, \quad (3)$$

is the parametric equation of  $\Gamma$ , where  $R(\varphi)$  is absolutely continuous and positive,  $R(0) = R(2\pi)$  and

$$|R'(\varphi)|[R^2(\varphi) + R'^2(\varphi)]^{-1/2} \leq k < 1 \quad (4)$$

almost everywhere in  $[0, 2\pi]$ . Then the mapping

$$\phi(w) = R^2(\varphi)/\bar{w}, \quad \varphi = \arg w, \quad (5)$$

is an anticonformal mapping of  $G$  onto  $\hat{\mathbb{C}} \setminus \bar{G}$  whose complex dilatation  $\phi_w/\phi_{\bar{w}}$  is bounded by  $k$  in absolute value.

*Proof.* Obviously  $\phi$  is a sense-reversing homeomorphism in  $G$ . Moreover, if  $w = re^{i\varphi}$ ,  $0 < r < R(\varphi)$ , then

$$\Phi(w) = \Phi(re^{i\varphi}) = R^2(\varphi)e^{i\varphi}/r$$

and we have for almost all  $\varphi$  in  $[0, 2\pi]$ :

$$\mu_\phi = \frac{\phi_w}{\phi_{\bar{w}}} = e^{-2i\varphi} \frac{\phi_r + \phi_\varphi/ir}{\phi_r - \phi_\varphi/ir} = -e^{-2i\varphi} \frac{R'(\varphi)}{R'(\varphi) + iR(\varphi)}$$

so that  $|\mu_\phi(w)| \leq k$  almost everywhere in  $G$  by (4).

We now prove that  $\phi$  has the ACL-property in  $G \setminus \{0\}$ . The function  $R^2(\varphi)e^{i\varphi}/r$  is absolutely continuous in  $\varphi$  with fixed  $r > 0$  because by (4)  $R'(\varphi)$  is essentially bounded and also absolutely continuous in  $r$ ,  $r \in [\delta, R(\varphi)]$ , for fixed  $\varphi$ ,  $\delta > 0$ . Thus the ACL-property holds in the log  $w$ -plane. Since the ACL-property is invariant under composition with conformal mapping,  $\phi$  has in fact the ACL-property in  $G \setminus \{0\}$ . This ends the proof.

The condition (4) has a simple geometrical interpretation. Suppose that  $R'(\varphi)$  does exist. Then  $\Gamma$  has a tangent intersecting the radius vector at an angle  $\psi = \arctan R/R'$  and consequently

$$R'(R^2 + R'^2)^{-1/2} = \cos \psi.$$

Hence (4) means that the angle  $\psi$  is bounded away from 0 and  $\pi$  at points where the tangent does exist.

The mapping  $\phi(w)$  will be called a reflection with respect to the starshaped curve  $\Gamma$ . It is a sense-reversing homeomorphism for any starshaped Jordan curve  $\Gamma$ . Moreover, if the angle between the radius vector and the tangent of  $\Gamma$  is bounded away from 0 and  $\pi$  a.e., the reflection  $\phi(w)$  is an anti-quasiconformal mapping.

*Proof of Theorem 1.* If  $f \in S^*(\alpha)$  with  $0 \leq \alpha < 1$  then  $f$  has a continuous extension on  $\bar{\Delta}$ ,  $f(e^{i\theta})$  is absolutely continuous and  $d/d\theta f(e^{i\theta}) = ie^{i\theta} f'(e^{i\theta})$  a.e. in  $[0, 2\pi]$ , cf. [1]. Hence the definition of  $F$  in (2) makes sense. Let  $\Gamma$  be the Jordan curve  $w = f(e^{i\theta}) = R(\varphi)e^{i\varphi}$ ,  $0 \leq \theta \leq 2\pi$ . After differentiation with respect to  $\theta$  of the identity:

$$\log f(e^{i\theta}) = \log R(\varphi) + i\varphi,$$

we obtain

$$\frac{e^{i\theta} f'(e^{i\theta})}{f(e^{i\theta})} = \left[ 1 - \frac{iR'(\varphi)}{R(\varphi)} \right] \frac{d\varphi}{d\theta}$$

and hence by (1)

$$\left| \arg \left\{ 1 - i \frac{R'(\varphi)}{R(\varphi)} \right\} \right| \leq \frac{\alpha\pi}{2},$$

or

$$|R'(\varphi)|[R^2(\varphi) + R'^2(\varphi)]^{-1/2} \leq \sin \frac{\alpha\pi}{2} \quad \text{a.e.}$$

This means that  $\Gamma$  satisfies the condition (4) with  $k = \sin \alpha\pi/2$  and therefore the reflection with respect to  $\Gamma$  is anti-quasiconformal with complex dilatation bounded by  $\sin \alpha\pi/2$ . Now, the mapping  $F(z)$  for  $|z| > 1$  is composed of the following mappings: reflection in  $|z| = 1$ , conformal mapping  $f$  and a reflection with respect to  $\Gamma$ . Complex dilatation of  $F$  has the form

$$\mu_F = \left( \frac{z}{\bar{z}} \right)^2 \frac{f'(1/\bar{z})}{\overline{f(1/\bar{z})}} \frac{\phi_w}{\phi_{\bar{w}}}.$$

Therefore  $F$  is a quasiconformal mapping in  $\{z : |z| > 1\}$  and  $|\mu_F(z)| \leq \sin \alpha\pi/2$  a.e. by Lemma 1. Obviously  $F$  as defined by (2), is a homeomorphism of the sphere  $\hat{\mathcal{C}}$  onto itself which is conformal in  $\Delta$  and quasiconformal in  $\mathcal{C} \setminus \bar{\Delta}$ . Since  $\partial\Delta$ ,  $\{\infty\}$ ,  $\{0\}$  are removable sets, cf. [4],  $F$  is quasiconformal in  $\hat{\mathcal{C}}$ .

**COROLLARY 1.** *If  $\Gamma$  is a Jordan curve starshaped with respect to  $w = 0$  and intersecting the radius vectors at an angle bounded away from 0 and  $\pi$  by  $\beta\pi/2$ ,  $0 < \beta \leq 1$ , then  $\Gamma$  is a  $k$ -circle with  $k \leq \cos \beta\pi/2$ .*

### 3. Some Applications of Theorem 1

It is well-known that, if

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad \text{in } \Delta \tag{5}$$

and

$$\sum_{n=2}^{\infty} n |a_n| \leq 1 \tag{6}$$

then  $f$  is a starlike univalent function. The condition (6) does not imply the possibility of quasiconformal extension of  $f$  (e.g.  $f(z) = z + \frac{1}{2}z^2$  satisfies (6) and obviously has no quasiconformal extension on  $\hat{\mathbb{C}}$ ).

Consider the class  $\tilde{S}(k)$  of functions  $f$  of the form (5) that satisfy the condition

$$\sum_{n=2}^{\infty} n |a_n| \leq k < 1. \quad (6')$$

We prove

LEMMA 2. *If  $f \in \tilde{S}(k)$ , then  $f \in S^*(\alpha)$  with  $\alpha = (2/\pi) \arcsin k$ .*

*Proof.* The condition (6') implies

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| \leq k,$$

because

$$\left| \frac{zf'(z)}{f(z)} - 1 \right| = \left| \frac{\sum_{n=2}^{\infty} (n-1)a_n z^{n-1}}{1 + \sum_{n=2}^{\infty} a_n z^{n-1}} \right| \leq \frac{\sum_{n=2}^{\infty} (n-1)|a_n|}{1 - \sum_{n=2}^{\infty} |a_n|} \leq k.$$

Hence  $f$  satisfies (1) with  $\alpha = (2/\pi) \arcsin k$ .

From Lemma 2 and Theorem 1 we immediately obtain

THEOREM 2. *If  $f \in \tilde{S}(k)$  then  $f \in S_k$ .*

Another quasiconformal extension of  $f \in \tilde{S}(k)$  can be obtained in a different way, similarly as in [2].

THEOREM 2'. *Let  $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$  belong to  $\tilde{S}(k)$ . Then the mapping  $G(z)$  defined by the formula*

$$G(z) = \begin{cases} z + \sum_{n=2}^{\infty} a_n z^n & \text{for } |z| \leq 1, \\ z + \sum_{n=2}^{\infty} a_n \bar{z}^{-n} & \text{for } |z| \geq 1 \end{cases} \quad (7)$$

*is a quasiconformal extension of  $f$  onto  $\hat{\mathbb{C}}$  and  $|\mu_G(z)| \leq k$ .*

The mapping  $G$  satisfies the following condition:

$$|z_1 - z_2| (1 - k) \leq |G(z_1) - G(z_2)| \leq |z_1 - z_2| (1 + k) \quad (8)$$

for  $z_1, z_2 \in \Delta$  and also for  $z_1, z_2 \in \mathbb{C} \setminus \bar{\Delta}$ . It is well known that a function lipschitzian in  $\Delta$  has a continuous extension on  $\bar{\Delta}$  that satisfies (8) also in  $\bar{\Delta}$ . Hence  $G$  as defined by (7) is a sense-preserving homeomorphism in  $\hat{\mathcal{C}}$ . Its complex dilatation satisfies

$$|\mu_G(z)| = |G_{\bar{z}}/G_z| = \left| \sum_{n=2}^{\infty} n a_n z^{-n-1} \right| \leq \sum_{n=2}^{\infty} n |a_n| \leq k$$

in  $\mathbb{C} \setminus \bar{\Delta}$ . Since  $\partial\Delta$  is a removable set,  $G$  is a quasiconformal in  $\hat{\mathcal{C}}$ .

Let  $C(B)$  denote the subclass of  $S$  consisting of all convex functions for which  $|f(z)| \leq B$ ,  $z \in \Delta$ . Next, let  $V_\lambda(B)$  denote the subclass of  $S$  consisting of all bounded functions  $|f(z)| \leq B$  for which  $f(\Delta)$  has boundary rotation at most  $\lambda\pi$ , cf. [3].

Moreover, let  $d_f$  denote the radius of the largest open disc centered at the origin which is contained in  $f(\Delta)$ .

In [1] Brahnan and Kirwan have found the following relations between  $C(B)$ ,  $V_\lambda(B)$  and  $S^*(\alpha)$ .

- (i) If  $f \in C(B)$ , then  $f \in S^*(\alpha)$  with  $\alpha = 1 - (2/\pi) \arcsin d_f/B$ .
- (ii) If  $f \in V_\lambda(B)$  and  $(\lambda - 2)\pi < 2 \arcsin (d_f/B)$ , then  $f \in S^*(\alpha)$  with  $\alpha = \lambda - 1 - (2/\pi) \arcsin (d_f/B)$ .

The above stated relations yield at once as immediate consequences of Theorem 1 the following results.

**THEOREM 3.** *If  $f \in C(B)$ , then  $f$  has a quasiconformal extension  $F$  on the whole plane defined by the formula (2) and*

$$|\mu_F(z)| \leq \sqrt{1 - \left(\frac{d_f}{B}\right)^2}.$$

**COROLLARY 2.** *If  $\Gamma$  is a convex Jordan curve contained in the annulus  $\{w : r \leq |w| \leq R\}$ , then  $\Gamma$  is a  $k$ -circle with  $k \leq \sqrt{1 - (r/R)^2}$ .*

**THEOREM 4.** *If  $f \in V_\lambda(B)$  and  $(\lambda - 2)\pi < 2 \arcsin (d_f/B)$ , then the function  $F$  defined by (2) is a quasiconformal extension of  $f$  and*

$$|\mu_F(z)| \leq \sin \left[ (\lambda - 1) \frac{\pi}{2} - \arcsin \frac{d_f}{B} \right].$$

## REFERENCES

- [1] BRANNAN, D. A. and KIRWAN, W. E., *On some classes of bounded univalent functions*, J. London Math. Soc. (2), 1 (1969), 431–443.
- [2] KRZYZ, J. G., *Convolution and quasiconformal extension*, Comment Math. Helv.
- [3] LEHTO, O., *On the distortion of conformal mapping with bounded boundary rotation*, Ann. Acad. Sci. Fenn., Ser. A 1, 124 (1952).
- [4] LEHTO, O. and VIRTANEN, K. I., *Quasiconformal mappings in the plane*, Berlin-Heidelberg-New York 1973.
- [5] STANKIEWICZ, J., *Quelques problèmes extrémaux dans les classes des fonctions  $\alpha$ -angulairement étoilées*, Ann. Univ. Marie Curie-Skłodowska Sect. A. XX (1966), 59–75.

*Department of Mathematics*  
*Marie Curie-Skłodowska University*  
*Lublin, Poland*

Received December 30, 1975







