

# Fundamental Groups, ...-groups and Codimension two Submanifolds

Autor(en): **Cappell, Sylvain E. / Shaneson, Julius L.**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **51 (1976)**

PDF erstellt am: **28.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-39453>

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# Fundamental Groups, $\Gamma$ -groups, and Codimension two Submanifolds

SYLVAIN E. CAPPELL AND JULIUS L. SHANESON

## Introduction and Statement of Results

Let  $\phi: M^n \rightarrow W^{n+2}$  be a smooth or piecewise linear (P.L.) embedding (possibly not locally flat) of the connected, compact, oriented smooth or P.L. manifold  $M$  in the compact connected smooth or P.L. manifold  $W$ . Assume that a regular neighborhood of  $\phi(M)$  meets the boundary in a regular neighborhood of  $\phi(\partial M) = \partial W \cap \phi(M)$ ; possibly  $\partial M = \emptyset$  or  $\partial W = \emptyset$ . If  $\phi(M)$  is smooth or P.L. locally flat, the usual geometric procedure, using transversality and counting intersection points together with orientations and associated elements of  $\pi_1 W$  (see, e.g. [W, §5]), gives an intersection number  $x \cdot \phi_*[M]$ ,  $x \in \pi_2 W$ , with values in  $Z \otimes_{Z\pi_1 M} Z\pi_1 W$ . This tensor product is defined using the left  $Z\pi_1 M$ -module structure on the integral group ring  $Z\pi_1 W$  given by<sup>1</sup>  $\alpha\beta = \beta\phi_*(\alpha)$ ,  $\alpha \in Z\pi_1 M$ ,  $\beta \in Z\pi_1 W$ . For the general  $\phi$ ,  $x \cdot \phi_*[M]$  is still defined homologically as in §1 or, alternatively, using geometric intersection numbers defined on the chain level [ST].

**THEOREM 1.** *Let  $K$  be the kernel of the map  $\pi_1(W - \phi M) \rightarrow \pi_1 W$  induced by inclusion. Then  $K_{ab} = K/[K, K]$ ,  $[K, K]$  the commutator subgroup, is isomorphic as a  $Z\pi_1 W$  module ( $\pi_1 W$  acts by conjugation) to*

$$Z \otimes_{Z\pi_1 M} Z\pi_1 W / \{x \cdot \phi_*[M] / x \in \pi_2 W\}.$$

In theorem 1,  $Z$  has the trivial  $Z\pi_1 M$ -module structure. In the non-orientable case the result is still valid, but the  $Z\pi_1 M$ -module structure on  $Z$  is given by the difference of the orientation characters of  $M$  and  $W$ . Theorem 1 also applies to Poincaré embeddings of Poincaré complexes, and hence to locally flat topological

---

<sup>1</sup> In  $Z\pi_1 W$ ,  $(\sum \alpha_g g)^- = \sum \alpha_g g^{-1}$ . Thus  $Z \otimes_{Z\pi_1 M} Z\pi_1 W$  is the free abelian group on the left cosets of  $\phi_*(\pi_1 M)$  in  $\pi_1 W$ . If we used the module structure  $\alpha\beta = \phi_*(\alpha)\beta$ , we would get right cosets, corresponding to writing covering translations as right rather than left operators.

embeddings. In [CS2, §5], it is shown how a general P.L. embedding, locally flat or not, induces an underlying Poincaré embedding.

We apply theorem 1 to the codimension two splitting problem. Suppose  $\phi(M) = P$  is a smooth, P.L. locally flat or topologically locally flat submanifold of  $W$ . (Again, it suffices to consider Poincaré embeddings of Poincaré complexes.) Let

$$f: (Q^{n+2}, \partial Q) \rightarrow (W, \partial W)$$

be a degree 1 map that induces a (simple) homotopy equivalence<sup>(2)</sup> of  $Q$  with  $W$ . Assume that  $f|_{\partial Q}$  is transverse to  $\partial P$  and that  $f|_{f^{-1}(\partial P)}: f^{-1}(\partial P) \rightarrow \partial P$  is a (simple) homology equivalence [CS1] over  $Z\pi_1 P$ . In [CS1, §8] we studied the codimension two splitting problem: when is  $f$  homotopic to  $f_1$ , relative  $\partial Q$ , transverse to  $P$ , with  $f_1|_{f_1^{-1}P}: f_1^{-1}P \rightarrow P$  a simple homotopy equivalence. If the answer is affirmative,  $f$  was said to be (simply) *splittable along  $P$*  (relative boundary.)

Let  $\sum_e(f)$ ,  $e = s, h$ , be the *abstract surgery obstruction* of  $f$ , as defined on page 322 of [CS1] (see also [B1], [W, §11]); i.e.  $\sum_e(f)$  is defined as the surgery obstruction of the normal map  $f$  determines on a transverse inverse image of  $P$ . In 8.2 of [CS1], we saw that for  $n \geq 5$  odd,  $f$  is simply splittable (splittable) if and only if  $\sum_s(f) = 0$  (resp.  $\sum_h(f) = 0$ .) In Thm 8.3 of [CS2], we proved the same result in even dimensions, under some additional hypotheses. These results were applied to the study of the existence and uniqueness of invariant spheres of group actions. In 8.5 of [CS2], we gave a general result on the even dimensional case; there is a further obstruction, in a quotient of a suitable  $\Gamma$ -group, defined if  $\sum_e(f) = 0$ , whose vanishing is necessary and sufficient for  $f$  to be (simply) splittable. The constructions of [CS3] imply that all these obstructions actually arise in codimension two splitting problems.

If  $M$  is closed,  $\phi_*[M]$ ,  $[M] \in H_n(M, \mathbb{Z})$  the orientation, can be considered<sup>(3)</sup> as an element of  $H_n(W, A)$ ,  $A = \mathbb{Z} \otimes_{Z\pi_1 M} Z\pi_1 W$ . Let  $\tau_P = \tau_{\phi(M)}: H_2(W, Z\pi) \rightarrow A$ ,  $\pi = \pi_1 W$ , be given by evaluation (Kronecker product) of the image in  $H^2(W, A)$  of the Poincaré dual in  $H^2(W, \partial W; A)$  of  $\phi_*[M]$ . If  $\partial M \neq \emptyset$ ,  $\phi_*[M]$  is in  $H_n(W, \partial W; A)$ ; again let  $\tau_P$  be the evaluation of its Poincaré dual.

**THEOREM 2.** *Suppose  $n \geq 6$  is even and  $\tau_P$  is surjective. Suppose also that*

<sup>2</sup> By Poincaré duality, it therefore induces a (simple) homology equivalence of  $(Q, \partial Q)$  and  $(W, \partial W)$ , with coefficients in  $Z\pi_1 W$ .

<sup>3</sup> This assumes base points chosen, so that a lift of  $P$  to the covering space of  $W$  with fundamental group  $\phi_*\pi_1 M$  is determined.

$K = \ker(\pi_1(W - P) \rightarrow \pi_1 W)$  is finitely generated. Then  $f$  is (simply) splittable if and only if  $\sum_e(f) = 0$ ,  $e = s$  or  $h$  as appropriate.

Note that if  $\pi_1 W$  is finite,  $K$  is automatically finitely presented. (For example, consider the covering space with  $\pi_1 = K$ .)

For example, the hypotheses of theorem 2 are satisfied in the following special cases:

2a.  $\phi_*$  surjective on fundamental groups,  $H_2(\pi_1 W) = 0$ ,  $K$  finitely presented, and  $\phi_*(M) \in H_n(W, Z)$  a primitive class. (e.g.  $\pi_1 W = \{e\}$  and  $\phi_*[M]$  primitive. Recall that primitive means of infinite order and indivisible.)

2b.  $\pi_1 M = \{e\}$ ,  $K$  finitely presented, there exists  $\xi \in H^n(W; Z\pi)$  or  $H^n(W, \partial W; Z\pi)$  if  $\partial M \neq \emptyset$ , with  $\langle \xi, \phi_*[M] \rangle = 1 \in Z\pi$ .

To prove theorem 2, a result of independent interest on  $\Gamma$ -groups is also needed. Let  $G \rightarrow \pi$  be a surjective homomorphism with kernel  $K$ , and let  $\mathcal{F}: ZG \rightarrow Z\pi$  be the map induced on integral group rings. Let  $\Gamma_n^e(\mathcal{F})$ ,  $e = s, h$  be the homology surgery groups of [CS1], and let

$$j_*: \Gamma_n^e(\mathcal{F}) \rightarrow L_n^e(\pi)$$

always denote the natural map. The next result can be found in a paper of Hausmann [H1].

**THEOREM 3.** *Suppose  $K$  is normally generated in  $G$  by a finitely generated subgroup  $N$  with  $N_{ab} = 0$ . Then*

$$j_*: \Gamma_n^e(\mathcal{F}) \rightarrow L_n^e(\pi)$$

*is an isomorphism.*

A purely algebraic proof of this result has been given recently by Justin Smith.

Using similar ideas, we also derive a new result on geometrically realizing codimension two Poincaré embeddings by P.L. embeddings (not necessarily locally flat.) Let  $\Theta$  be a Poincaré or  $h$ -Poincaré embedding (see [CS2, §5] for the definitions) of the compact oriented closed P.L. manifold  $M^n$  in the compact P.L. manifold  $W^{n+2}$ . In [CS2] it was shown that if  $n \geq 3$  is odd or if  $\pi_1 W = \{e\}$  and  $\Theta$  is cyclic, then  $\Theta$  can be realized by a P.L. embedding. Let

$$A_\Theta = Z \otimes_{Z\pi_1 M} Z\pi_1 W / \{x \cdot [M] / x \in \pi_2 W\}.$$



**THEOREM 4.** *Suppose that  $A_\Theta = 0$ . Let  $n \geq 4$  be even. Assume that the kernel of the natural map  $\pi_1 E_\Theta \rightarrow \pi_1 W$ ,  $E_\Theta$  the Poincaré complement [CS2], is finitely generated. Then  $\Theta$  can be realized by a P.L. embedding of  $M^n$  in  $W^{n+2}$ .*

Under the hypotheses of Theorem 4, the theory of [CS2] applies to study the singularities of embeddings realizing  $\Theta$ . In particular, the singularities will carry the Poincaré duals of the total  $L$ -class  $L(\Theta)$  (see [CS2, §6]).

The same assumptions as in special case 2a and 2b of theorem 2 also guarantee that the hypotheses of theorem 4 are satisfied. For example, essentially as a special case of Theorem 4, we have

**THEOREM 5.** (Compare [CS3], [CS1, §6]). *Let  $h: M^n \rightarrow W^{n+2}$ ,  $n \geq 4$  even, be a homotopy equivalence,  $M^n$  a closed oriented P.L. manifold,  $W$  a compact oriented P.L. manifold. Suppose  $H_2(\pi_1 W; \mathbb{Z}) = 0$ . Assume that the Euler class  $\chi(W) \in H^2(M)$  is a primitive generator. Then  $h$  is homotopic to a P.L. embedding.*

Recall that  $\chi(W)$  is defined in [CS2] as follows: restrict the Poincaré dual of  $h_*[M]$  to  $H^2(W) \cong_{h^*} H^2(M)$ .

In an appendix, we use Theorem 1 to give the proof of Theorem 5.1 of [CS2].

## 1. Proof of Theorem 1.

Let  $R$  be a regular neighborhood of  $\phi(M)$  that meets the boundary regularly [H3]. Let  $\pi = \pi_1 W$ . Then

$$K_{ab} = H_1(W - R; \mathbb{Z}\pi) = H_1(\widetilde{W - R})$$

$\widetilde{W - R}$  the covering space associated to  $K$ . The following sequence is exact:

$$\pi_2 W = H_2(W; \mathbb{Z}\pi) \rightarrow H_2(W, W - R; \mathbb{Z}\pi) \rightarrow H_1(W - R; \mathbb{Z}\pi) \rightarrow H_1(W; \mathbb{Z}\pi),$$

and  $H_1(W; \mathbb{Z}\pi) = H_1(\hat{W}) = 0$ ,  $\hat{W}$  the universal cover of  $W$ .

By excision  $H_2(W, W - R; \mathbb{Z}\pi) \cong H_2(R, \dot{R}; \mathbb{Z}\pi)$ ,  $\dot{R} = \partial R - \text{Int}(R \cap \partial W)$ . Here  $\mathbb{Z}\pi$  is a  $\pi_1 R = \pi_1 M$  module via the inclusion induced map. By Poincaré duality

$$H_2(R, \dot{R}; \mathbb{Z}\pi) \cong H^n(R, R \cap \partial W; \mathbb{Z}\pi).$$

But the inclusion  $(M, \partial M) \subset (R, \dot{R} \cap \partial W)$  is a (simple) homotopy equivalence. Therefore, applying Poincaré duality of  $M$ , we finally get an isomorphism  $H_2(W, W - R; \mathbb{Z}\pi) \cong H_0(M, \mathbb{Z}\pi)$ .

Now,  $H_0(M; Z\pi) = H_0(\tilde{M})$ , where  $\tilde{M}$  is the covering space induced via  $\phi$  from the universal covering space of  $W$ . From this, it is not hard to see that  $H_0(M; Z\pi) \cong Z \otimes_{Z\pi_1 M} Z\pi_1 W$ , as  $Z\pi$  modules. Further, well known standard arguments show that if  $\phi(M)$  is a smooth or P.L. locally flat submanifold, (so that  $R$  is the total space of a bundle over  $M$ ), then the map  $\pi_2 W \rightarrow H_2(W, W - R; Z\pi) \cong Z \otimes_{Z\pi_1 M} Z\pi$  is given by taking the (geometric) intersection number with  $\phi_*[M]$ . In general, we take this as the definition of  $x \circ \phi_*[M]$ ,  $x \in \pi_2 W$ . Theorem 1 now follows.

Clearly a similar argument works for a Poincaré embedding  $\Theta$  of  $M$  in  $W$ . More generally, suppose  $M = \bigcup_{i=1}^t M_i$ ,  $M_i$  components of  $M$ , but  $W$  still connected. Dropping orientability hypotheses, let  $Z_{\Theta_i}$  be the  $Z\pi_1 M_i$  module consisting of the integers with

$$g \cdot t = \omega_W(\phi_* g) \omega_{M_i}(g) t,$$

where  $g \in \pi_1 M_i$ ,  $t \in Z$ , and  $\omega_W: \pi_1 W \rightarrow \{\pm 1\}$ ,  $\omega_{M_i}: \pi_1 M_i \rightarrow \{\pm 1\}$  are the orientation characters. Let  $K$  be as in Theorem 1. Then one easily extends the above argument to show

**PROPOSITION 1.**  $K_{ab} = \bigoplus_{i=1}^t (Z_{\Theta_i} \otimes_{Z\pi_1 M_i} Z\pi_1 W / (\pi_2 W) \circ \phi_*[M_i])$ .

*Some examples:* 1.  $M$  a point in a 2-manifold  $W$  other than  $S^2$  or  $P^2$ . Then  $\pi_1(W - M)$  is a free group  $F$ , and  $\pi_2 W = 0$ . Hence we have the presentation

$$1 \rightarrow K \rightarrow F \rightarrow \pi \rightarrow 1,$$

with  $K_{ab} \cong Z\pi$ .

2. If  $W$  is a prime orientable 3-manifold (so that  $\pi_2 W = 0$  by the sphere theorem), and if  $S^1 \subset W$  represents  $\alpha \in \pi_1 W$ , then

$$K_{ab} \cong Z \otimes_{Z[\alpha]} Z\pi_1 W,$$

$K$  the kernel of  $\pi_1(W - S^1) \rightarrow \pi_1 W$ . In particular, if  $\alpha = 0$ ,  $K_{ab} \cong Z\pi_1 W$ .

3. Let  $k$  be a smooth knotted circle in  $S^3$ , and let  $l$  be a longitude and  $m$  a meridian; e.g.  $l$  is obtained by pushing  $k$  off itself slightly and linking it with linking number zero. Let  $W$  be obtained from  $S^3$  by zero-framed surgery on  $k$ , i.e. surgery killing  $l$ . Let  $G = \pi_1(S^3 - k)$ , and let  $N(l)$  be the normal subgroup of  $G$  generated by the class of  $l$ . Since  $l$  lies on a Seifert surface of  $k$  and near its

boundary,  $N(l) \subset G' = [G, G]$ . Clearly  $\pi_1 W = G/N(l)$ . Thus if  $\pi_1 W = Z$ ,  $N(l) = G'$  and so  $G'' = G' = N(l) = N(l)'$ .

Conversely suppose  $N(l) \subseteq N(l)'$ . Then by Theorem 1,  $0 = Z \otimes_{Z[m]} Z\pi_1 W / \pi_2 W \cdot [m]$ ,  $[m] \in \pi_1 W$  represented by  $m$ . Applying the sphere theorem, it follows that  $W = S^1 \times S^2 \neq Q$ . Since  $G$ , and hence  $\pi_1 W$ , is normally generated by  $[m]$ , we must have  $\pi_1 Q = \{e\}$ . Thus we have the following (possibly known) result

**PROPOSITION.** *Let  $k \subset S^3$  be a knot. Then the following are equivalent:*

1.  $\pi_1 W = Z$ ,  $W$  obtained by framed (necessarily zero-) surgery on  $k$ .
2.  $N(l) = N(l)'$ ,  $l$  a longitude.
3.  $N(l) = G'$ ,  $G = \pi_1(S^3 - k)$ .

From this proposition it follows in particular that if  $W = S^1 \times S^2$ , then the Alexander polynomial of  $k$  is identically 1. (See [Mo].)

## 2. Proof of Theorems 2, 4, and 5.

First consider theorem 2. It is known and not hard to prove that if  $h$  is (simply) splittable, then  $\sum_e(h) = 0$ ,  $e = s$  or  $h$  as appropriate. (See e.g. [B1], [CS1]) Conversely suppose  $\sum_e(h) = 0$ . Let  $\Phi$  be the diagram

$$\begin{array}{ccc} Z\pi_1(W - P) & \xrightarrow{\mathcal{F}} & Z\pi_1 W \\ \downarrow & & \downarrow \\ Z\pi_1 W & \longrightarrow & Z\pi_1 W, \end{array}$$

consisting of identities and inclusion induced maps. According to Theorem 8.5 of [CS1],  $h$  is splittable if and only if an obstruction in a quotient of  $\Gamma_{n+3}^e(\Phi)$  vanishes.<sup>(4)</sup>

We have the exact sequence [CS1, §3]:

$$\Gamma_{n+3}^e(\mathcal{F}) \rightarrow L_{n+3}^e(\pi_1 W) \rightarrow \Gamma_{n+3}^e(\Phi) \rightarrow \Gamma_{n+2}^e(\mathcal{F}) \rightarrow L_{n+2}^e(\pi_1 W).$$

---

<sup>4</sup> Actually in §8 of [CS1] we supposed  $h$  induces a (simple) homotopy equivalence of boundaries  $\partial Q$  and  $\partial W$ . However, everything in §8 goes through without change if  $h$  induces a (simple)  $Z\pi_1 W$ -homology equivalence of boundaries.

Let  $K$  be the kernel of  $\pi_1(W-P) \rightarrow \pi_1(W)$ . By hypothesis,  $K$  is finitely presented. By Theorem 1,

$$K_{ab} = Z \otimes_{Z\pi_1 P} Z\pi_1 W / \{x \cdot [P] / x \in \pi_2 W\}; \quad i.e.$$

$$K_{ab} = \text{coker } \tau_P.$$

(Here  $[P] = \phi_*[M]$ .) Thus  $K_{ab} = 0$ . Therefore, by Theorem 3, the two extreme maps in the above sequence are isomorphisms, and hence  $\Gamma_{n+3}^e(\Phi)$  vanishes. Therefore, by 8.5 of [CS1],  $h$  is (simply) splittable.

The special case 2a. follows easily. For if  $\pi_1 P \rightarrow \pi_1 W$  is surjective, then  $Z \otimes_{Z\pi_1 P} Z\pi_1 W = Z$ . If  $[P] = \phi_*[M]$  is primitive, then by Poincaré duality and universal coefficients, there is a class  $\bar{x} \in H_2 W$  with  $\bar{x} \cdot [P] = 1$ . Since  $H_2(\pi_1 W) = 0$ , the Hurewicz homomorphism  $\pi_2 W \rightarrow H_2 W$  is surjective. Therefore  $\tau_P$  is surjective in this case.

For 2b.,  $Z \otimes_{Z\pi_1 P} Z\pi_1 W = Z\pi_1 W$ , and  $\xi$  is Poincaré dual to  $x \in H_2(W, Z\pi) = \pi_2 W$ . By definition,

$$x \cdot \phi_*[M] = \langle \xi, \phi_*[M] \rangle = 1;$$

so again  $\tau_P$  will be surjective.

Theorem 4 is proven using Theorems 1 and 3 and the same method as for Theorem 5.3 of [CS2]. Given a Poincaré embedding  $\Theta$ , one obtains, as in the section of §5 of [CS2] just preceding lemma 5.5, an element  $\Sigma \in \Gamma_{n+2}^s(\mathcal{F}_\Theta)$ , where  $\mathcal{F}_\Theta$  is the natural map  $Z\pi_1 E_\Theta \rightarrow Z\pi_1 W$ . Using the result from Theorem 3 that  $j_*: \Gamma_{n+3}^s(\mathcal{F}_\Theta) \rightarrow L_{n+3}^s(\pi_1 W)$  is surjective, one shows that the vanishing of  $\Sigma$  implies the existence of the desired P.L. embedding (actually with any regular neighborhood with the correct normal invariant and associated bundle). This is proven using same argument as for the proof of Lemma 5.5d of [CS2]. Theorem 3 applies in present case because, by Theorem 1 (for a Poincaré embedding) and the hypothesis  $A_\Theta = 0$ , the kernel of  $\pi_1 E_\Theta \rightarrow \pi_1 W$  has trivial abelianization.

However, lemma 5.5a of [CS2] is also valid in the present situation, by the same proof. Thus  $j_*\Sigma = 0$ . Hence, by theorem 3 again,  $\Sigma = 0$ . Therefore the desired P.L. embedding exists. If  $\Theta$  was only a  $h$ -Poincaré embedding to begin with, this argument only leads to an embedding of  $M$  in a manifold  $h$ -cobordant to  $W$ . As in [CS2], one then uses the same type argument as in 8.1 of [CS1] to correct the torsion, in the complement of  $M$ , to obtain the desired embedding of  $M$  in (a manifold  $s$ -cobordant to)  $W$ .

Finally, to prove Theorem 5, recall that in 6.1 of [CS2] we saw that, under the hypotheses of Theorem 5, there is an  $h$ -Poincaré embedding  $\Theta$  of  $M$  in  $W$  whose underlying map  $M \rightarrow W$  is precisely  $h$ . Further, the natural map  $j_*: \pi_1(E_\Theta) \rightarrow \pi_1(W)$  is in this case just the composition with  $(h_*)^{-1}$  on  $\pi_1$  of  $p_*: \pi_1(S(\xi)) \rightarrow \pi_1 M$ ,  $p$  the projection of an orientable circle bundle over  $M$  with Euler class

$\chi(W) \in H^2(M)$ . In particular, the kernel  $K$  of  $i_*$  is a cyclic group. Therefore, by Theorem 1,

$$K = K_{ab} = \mathbb{Z}/\{x \cdot h_*[M]/x \in \pi_2 W\}.$$

For  $\bar{x} \in H_2 W$ , the evaluation  $\langle \chi(W)m \bar{x} \rangle$  is just the intersection number  $\bar{x} \circ h_*[M]$ . Hence, if  $\chi(W)$  is primitive, it follows by universal coefficients that  $\exists \bar{x}$  with  $\bar{x} \cdot h_*[M] = 1$ . Since  $H_2(\pi_1 W) = 0$  by assumption,  $\bar{x}$  is the image of  $x \in \pi_2 W$  under the Hurewicz map. Thus  $K = 0$ , and the result now follows from Theorem 5.3 of [CS2].

*Note:* Under the hypotheses of Theorem 5, one can use the methods of [CS2] to try to find a locally flat spine of  $W$ ; i.e. a locally flat embedding  $h': M' \rightarrow W$  that is a homotopy equivalence. It can be shown that it is possible if and only if the normal invariant  $\eta(\Theta)$  in  $[M; G/PL]$  has trivial surgery obstruction in  $L_n(\pi_1 M)$  (Compare [CS2, h. 2]). If  $\eta(\Theta)$  has trivial surgery obstruction in the reduced Wall group  $\tilde{L}_n(\pi_1 M)$ , then there exists an almost locally flat spine (i.e. locally flat except at one point.) This assertions can also be derived from Theorem 2, and should be compared to the results of [CS3] on the existence of totally spinless manifolds, as well as [Ma] and the exposition [Sh2]. For the case  $\pi_1 M = \{e\}$ , results on the existence of locally flat spines were obtained in [KM], by different methods.

## Appendix

The next result was given without proof in [CS2]. It is proven using Theorem 1.

**THEOREM (5.1 of [CS2]).** *Let  $\Theta$  be an (oriented) Poincaré or h-Poincaré embedding of  $(M^n, \partial M)$  in  $(W^{n+2}, \partial W)$ ,  $W$  connected,  $M = M_1 \cup \cdots \cup M_i$ ,  $M_i$  connected. Then there is an (oriented) cyclic Poincaré or h-Poincaré embedding of  $(M, \partial M)$  in  $(W, \partial W)$ ,  $\Theta'$  say, and a map  $\Theta \rightarrow \Theta'$ , if and only if*

$$A_\Theta = \bigoplus_{i=1}^t (Z_{\Theta_i} \otimes_{Z\pi_1 M_i} Z\pi_1 W) / \pi_2 W \cdot [M_i])$$

*is a cyclic abelian group.*

Recall that  $\Theta'$  is by definition [CS2, §5] cyclic if and only if the kernel of  $\pi_1(E_{\Theta'}) \rightarrow \pi_1(W)$ ,  $E_{\Theta'}$  the Poincaré “complement”, is cyclic. Also, see [CS2, §5] for the definition of a map  $\Theta \rightarrow \Theta'$ .

To prove this result, let  $K$  be the kernel of  $\pi_1 E_\Theta \rightarrow \pi_1 W$ . A map  $\Theta \rightarrow \Theta'$ , if it exists, restricts to a homology equivalence  $E_\Theta \rightarrow E_{\Theta'}$  with coefficients in  $Z\pi_1 W$ . Therefore

$$K_{ab} = H_1(E_\Theta, Z\pi_1 W) = H_1(E_{\Theta'}, Z\pi_1 W) = K'_{ab}.$$

Therefore, Theorem 1, applied to Poincaré embeddings shows that a cyclic  $\Theta'$  and a map  $\Theta \rightarrow \Theta'$  exist only if  $A_\Theta$  is cyclic.

Suppose that  $A_\Theta$  is cyclic. Let  $G = \pi_1 E_\Theta$ . The sequence

$$1 \rightarrow K_{ab} \rightarrow G/[K, K] \rightarrow G/K = \pi_1 W \rightarrow 1$$

is exact. (That  $G \rightarrow \pi_1 W$  is surjective follows either by general position or an argument using Van Kampen's theorem and the exact homotopy sequence of a circle bundle.) Hence, as  $K_{ab}$  is cyclic by Theorem 1 and the hypothesis on  $A_\Theta$ ,  $G/[K, K]$  is finitely presented. Therefore  $[K, K]$  is normally generated by a finite number of elements,  $\alpha_1, \dots, \alpha_r$  say.

Let  $Y \supset E_\Theta$  be obtained by attaching 2-cells along circles representing  $\alpha_1, \dots, \alpha_r$ . Then we may identify  $\pi_1 Y$  with  $G/[K, K]$ . Let  $\pi = G/K$ . We have the exact sequence

$$H_i(E_\Theta; Z\pi) \rightarrow H_i(Y, Z\pi) \rightarrow H_i(Y, E_\Theta; Z\pi) \rightarrow H_{i-1}(E_\Theta; Z\pi),$$

and

$$H_i(Y; E_\Theta; Z\pi) = \begin{cases} 0, & i \neq 2 \\ Z\pi \oplus \dots \oplus Z\pi (r \text{ summands}), & i = 2. \end{cases}$$

Further, the connecting homomorphism

$$H_2(Y, E_\Theta; Z\pi) \rightarrow H_1(E_\Theta; Z\pi) = K_{ab}$$

is trivial. Hence

$$H_2(Y; Z\pi) \cong H_2(E_\Theta; Z\pi) + (Z\pi)',$$

and the inclusion  $E_\Theta \subset Y$  induces isomorphisms of homology groups in other dimensions over  $Z\pi$ , and is in fact a simple equivalence over  $Z\pi$  with respect to the basis of  $H_2(Y; E_\Theta; Z\pi)$  determined by the 2-cells attached.

We assert that the Hurewicz homomorphism  $\pi_2 Y \rightarrow H_2(Y; Z\pi)$  is surjective. In fact  $H_2(Y; Z\pi) = H_2 \tilde{Y}$ ,  $\tilde{Y}$  the covering space of  $Y$  with  $\pi_1 \tilde{Y} = K_{ab}$ , and one has the (Hopf) exact sequence

$$\pi_2 Y = \pi_2 \tilde{Y} \rightarrow H_2 \tilde{Y} \rightarrow H_2(K_{ab}, Z) = 0.$$

Now let  $E_{\Theta'} \supset Y$  be obtained by attaching 3 cells along two-spheres representing the elements of the obvious basis of  $Z\pi \oplus \cdots \oplus Z\pi$ . The inclusion  $E_{\Theta} \rightarrow E_{\Theta'}$  will be a (simple) homology equivalence over  $Z\pi$ ,  $\pi_1 E_{\Theta} \rightarrow \pi_1 W$  is just  $G/[K, K] \rightarrow \pi_1 W$ , with kernel  $K_{ab}$ . Let  $F_{\Theta'} = F_{\Theta} \subset E_{\Theta'}$  (this is the Poincaré complement of  $\partial M$  in  $\partial W$ ; see [CS2]) and let  $h_{\Theta'}$  be the composite of  $h_{\Theta}: W \rightarrow E(\xi_{\Theta}) \cup_{s(\xi_{\Theta})} E_{\Theta}$  and the union of the identity on  $E(\xi_{\Theta})$  and the inclusion  $E_{\Theta} \subset E_{\Theta'}$ . Let  $\xi_{\Theta'} = \xi_{\Theta}$ . Then  $h_{\Theta'}$  is a (simple) homotopy equivalence, and it is not hard to see that  $\Theta' = (\xi_{\Theta'}, (E_{\Theta'}, F_{\Theta'}), h_{\Theta'})$  is a cyclic Poincaré or  $h$ -Poincaré embedding, as appropriate, with a map  $\Theta \rightarrow \Theta'$  defined by the inclusion  $E_{\Theta} \subset E_{\Theta'}$ .

## REFERENCES

- [B1] W. BROWDER, *Embedding Smooth Manifolds*, Proc. Int. Cong. of Mathematicians, Moscow, 1966; Mir 1968 (712–719).
- [B2] —, *Surgery on simply-connected manifolds*, Ergebnisse der Mathematik, Springer Verlag, 1972.
- [CS1] S. E. CAPPELL and J. L. SHANESON, *The codimension two placement problem and homology equivalent manifolds*, Annals of Math 99 (1974), 277–348.
- [CS2] —, *P.L. embeddings and their singularities*, Annals of Math. 103 (1976), 163–228.
- [CS3] —, *Totally spineless manifolds*, Illinois Journal of Mathematics, to appear.
- [H1] J. HAUSMANN, *Homological Surgery*, Ann. of Math., to appear.
- [H2] —, *Manifolds up to homology*, to appear.
- [H3] J. F. P. HUDSON, *Piecewise Linear Topology*, Benjamin, 1970.
- [KM] M. KATO and Y. MATSUMOTO, *Simply connected surgery of submanifolds in codimension two*, I. J. Math. Soc. Japan 24 (1974), 586–608.
- [Ma] Y. MATSUMOTO, *A 4-manifold which admits no spine*, Bull. Amer. Math. Soc. 81 (1975), 467–470.
- [Mo] LOUISE E. MOSER, *On the impossibility of obtaining  $S^2 \times S^1$  by elementary surgery along a knot*, Pac. J. Math. 53 (1974), 519–523.
- [ST] H. SEIFERT and W. THRELLFALL, *Lehrbuch der Topology*, Chelsea Publishing Co., Bronx, N.Y., 1947.
- [Sh1] J. L. SHANESON, *Wall's surgery obstruction groups for  $Z \times G$* , Ann. of Math. 90 (1969), 296–334.
- [Sh2] —, *Spines and Spinelessness*, in Vol. 438, Lecture Notes in Mathematics (Springer-Verlag), Geometric Topology, p. 431–439.
- [W] C. T. C. WALL, *Surgery on compact manifolds*, Academic Press, London, 1970.

*Courant Institute of Mathematical Sciences,  
Rutgers University*

Received May 21, 1976