

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 52 (1977)

Artikel: Groups with cyclic Sylow subgroups and finiteness conditions for certain complexes.
Autor: Mislin, G.
DOI: <https://doi.org/10.5169/seals-40005>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 11.07.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Groups with cyclic Sylow subgroups and finiteness conditions for certain complexes

G. MISLIN

Introduction

Let π denote a finite group of order n whose Sylow subgroups are all cyclic and let $N = \sum x \in \mathbf{Z}\pi$, $x \in \pi$, denote the norm element. The augmentation $\mathbf{Z}\pi \rightarrow \mathbf{Z}$ induces a map $j: \mathbf{Z}\pi/N \rightarrow \mathbf{Z}/n$ which we use to consider \mathbf{Z}/n as a $\mathbf{Z}\pi/N$ -module. We show (Theorem 1.3) that

$$\text{proj. dim}_{\mathbf{Z}\pi/N} (\mathbf{Z}/n) < \infty.$$

Thus there is a *transfer map*

$$j^*: K_0(\mathbf{Z}/n) \rightarrow K_0(\mathbf{Z}\pi/N)$$

between projective class groups. It turns out that $\text{im}(j^*) \subset \text{im}(pr_*: K_0(\mathbf{Z}\pi) \rightarrow K_0(\mathbf{Z}\pi/N))$ and, since $\text{im}(K_0(\mathbf{Z}\pi) \rightarrow K_0(\mathbf{Z}\pi/N)) \cong (K_0\mathbf{Z}\pi)/\text{im } S$ where $S: K_1(\mathbf{Z}/n) \rightarrow K_0(\mathbf{Z}\pi)$ denotes the *Swan homomorphism* (cf. Section 2), we can think of the transfer map to map $K_0(\mathbf{Z}/n)$ into $(K_0\mathbf{Z}\pi)/\text{im } S$. If we compose this map with the obvious homomorphism

$$u\left(\mathbf{Z}\left[\frac{1}{n}\right]\right) \rightarrow K_0(\mathbf{Z}/n)$$

we obtain a “transfer” homomorphism

$$T: u\left(\mathbf{Z}\left[\frac{1}{n}\right]\right) \rightarrow (K_0\mathbf{Z}\pi)/\text{im } S$$

($u(\mathbf{Z}[1/n])$ denotes the group of units of $\mathbf{Z}[1/n]$). The homomorphism T is in general non-trivial, even if π is cyclic (in which case $\text{im } S = 0$). However, we

show that $T = 0$ if n is a prime power or if $n = 2p$, p an odd prime (cf. Theorem 2.5).

In the second half of the paper we make use of the homomorphism T to compute the Wall obstruction $wX \in K_0\mathbb{Z}\pi_1 X$ for certain complexes. We will consider spaces X for which $\pi_1 X$ operates nilpotently on $H_*\tilde{X}$ (i.e. X is homologically nilpotent in the sense of Brown–Kahn [3]). If such a space is dominated by a finite complex and has a finite fundamental group of order n , then the rational number

$$\rho(X) = \text{card } H_{\text{odd}}(X, \tilde{X}) / \text{card } H_{\text{ev}}(X, \tilde{X})$$

is well defined and is a unit in $\mathbb{Z}[1/n]$; $\rho(X)$ is related to the finiteness obstruction wX in the following way, (cf. Theorem 3.3).

THEOREM I. *Let X be a finitely dominated homologically nilpotent space with non-zero finite fundamental group of square free order. Then*

$$T\rho(X) = \bar{w}X$$

where $\bar{w}X$ denotes the image of wX in $(K_0\mathbb{Z}\pi_1 X)/\text{im } S$.

In particular, if the space X in Theorem I is supposed to be *nilpotent*, then $\pi_1 X$ —being nilpotent and of square free order—is necessarily cyclic and therefore $\text{im } S = 0$ by a result of Swan [14]. The formula reduces then to

$$T\rho(X) = wX$$

yielding new information concerning the Wall obstruction for nilpotent spaces.

Under suitable conditions on X the rational number $\rho(X)$ depends only upon H_*X : Suppose that $\pi_1 X$ is cyclic of square free order n operating trivially on $H_*^\pi(X; I\mathbb{Z}\pi)$. Then we show that

$$\rho(X) = \prod_{p|n} p^{e_p(X)}$$

the product being taken over all prime divisors of n , and $e_p(X)$ denoting the value at -1 of the derivative of the *Poincaré polynomial* of X with respect to \mathbb{Z}/p -coefficients, a quantity depending only upon H_*X .

As an illustration we show that for X an H -space of rank ≥ 2 one has $e_p(X) = 0$ for all primes p , and hence $\rho(X) = 1$. The following vanishing theorem for the Wall obstruction for H -spaces then follows.

THEOREM II. *Let X be a finitely dominated H -complex with finite fundamental group of square free order. Then $wX=0$ and X is therefore of the homotopy type of a finite complex.*

1. Groups with cyclic Sylow subgroups and $\mathbb{Z}\pi/N$ -modules

Let π denote a finite group whose p -Sylow subgroups are cyclic of order p^k for a fixed prime p . Such a group π is p -periodic in the sense of Cartan–Eilenberg [4]. If q denotes the smallest p -period of π , then $H^q(\pi; \mathbb{Z}_{(p)}) \cong \mathbb{Z}/p^k$, where $\mathbb{Z}_{(p)}$ denotes the integers localized at p . Furthermore, if $H^i(\pi; \mathbb{Z}_{(p)}) \cong \mathbb{Z}/p^k$ for some $i > 0$, then i is necessarily a multiple of q (see Swan [15]). It has been observed by Lundmark [8] that

$$H^i(\pi; \mathbb{Z}_{(p)}) = 0 \quad \text{for } 0 < i < q.$$

Namely, suppose i is an integer with $0 < i < q$ and let π_p denote a p -Sylow subgroup of π . Then from the decomposition

$$H^i(\pi_p; \mathbb{Z}) \cong \text{im } \iota(\pi_p, \pi) \oplus \ker t(\pi, \pi_p)$$

(cf. [4]) and the fact that the map induced by inclusion $\iota(\pi_p, \pi): H^i(\pi; \mathbb{Z}) \rightarrow H^i(\pi_p; \mathbb{Z})$ is monic on the p -primary subgroup, we infer, because π_p is cyclic, that $H^i(\pi; \mathbb{Z}_{(p)}) \cong \mathbb{Z}/p^k$ or $H^i(\pi; \mathbb{Z}_{(p)}) = 0$. The former case is impossible since i is not a multiple of q and hence $H^i(\pi; \mathbb{Z}_{(p)}) = 0$ for $0 < i < q$.

Let π be an arbitrary finite group of order n and $N = \sum x \in \mathbb{Z}\pi$, $x \in \pi$. Then

$$\begin{array}{ccc} \mathbb{Z}\pi & \longrightarrow & \mathbb{Z}\pi/N \\ \downarrow & & \downarrow \\ \mathbb{Z} & \longrightarrow & \mathbb{Z}/n \end{array}$$

is a pullback square of rings (with obvious maps). Hence there is a short exact sequence of $\mathbb{Z}\pi/N$ -modules

$$0 \rightarrow I\mathbb{Z}\pi \rightarrow \mathbb{Z}\pi/N \rightarrow \mathbb{Z}/n \rightarrow 0$$

where $I\mathbb{Z}\pi$ denotes the augmentation ideal. Notice that a $\mathbb{Z}\pi/N$ -module may be considered as a π -module via the projection $\mathbb{Z}\pi \rightarrow \mathbb{Z}\pi/N$.

DEFINITION 1.1. A $\mathbf{Z}\pi/N$ -module M is said to be *trivial*, if it is trivial as a π -module; M is called *nilpotent*, if M possesses a finite filtration with associated graded module a trivial $\mathbf{Z}\pi/N$ -module.

If M is a $\mathbf{Z}\pi/N$ -module, then we will write IM for $(I\mathbf{Z}\pi)M$ and $I^k M$ for $I(I^{k-1}M)$, $k \geq 2$. Obviously, M is then nilpotent if and only if $I^k M = 0$ for some k , (if and only if M is nilpotent as a π -module, respectively). Furthermore, M is a trivial $\mathbf{Z}\pi/N$ -module if and only if $IM = 0$; hence a trivial $\mathbf{Z}\pi/N$ -module is the same as a \mathbf{Z}/n -module. It is plain that the underlying abelian group of a nilpotent $\mathbf{Z}\pi/N$ -module is an n -torsion group.

LEMMA 1.2. Let π denote a finite group whose p -Sylow subgroups are cyclic of order p^k , p a fixed prime. Then, for \mathbf{Z}/p^k considered as a trivial $\mathbf{Z}\pi/N$ -module

$$\text{proj. dim}_{\mathbf{Z}\pi/N} (\mathbf{Z}/p^k) \leq q$$

where q denotes the minimal p -period of π .

Proof. By [14] there exists a periodic resolution

$$\cdots \rightarrow P_i \rightarrow P_{i-1} \rightarrow \cdots \rightarrow P_0 \rightarrow \mathbf{Z}_{(p)} \rightarrow 0$$

with P_i projective $\mathbf{Z}_{(p)}\pi$ -modules, $P_i = P_{i+q}$ and $P_q \rightarrow P_{q-1}$ factoring through $\mathbf{Z}_{(p)}$. Let $\Lambda = \mathbf{Z}\pi/N$ and $\Lambda_p = \Lambda \otimes \mathbf{Z}_{(p)}$. From the short exact sequence $\mathbf{Z}_{(p)} \xrightarrow{N} \mathbf{Z}_{(p)}\pi \rightarrow \Lambda_p$ we deduce $H_i(\pi; \Lambda_p) \cong H_{i-1}(\pi; \mathbf{Z}_{(p)})$ for $i \geq 2$, and an exact sequence

$$0 \rightarrow H_1(\pi; \Lambda_p) \rightarrow \mathbf{Z}_{(p)} \xrightarrow{n} \mathbf{Z}_{(p)} \rightarrow H_0(\pi; \Lambda_p) \rightarrow 0.$$

Since $H_i(\pi; \mathbf{Z}_{(p)}) = 0$ for $0 < i < q-1$ we conclude that

$$H_i(\pi; \Lambda_p) = \begin{cases} \mathbf{Z}/p^k & \text{if } i = 0, q \\ 0 & \text{if } 0 < i < q. \end{cases}$$

These groups are the homology groups of the complex $\cdots \rightarrow Q_i \rightarrow Q_{i-1} \rightarrow \cdots \rightarrow Q_0 \rightarrow 0$ in dimension $\leq q$, where $Q_i = \Lambda_p \otimes_{\pi} P_i$. Notice that Q_i is torsionfree as an abelian group, since it is Λ_p -projective. We know that $d_q: Q_q \rightarrow Q_{q-1}$ factors through $\Lambda_p \otimes_{\pi} \mathbf{Z}_{(p)} \cong \mathbf{Z}/p^k$ and therefore, since $\text{im}(d_q)$ is a torsionfree abelian

group, we infer $d_q = 0$. Thus

$$0 \rightarrow Q_{q-1} \rightarrow Q_{q-2} \rightarrow \cdots \rightarrow Q_0 \twoheadrightarrow \mathbf{Z}/p^k$$

is a projective resolution of the trivial Λ_p -module \mathbf{Z}/p^k . As a result $\text{proj. dim}_{\Lambda_p}(\mathbf{Z}/p^k) \leq q-1$. Of course $\text{proj. dim}_{\Lambda}(\Lambda_p) = 1$, as one can see by tensoring a free abelian presentation of $\mathbf{Z}_{(p)}$ with Λ . As a consequence

$$\text{proj. dim}_{\Lambda}(\mathbf{Z}/p^k) \leq \text{proj. dim}_{\Lambda_p}(\mathbf{Z}/p^k) + \text{proj. dim}_{\Lambda}(\Lambda_p) \leq q$$

which completes the proof of the lemma.

An immediate consequence is the following theorem which was mentioned in the introduction.

THEOREM 1.3. *Suppose π is a finite group of order n with cyclic Sylow subgroups. Then \mathbf{Z}/n considered as a trivial $\mathbf{Z}\pi/N$ -module has finite projective dimension.*

Proof. Write $\mathbf{Z}/n = \bigoplus \mathbf{Z}/p^{k(p)}$, the sum taken over all prime divisors of n . Then

$$\text{proj. dim}_{\mathbf{Z}\pi/N}(\mathbf{Z}/n) = \max(\text{proj. dim}_{\mathbf{Z}\pi/N}(\mathbf{Z}/p^{k(p)}) \mid p|n) < \infty$$

Remark. From the short exact sequence $I\mathbf{Z}\pi \rightarrow \mathbf{Z}\pi/N \rightarrow \mathbf{Z}/n$ we see that $\text{proj. dim}_{\mathbf{Z}\pi/N}(I\mathbf{Z}\pi) = \text{proj. dim}_{\mathbf{Z}\pi/N}(\mathbf{Z}/n) - 1$. Hence, if π has cyclic Sylow subgroups, we get from Theorem 1.3

$$\text{proj. dim}_{\mathbf{Z}\pi/N}(I\mathbf{Z}\pi) < \infty$$

This generalizes a well known fact on the augmentation ideal of a finite cyclic group, in which case $I\mathbf{Z}\pi$ is free of rank 1 over $\mathbf{Z}\pi/N$.

We will apply later Lemma 1.2 and Theorem 1.3 in case π has square free order; for such a π the Sylow subgroups are of course cyclic of prime order.

LEMMA 1.4. *Let π be a finite group of square free order n and let M denote a nilpotent $\mathbf{Z}\pi/N$ -module. Then*

$$(i) \text{ proj. dim}_{\mathbf{Z}\pi/N}(M) < \infty;$$

if, in addition, M is finitely generated, then

(ii) M is of type FP and $\text{card}(M)$ is a unit in $\mathbf{Z}[1/n]$.

Proof. We first assume that M is a trivial $\mathbf{Z}\pi/N$ -module. Then M is a direct sum of modules of the form \mathbf{Z}/p , p dividing n . From Lemma 1.2 we see then that $\text{proj. dim } M < \infty$. If M is a general nilpotent $\mathbf{Z}\pi/N$ -module, we choose a finite filtration of M such that $\text{gr}(M)$ is a trivial $\mathbf{Z}\pi/N$ -module. Clearly $\text{proj. dim } \text{gr}(M) \geq \text{proj. dim } M$ and *i*) follows. If M is finitely generated then, $\mathbf{Z}\pi/N$ being noetherian, we can find a projective resolution of M of finite length, which is also of finite type; by definition, M is therefore of type FP . Finally, a finitely generated nilpotent $\mathbf{Z}\pi/N$ -module has as underlying abelian group a finitely generated n -torsion group. Hence $\text{card}(M)$ is a unit in $\mathbf{Z}[1/n]$.

2. The transfer homomorphism $T: u(\mathbf{Z}[1/n]) \rightarrow (K_0\mathbf{Z}\pi)/\text{im } S$

Let π denote a finite group of order n with cyclic Sylow subgroups. Then according to Theorem 1.3, $\text{proj. dim}_{\mathbf{Z}\pi/N}(\mathbf{Z}/n) < \infty$, and therefore the canonical projection $j: \mathbf{Z}\pi/N \rightarrow \mathbf{Z}/n$ gives rise to a transfer map (cf. Bass [1, Chapter IX, 1.7])

$$j^*: K_0(\mathbf{Z}/n) \rightarrow K_0(\mathbf{Z}\pi/N).$$

The map j^* is defined on a generator $[\mathbf{Z}/p^k]$ of $K_0(\mathbf{Z}/n)$ by choosing a $\mathbf{Z}\pi/N$ -projective resolution of finite type

$$0 \rightarrow P_m \rightarrow P_{m-1} \rightarrow \cdots \rightarrow P_0 \rightarrow \mathbf{Z}/p^k \rightarrow 0$$

of the trivial $\mathbf{Z}\pi/N$ -module \mathbf{Z}/p^k , and setting

$$j^*[\mathbf{Z}/p^k] = \sum (-1)^i [P_i] \in K_0(\mathbf{Z}\pi/N).$$

Let $j_*: K_0(\mathbf{Z}\pi/N) \rightarrow K_0(\mathbf{Z}/n)$ denote the map induced by the projection $j: \mathbf{Z}\pi/N \rightarrow \mathbf{Z}/n$.

LEMMA 2.1. $j_*j^*: K_0(\mathbf{Z}/n) \rightarrow K_0(\mathbf{Z}/n)$ is the 0-homomorphism.

Proof. Let q denote the minimal p -periode of π and let $[\mathbf{Z}/p^k] \in K_0(\mathbf{Z}/n)$ denote a generator. Choose a $\mathbf{Z}\pi/N$ -projective resolution of finite type of \mathbf{Z}/p^k

which has length q (cf. Lemma 1.2)

$$0 \rightarrow L_q \rightarrow L_{q-1} \rightarrow \cdots \rightarrow L_0 \rightarrow \mathbf{Z}/p^k \rightarrow 0.$$

Then

$$\begin{aligned} j_* j^* [\mathbf{Z}/p^k] &= j_* \left(\sum (-1)^i [L_i] \right) \\ &= \sum (-1)^i [\mathbf{Z}/n \otimes_{\pi} L_i] \\ &= \sum_{r|n} \left(\sum (-1)^i [\mathbf{Z}/n(r) \otimes_{\pi} L_i] \right) \end{aligned}$$

where $n(r)$ stands for the highest power of the prime r , which divides n . For $r \neq p$ we have

$$\mathrm{Tor}_{\mathbf{Z}\pi/N}^* (\mathbf{Z}/n(r), \mathbf{Z}/p^k) = 0$$

and therefore the complex

$$0 \rightarrow \mathbf{Z}/n(r) \otimes_{\pi} L_q \rightarrow \cdots \rightarrow \mathbf{Z}/n(r) \otimes_{\pi} L_0 \rightarrow 0$$

is exact. Hence $\sum (-1)^i [\mathbf{Z}/n(r) \otimes_{\pi} L_i] = 0$ for $r \neq p$, and therefore $j_* j^* [\mathbf{Z}/p^k] = \sum (-1)^i [\mathbf{Z}/p^k \otimes_{\pi} L_i]$. To compute $\sum (-1)^i [\mathbf{Z}/p^k \otimes_{\pi} L_i]$ and the homology of $\{\mathbf{Z}/p^k \otimes_{\pi} L_i\}$ we can as well use the $\mathbf{Z}\pi/N \otimes \mathbf{Z}_{(p)}$ -projective resolution $\{Q_i\}$ of \mathbf{Z}/p^k , which was considered in the proof of Lemma 1.2. Hence

$$\sum (-1)^i [\mathbf{Z}/p^k \otimes_{\pi} L_i] = \sum_{i=0}^{q-1} (-1)^i [\mathbf{Z}/p^k \otimes_{\pi} Q_i]$$

and plainly for $0 \leq i \leq q-1$ one has

$$\mathrm{Tor}_{\mathbf{Z}\pi/N}^i (\mathbf{Z}/p^k, \mathbf{Z}/p^k) = H_i(\pi; \mathbf{Z}/p^k) = \begin{cases} 0 & \text{for } 0 < i < q-1 \\ \mathbf{Z}/p^k & \text{for } i = 0, q-1 \end{cases}$$

Therefore $j_* j^* [\mathbf{Z}/p^k] = [\mathbf{Z}/p^k] + (-1)^{q-1} [\mathbf{Z}/p^k] = 0$ because the p -period q of π is an even number [15].

If π denotes an arbitrary group of order n then associated with the square of rings

$$\begin{array}{ccc} \mathbf{Z}\pi & \xrightarrow{pr} & \mathbf{Z}\pi/N \\ \downarrow & & \downarrow j \\ \mathbf{Z} & \longrightarrow & \mathbf{Z}/n \end{array}$$

there is an exact sequence (cf. Milnor [9]) which reduces to

$$u(\mathbf{Z}/n) \xrightarrow{S} K_0(\mathbf{Z}\pi) \xrightarrow{pr_*} K_0(\mathbf{Z}\pi/N) \xrightarrow{j_*} \tilde{K}_0(\mathbf{Z}/n) \rightarrow 0 \quad (2.2)$$

We call S the *Swan homomorphism* (cf. [14]). S can be described in the following way: for k a unit mod n , $S(k) = [(k, N)]$ where (k, N) denotes the projective ideal in $\mathbf{Z}\pi$ generated by k and N .

Consider now the case of a π with cyclic Sylow subgroups. Then $j_*j^* = 0$ by Lemma 2.1 and, by the exactness of (2.2), the transfer j^* gives therefore rise to a homomorphism

$$t: K_0(\mathbf{Z}/n) \rightarrow (K_0\mathbf{Z}\pi)/\text{im } S$$

such that $\bar{pr}_*t = j^*$, $\bar{pr}_*: (K_0\mathbf{Z}\pi)/\text{im } S \rightarrow K_0(\mathbf{Z}\pi/N)$ denoting the map induced by pr_* .

If $n = p_1^{k_1} \cdots p_m^{k_m}$ then $K_0(\mathbf{Z}/n)$ is a free abelian group, freely generated by $\{[\mathbf{Z}/p_i^{k_i}], 1 \leq i \leq m\}$. Hence there is a unique group homomorphism

$$\varphi: u\left(\mathbf{Z}\left[\frac{1}{n}\right]\right) \rightarrow K_0(\mathbf{Z}/n)$$

such that $\varphi(\pm p_i) = [\mathbf{Z}/p_i^{k_i}]$. If we compose φ with t we get a map $T = t\varphi$ which we will also call a *transfer*, since it is induced by j^* . For π a group with cyclic Sylow subgroups we get therefore a commutative diagram

$$\begin{array}{ccc} u(\mathbf{Z}[1/n]) & \xrightarrow{T} & (K_0\mathbf{Z}\pi)/\text{im } S \\ \downarrow \varphi & \nearrow t & \downarrow \bar{pr}_* \\ K_0(\mathbf{Z}/n) & \xrightarrow{j^*} & K_0(\mathbf{Z}\pi/N) \end{array} \quad (2.3)$$

We will sometimes consider $K_0(\mathbf{Z}\pi/N)$ to be the range of T ; this should not give rise to any confusion, since \bar{pr}_* is injective.

It is well known that if R is a ring and M an R -module of type FP , then M defines an element $[M] \in K_0R$ (depending only upon the isomorphism class of M) by choosing any finite projective resolution of finite type

$$0 \rightarrow P_m \rightarrow P_{m-1} \rightarrow \cdots \rightarrow P_0 \rightarrow M \rightarrow 0$$

and setting $[M] = \sum (-1)^i [P_i] \in K_0 R$; if $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ is an exact sequence of modules of type FP , then $[M] = [M'] + [M'']$ (cf. [1] and [11]).

LEMMA 2.4. *Let π denote a finite group of square free order n and let M denote a finitely generated nilpotent $\mathbf{Z}\pi/N$ -module. Then*

$$T(\text{card } M) = [M] \in K_0(\mathbf{Z}\pi/N).$$

Proof. Notice that M is of type FP over $\mathbf{Z}\pi/N$ and $\text{card } M \in u(\mathbf{Z}[1/n])$ by Lemma 1.4. Hence $T(\text{card } M)$ and $[M]$ are well defined elements of $K_0(\mathbf{Z}\pi/N)$. If M is a trivial $\mathbf{Z}\pi/N$ -module, then $T(\text{card } M) = j^* \varphi(\text{card } M) = [M]$ where the second equation follows from the definition of φ , j^* and $[M]$ respectively. For the general case we choose a finite filtration of M with $\text{gr}(M)$ a trivial $\mathbf{Z}\pi/N$ -module. Clearly $\text{card } M = \text{card } \text{gr}(M)$ and $[M] = [\text{gr}(M)]$; therefore $T(\text{card } M) = [M]$.

For the applications in the next section we will be particularly interested in groups π for which $\text{im } S = 0$. The following theorem gives some information on T for such cases.

THEOREM 2.5. *Let π denote a finite group of order n with cyclic Sylow subgroups. Then*

- (i) $T \equiv 0$ in case n is a prime power or $n = 2p$, p an odd prime.
- (ii) $T(p_1 \cdots p_m) = 0$ if π is cyclic of order $p_1^{k_1} \cdots p_m^{k_m}$.
- (iii) $T(3) = T(5) \neq 0$ if $n = 15$, and $T(3)$ has order 2.

Furthermore, in all three cases listed above one has $\text{im } S = 0$, and T can therefore be considered as a map $T: u(\mathbf{Z}[1/n]) \rightarrow K_0 \mathbf{Z}\pi$.

We will break the proof up into a couple of lemmas.

LEMMA 2.6. *Let π be a cyclic group of order n . Then $j^*: K_0(\mathbf{Z}/n) \rightarrow K(\mathbf{Z}\pi/N)$ factors through $\tilde{K}_0(\mathbf{Z}/n)$.*

Proof. We may assume $n > 1$. Let x denote a generator of π . Then $I\mathbf{Z}\pi$ is freely generated by $(1-x)$ over $\mathbf{Z}\pi/N$ and hence there is an exact sequence $0 \rightarrow \mathbf{Z}\pi/N \rightarrow \mathbf{Z}\pi/N \rightarrow \mathbf{Z}/n \rightarrow 0$, from which we infer that $j^*[\mathbf{Z}/n] = [\mathbf{Z}\pi/N] - [\mathbf{Z}\pi/N] = 0$. Thus j^* factors through $\tilde{K}_0(\mathbf{Z}/n) = K_0(\mathbf{Z}/n)/\langle [\mathbf{Z}/n] \rangle$.

LEMMA 2.7. *Let p denote an odd prime and $\pi = \mathbf{Z}/2p$ or the dihedral group D_{2p} . Then*

$$j^* = 0: K_0(\mathbf{Z}/2p) \rightarrow K_0(\mathbf{Z}\pi/N)$$

Proof. We will first consider the case $\pi = \mathbf{Z}/2p$. Since j^* factors through $\tilde{K}_0(\mathbf{Z}/2p)$ which is cyclic, generated by the equivalence class of $[\mathbf{Z}/2]$, it suffices to prove that $j^*[\mathbf{Z}/2] = 0$. Let $\pi = \langle x, y \mid x^2 = y^p = 1, xy = yx \rangle$, $R = \mathbf{Z}[\omega]$ with $\omega = \exp(2\pi i/p)$ and $R[\mathbf{Z}/2] \cong \mathbf{Z}\pi/(1 + y + \cdots + y^{p-1})$ the obvious isomorphism (mapping ω to y). Consider the pullback square of rings

$$\begin{array}{ccc} \mathbf{Z}\pi & \xrightarrow{\lambda} & R[\mathbf{Z}/2] \\ \downarrow & & \downarrow \\ \mathbf{Z}[\mathbf{Z}/2] & \longrightarrow & \mathbf{F}_p[\mathbf{Z}/2] \end{array}$$

with obvious maps. Since $u(R[\mathbf{Z}/2]) \rightarrow u(\mathbf{F}_p[\mathbf{Z}/2])$ is surjective (cf. Reiner–Ullom [12, §7]) we get from the associated Milnor–Mayer–Vietoris sequence a monomorphism

$$\lambda_*: K_0\mathbf{Z}\pi \rightarrow K_0R[\mathbf{Z}/2].$$

Let $P \subset \mathbf{Z}\pi$ be the ideal generated by $(1 - y)$ and 2 . Then $\mathbf{Z}\pi/P \cong \mathbf{F}_2[\mathbf{Z}/2]$ is certainly cohomologically trivial and hence P is projective (cf. Rim [13]). Since $\mathbf{F}_2[\mathbf{Z}/2]/N = \mathbf{Z}/2$ we see that $j^*[\mathbf{Z}/2] = [\mathbf{Z}\pi/N] - [\mathbf{Z}\pi/N \otimes_\pi P]$. It suffices therefore to show that $[P] = [\mathbf{Z}\pi] \in K_0\mathbf{Z}\pi$. But $\lambda_*[P] = [(1 - \omega, 2)] = [R[\mathbf{Z}/2]]$ since $R/(1 - \omega)R \cong \mathbf{Z}/p$ and p odd. Hence $[P] = [\mathbf{Z}\pi]$ because λ_* is injective, from where we conclude that $j^*[\mathbf{Z}/2] = 0$. In case $\pi = D_{2p}$ we proceed in a similar way. Notice that $K_0(\mathbf{Z}/2p)$ is freely generated by $[\mathbf{Z}/2]$ and $[\mathbf{Z}/p]$. From Corollary 3.5 we infer that $j^*[\mathbf{Z}/p] = 0$ and we are therefore left showing that $j^*[\mathbf{Z}/2] = 0$. Let $D_{2p} = \langle x, y \mid x^2 = y^p, yxy = x \rangle$. Notice that $P = (1 - y)\mathbf{Z}\pi + 2\mathbf{Z}\pi$ is a two-sided ideal with $\mathbf{Z}\pi/P \cong \mathbf{F}_2[\mathbf{Z}/2]$, which is cohomologically trivial. Hence P is a projective π -module and clearly $j^*[\mathbf{Z}/2] = [\mathbf{Z}\pi/N] - [\mathbf{Z}\pi/N \otimes_\pi P]$. In order to see that $[P] = [\mathbf{Z}\pi]$ we consider the square of rings

$$\begin{array}{ccc} \mathbf{Z}\pi & \xrightarrow{\lambda} & R_t[\mathbf{Z}/2] \\ \downarrow & & \downarrow \\ \mathbf{Z}[\mathbf{Z}/2] & \longrightarrow & \mathbf{F}_p[\mathbf{Z}/2] \end{array}$$

with $R_t[\mathbf{Z}/2] = \mathbf{Z}\pi/(1 + y + \cdots + y^{p-1})$ a twisted group ring. By [12, §7] $u(R_t[\mathbf{Z}/2]) \rightarrow u(\mathbf{F}_p[\mathbf{Z}/2])$ is surjective and hence

$$\lambda_*: K_0\mathbf{Z}\pi \rightarrow K_0R_t[\mathbf{Z}/2]$$

is injective. Since $\mathbf{F}_2[\mathbf{Z}/2] \otimes_{\pi} R_t[\mathbf{Z}/2] \cong \mathbf{F}_2[\mathbf{Z}/2] \otimes_{\pi} \mathbf{F}_p[\mathbf{Z}/2] = 0$ we infer that $\lambda_*[P] = [R_t[\mathbf{Z}/2]]$ and whence $[P] = [\mathbf{Z}\pi]$ from the injectivity of λ_* . Therefore $j^*[\mathbf{Z}/2] = 0$, which completes the proof.

LEMMA 2.8. *Let $\pi = \mathbf{Z}/15$. Then $T(3)$ is the element of order 2 in $K_0\mathbf{Z}[\mathbf{Z}/15]$.*

Proof. Let $\pi = \langle x, y \mid x^3 = y^5 = 1, xy = yx \rangle$ and let $P = (x+2)\mathbf{Z}\pi + (y-1)\mathbf{Z}\pi$. Then $\mathbf{Z}\pi/P = M$ is cyclic of order 9 with y operating trivially and x operating by multiplication with 7 mod 9. One checks easily that M is cohomologically trivial using the criterion of [13]. Hence P is projective. Since $M/NM \cong \mathbf{Z}/3$ as trivial $\mathbf{Z}\pi/N$ -module we infer that

$$j^*[\mathbf{Z}/3] = [\mathbf{Z}\pi/N] - [\mathbf{Z}\pi/N \otimes_{\pi} P] \in K_0(\mathbf{Z}\pi/N).$$

Notice that $\text{im } S = 0$ since π is cyclic. Hence we can think of $T(3)$ to be the element $[\mathbf{Z}\pi] - [P] \in K_0(\mathbf{Z}\pi)$. Recall that $K_0\mathbf{Z}\pi \cong \mathbf{Z} \oplus \mathbf{Z}/2$ by Kervaire–Murthy [7]. Since P is projective of rank 1, it remains therefore to prove that $[P] \neq 0$ in $\tilde{K}_0(\mathbf{Z}\pi) \cong \mathbf{Z}/2$. For this we consider the pullback square (with obvious maps)

$$\begin{array}{ccc} \mathbf{Z}\pi & \longrightarrow & R[\mathbf{Z}/3] \\ \downarrow & & \downarrow \\ \mathbf{Z}[\mathbf{Z}/3] & \longrightarrow & \mathbf{F}_5[\mathbf{Z}/3] \end{array}$$

where $R = \mathbf{Z}[\exp(2\pi i/5)]$. The associated Milnor–Mayer–Vietoris sequence yields a map

$$\partial: u(\mathbf{F}_5[\mathbf{Z}/3]) \rightarrow \tilde{K}_0\mathbf{Z}[\mathbf{Z}/15]$$

By [7] ∂ factors through $u(\mathbf{F}_5(\omega)) \cong \mathbf{Z}/24$ where $\mathbf{F}_5(\omega)$ is the field $\mathbf{F}_5[\mathbf{Z}/3]/(1+x+x^2)$, ω is the residue class of the generator $x \in \mathbf{Z}/3$. Furthermore, ∂ is surjective (cf. [7]). Notice that $R[\mathbf{Z}/3] \otimes_{\pi} M = 0$ and therefore $P' = R[\mathbf{Z}/3] \otimes_{\pi} P \cong R[\mathbf{Z}/3]$. Furthermore $P'' = \mathbf{Z}[\mathbf{Z}/3] \otimes_{\pi} P \subset \mathbf{Z}[\mathbf{Z}/3]$ is the principal ideal generated by $(x+2)$. Hence (cf. [9])

$$P = \{((x+2)a, b) \in \mathbf{Z}[\mathbf{Z}/3] \times R[\mathbf{Z}/3] \mid \overline{(x+2)}\bar{a} = \bar{b} \in \mathbf{F}_5[\mathbf{Z}/3]\}.$$

Notice that $\overline{x+2} \in u(\mathbf{F}_5[\mathbf{Z}/3])$ corresponds to $(3, \omega+2) \in u(\mathbf{F}_5) \times u(\mathbf{F}_5(\omega))$ under the obvious isomorphism $\mathbf{F}_5[\mathbf{Z}/3] \cong \mathbf{F}_5 \times \mathbf{F}_5(\omega)$. It follows therefore that $\partial(\overline{x+2}) = \partial(\omega+2) = [P]$ and, since $(\omega+2)$ has order 24 in $u(\mathbf{F}_5(\omega))$, we conclude that $[P]$ must have order 2. This completes the proof of the lemma.

We can now complete the proof of Theorem 2.5: First, if n is a prime power, π (having cyclic Sylow subgroups) is necessarily cyclic and therefore $j^*: K_0(\mathbf{Z}/n) \rightarrow K_0(\mathbf{Z}\pi/N)$ factors through $\tilde{K}_0(\mathbf{Z}/n)$ by Lemma 2.6. But $\tilde{K}_0(\mathbf{Z}/n) = 0$ for n a prime power. Hence $T \equiv 0$ in this case. If $n = 2p$, p an odd prime, then $\pi = \mathbf{Z}/2p$ or D_{2p} and it follows from Lemma 2.7 that $T \equiv 0$. Thus (i) holds. Assume now that π is cyclic of order $\prod p_i^{k_i} = n$. Then $T(\prod p_i) = t\varphi(\prod p_i) = t[\mathbf{Z}/n] = 0$, since $j^*[\mathbf{Z}/n] = 0$ by Lemma 2.6. Therefore (ii) holds. For (iii) notice that in case $n = 15$, π is necessarily cyclic. Hence, by applying Lemma 2.6, $j^*[\mathbf{Z}/3] = -j^*[\mathbf{Z}/5]$ or $T(3) = -T(5)$. Moreover $T(3)$ has order 2 by Lemma 2.8 and therefore $T(3) = T(5) \neq 0$. Finally $\text{im } S = 0$ for π cyclic or $\pi = D_{2p}$, p an odd prime (cf. Ullom [16]) and hence $\text{im } S = 0$ for the groups considered in (i)–(iii) of Theorem 2.5.

3. Applications to homologically nilpotent spaces

Following [3] we call a connected space X *homologically nilpotent*, if $\pi_1 X$ operates nilpotently on $H_i \tilde{X}$ for all i . In particular, if X is a nilpotent space, then X is homologically nilpotent and conversely, a homologically nilpotent space X is nilpotent if and only if $\pi_1 X$ is a nilpotent group (cf. [10]). Let $H_*^\pi(X; I\mathbf{Z}\pi)$ denote the homology of the (left) $\mathbf{Z}\pi/N$ -complex $I\mathbf{Z}\pi \otimes_\pi C_* \tilde{X}$, where $\pi = \pi_1 X$; similarly for $H_*^\pi(X; \mathbf{Z}\pi/N)$.

LEMMA 3.1. *If X is homologically nilpotent with $\pi_1 X$ of finite order n , then $H_i^\pi(X; I\mathbf{Z}\pi)$ and $H_i^\pi(X; \mathbf{Z}\pi/N)$ are nilpotent $\mathbf{Z}\pi/N$ -modules for all i . If, in addition, n is square free and X finitely dominated, then $H_i^\pi(X; I\mathbf{Z}\pi)$ and $H_i^\pi(X; \mathbf{Z}\pi/N)$ are of type FP over $\mathbf{Z}\pi/N$.*

Proof. Consider the long exact homology sequence associated with the exact sequence of chain complexes $0 \rightarrow I\mathbf{Z}\pi \otimes_\pi C_* \tilde{X} \rightarrow C_* \tilde{X} \rightarrow C_* X \rightarrow 0$. Standard results on nilpotent actions (cf. [6, Chapter I.4]) imply then that $H_i^\pi(X; I\mathbf{Z}\pi)$ is a nilpotent π -module. Since the π -action on $H_i^\pi(X; I\mathbf{Z}\pi)$ factors through $\mathbf{Z}\pi/N$, we conclude that $H_i^\pi(X; I\mathbf{Z}\pi)$ is a nilpotent $\mathbf{Z}\pi/N$ -module. If X is dominated by a finite complex and $\pi_1 X$ finite, then certainly $H_i^\pi(X; I\mathbf{Z}\pi)$ is a finitely generated $\mathbf{Z}\pi/N$ -module. Hence, by Lemma 1.4, $H_i^\pi(X; I\mathbf{Z}\pi)$ is of type FP. The proof for $H_i^\pi(X; \mathbf{Z}\pi/N)$ is similar, using $0 \rightarrow C_* X \rightarrow C_* \tilde{X} \rightarrow \mathbf{Z}\pi/N \otimes_\pi C_* \tilde{X} \rightarrow 0$.

LEMMA 3.2. *Let X be a finitely dominated homologically nilpotent space with finite fundamental group of order n and let $H_*(X, \tilde{X})$ denote the homology of the mapping cylinder of $\tilde{X} \rightarrow X \bmod \tilde{X}$. Then the groups $H_i^\pi(X, I\mathbf{Z}\pi)$ and $H_{i+1}(X, \tilde{X})$ have the same finite cardinality $c(i)$ for $i \geq 0$, and $c(i)$ is a unit in $\mathbf{Z}[1/n]$.*

Proof. Since $H_i^\pi(X; I\mathbb{Z}\pi)$ is a finitely generated nilpotent $\mathbb{Z}\pi/N$ -module, it has by Lemma 1.4 a cardinality $c(i)$ which is a unit in $\mathbb{Z}[1/n]$. From long exact homology sequences it is obvious that $c(i)$ is also given by

$$c(i) = (\text{card coker } (H_{i+1}\tilde{X} \rightarrow H_{i+1}X)) \cdot (\text{card ker } (H_i\tilde{X} \rightarrow H_iX))$$

which equals $\text{card } H_{i+1}(X, \tilde{X})$.

It follows that for X as in Lemma 3.2, the rational number

$$\rho(X) = \text{card } H_{\text{odd}}(X, \tilde{X}) / \text{card } H_{\text{ev}}(X, \tilde{X})$$

is a well defined unit in $\mathbb{Z}[1/n]$. This unit $\rho(X)$ is related to the finiteness obstruction $wX \in K_0\mathbb{Z}\pi_1X$ of Wall (cf. [17], [10]) in the following way.

THEOREM 3.3. *Let X be a homologically nilpotent space with non-trivial fundamental group π of square free order n . Suppose further that X is dominated by a finite complex and let $\bar{w}X$ denote the image of wX in $(K_0\mathbb{Z}\pi_1X)/\text{im } S$. Then*

$$T\rho(X) = \bar{w}X$$

If X is in addition nilpotent (i.e. π_1X is cyclic) then

- (i) $T\rho(X) = wX$, and
- (ii) $wX = 0$ in case $\rho(X)$ is a power (positive or negative) of n .

Before proving Theorem 3.3 we will establish a different way of computing $\rho(X)$.

LEMMA 3.4. *Let X be as in (3.2) and let $H_*^\pi(X; \mathbb{Z}\pi/N)$ denote the homology of $\mathbb{Z}\pi/N \otimes_\pi C_*(\tilde{X})$. Then*

$$\rho(X) = \text{card } H_{\text{ev}}^\pi(X; \mathbb{Z}\pi/N) / \text{card } H_{\text{odd}}^\pi(X; \mathbb{Z}\pi/N)$$

Proof. We may of course assume that $n = \text{card } \pi_1X > 1$. Since $H_*(\tilde{X}; \mathbb{Q})$ is semisimple and nilpotent as π -module, we have $H_*(\tilde{X}; \mathbb{Q}) \cong H_*(X; \mathbb{Q})$ and therefore the Euler characteristic of X vanishes. Thus, for all primes p , $\chi_p(X) = \sum (-1)^i \dim H_i(X; \mathbb{Z}/p) = 0$. Since n is square free, we infer then that

$$\text{card } H_{\text{ev}}(X; \mathbb{Z}/n) = \text{card } H_{\text{odd}}(X; \mathbb{Z}/n).$$

The long exact homology sequence associated with the exact sequence $I\mathbf{Z}\pi \rightarrow \mathbf{Z}\pi/N \rightarrow \mathbf{Z}/n$ then yields

$$\begin{aligned} \text{card } H_{\text{ev}}^{\pi}(X; \mathbf{Z}\pi/N) / \text{card } H_{\text{odd}}^{\pi}(X; \mathbf{Z}\pi/N) \\ = \text{card } H_{\text{ev}}^{\pi}(X; I\mathbf{Z}\pi) / \text{card } H_{\text{odd}}^{\pi}(X; I\mathbf{Z}\pi) \end{aligned}$$

from where the result follows, using Lemma 3.2.

Proof of Theorem 3.3. Let $pr_*: K_0(\mathbf{Z}\pi) \rightarrow K_0(\mathbf{Z}\pi/N)$ be the map induced by the projection. If \bar{C}_* denotes a chain complex of type *FP*, homotopy equivalent to the singular complex of \tilde{X} , then $pr_*wX = \sum (-1)^i [\mathbf{Z}\pi/N \otimes_{\pi} \bar{C}_i]$. Since $H_i^{\pi}(X; \mathbf{Z}\pi/N)$ is of type *FP* (cf. Lemma 3.1) we infer that (cf. [11])

$$\sum (-1)^i [\mathbf{Z}\pi/N \otimes_{\pi} \bar{C}_i] = \sum (-1)^i [H_i^{\pi}(X; \mathbf{Z}\pi/N)]$$

and therefore, since $[H_i^{\pi}(X; \mathbf{Z}\pi/N)] = T(\text{card } H_i^{\pi}(X; \mathbf{Z}\pi/N))$ by Lemma 2.4, we get

$$pr_*wX = \bar{w}X = T(\text{card } H_{\text{ev}}^{\pi}(X; \mathbf{Z}\pi/N) / \text{card } H_{\text{odd}}^{\pi}(X; \mathbf{Z}\pi/N)) = T\rho(X).$$

In case X is in addition nilpotent, $\pi_1 X$ is necessarily cyclic and therefore $\text{im } S = 0$. Hence $T\rho(X) = wX$ in this case. Furthermore, if $\rho(X) = n^k$, we conclude from Theorem 2.5(ii) that $wX = T(n^k) = 0$. This completes the proof of Theorem 3.3.

As a first application we will prove the following algebraic result, which is used to prove Lemma 2.7.

COROLLARY 3.5. *Let π denote the dihedral group D_{2p} , p an odd prime. Then $j^*[\mathbf{Z}/p] = 0 \in K_0(\mathbf{Z}D_{2p}/N)$.*

Proof. By Theorem A [14] there is a finite simplicial complex \tilde{X} of the homotopy type of S^3 on which D_{2p} acts freely and simplicially. Let $X = \tilde{X}/D_{2p}$. Then

$$H_i(X, \tilde{X}) = \begin{cases} 0, & \text{if } i = 0, 2 \\ \mathbf{Z}/2, & \text{if } i = 1 \\ \mathbf{Z}/2p, & \text{if } i = 3 \end{cases}$$

Thus $\rho(X) = 4p$. Since X is a finite complex with trivial action of $\pi_1 X$ on $H_*\tilde{X}$, we infer that $wX = 0$ and therefore $T(4p) = 0$. We have already observed in course of the proof of Lemma 2.7 that $j^*[\mathbf{Z}/2] = 0$. Hence

$$j^*[\mathbf{Z}/p] = j^*[(\mathbf{Z}/2) \oplus (\mathbf{Z}/2) \oplus (\mathbf{Z}/p)] = T(4p) = 0.$$

Recall that two nilpotent spaces X and Y of finite type are of the same *genus* (cf. [6]) if their p -localizations X_p, Y_p are homotopy equivalent for all primes p . In [10] one finds an example of two finitely dominated nilpotent spaces X and Y of the same genus with fundamental groups of order 8 such that $wX \neq 0$ but $wY = 0$. For fundamental groups of square free orders, such an example is impossible. Namely one has

COROLLARY 3.6. *Let X and Y be two finitely dominated nilpotent spaces of the same genus, with non trivial fundamental groups of square free order n . Then $\rho(X) = \rho(Y)$ and wX, wY have the same finite orders.*

Proof. Notice that $\pi_1 X$ and $\pi_1 Y$ are abelian and whence $\pi_1 X \cong \pi_1 Y$, since X and Y are of the same genus (cf. [6]). Furthermore, there are for all primes p commutative diagrams

$$\begin{array}{ccc} \tilde{X}_p & \longrightarrow & \tilde{Y}_p \\ pr \downarrow & & \downarrow pr \\ X_p & \longrightarrow & Y_p \end{array}$$

with the horizontal maps being homotopy equivalences. Thus

$$H_i(X, \tilde{X}; \mathbf{Z}_{(p)}) \cong H_i(X_p, \tilde{X}_p; \mathbf{Z}_{(p)}) \cong H_i(Y_p, \tilde{Y}_p; \mathbf{Z}_{(p)}) \cong H_i(Y, \tilde{Y}; \mathbf{Z}_{(p)})$$

and therefore $H_i(X, \tilde{X}) \cong H_i(Y, \tilde{Y})$ since the groups $H_i(X, \tilde{X})$ and $H_i(Y, \tilde{Y})$ are finite. Hence $\rho X = \rho Y$ and, since $\pi_1 X \cong \pi_1 Y$, it follows from Theorem 3.3(i) that wX and wY have the same orders; the orders must be finite, because X and Y must have vanishing Euler characteristic (cf. proof of Lemma 3.4).

Another application of Theorem 3.3 is the following vanishing theorem for the Wall obstruction.

COROLLARY 3.7. *Let X be a finitely dominated homologically nilpotent space with fundamental group of order p or $2p$, p a prime. Then $wX = 0$ and X is therefore of the homotopy type of a finite complex.*

Proof. First consider the case $p = 2$. By a result of Fröhlich (cf. [5, Theorem 6(i)]) we infer that $\tilde{K}_0 \pi_1 X = 0$ and, since the Euler characteristic of X is necessarily 0 (cf. proof of Lemma 3.4), it follows that $wX = 0$. Second, let p denote an odd prime. Then we may apply Theorem 3.3 and obtain $T\rho(X) = \bar{w}X$. Since $T \equiv 0$ and $\text{im } S = 0$ for the $\pi_1 X$ in question (cf. Theorem 2.5), we infer that $wX = 0$.

It is sometimes possible to compute $\rho(X)$ directly from H_*X , giving rise to a particular simple formula for wX . We will treat one such case in the next section and plan to treat other cases in a forthcoming paper.

4. The Wall obstruction for H -spaces

We want to prove the Theorem II mentioned in the introduction.

Suppose X is a space with $\oplus H_i(X; \mathbf{Z})$ finitely generated and let p denote a fixed prime. Then we write

$$\chi_p(X, t) = \sum (-1)^i \beta_i t^i, \quad \beta_i = \dim H_i(X; \mathbf{Z}/p)$$

for the *Poincaré polynomial* of X with respect to \mathbf{Z}/p . Define

$$e_p(X) = \left. \frac{d}{dt} \chi_p(X, t) \right|_{t=-1}$$

and, in case $\pi_1 X$ has finite order n , define

$$e(X) = \prod_{p|n} p^{e_p(X)}$$

THEOREM 4.1. *Let X be a finitely dominated nilpotent space with fundamental group π of square free order. Suppose that $H_i^\pi(X; I\mathbf{Z}\pi)$ is a trivial π -module for all i . Then*

$$\rho(X) = e(X)$$

Proof. We may assume that $\pi \neq \{1\}$. Let x denote a generator of the necessarily cyclic group π . Then there is a exact sequence of $\mathbf{Z}\pi/N$ -modules

$$0 \rightarrow I\mathbf{Z}\pi \xrightarrow{1-x} I\mathbf{Z}\pi \rightarrow \mathbf{Z}/n \rightarrow 0$$

where n denotes the order of π . Since $(1-x)$ induces 0 in $H_*^\pi(X; I\mathbf{Z}\pi)$, the associated long exact homology sequence breaks up into short exact sequences

$$0 \rightarrow H_i^\pi(X; I\mathbf{Z}\pi) \rightarrow H_i(X; \mathbf{Z}/n) \rightarrow H_{i-1}^\pi(X; I\mathbf{Z}\pi) \rightarrow 0$$

for $i \geq 0$. Since $H_i^\pi(X; I\mathbf{Z}\pi)$ is a trivial $\mathbf{Z}\pi/N$ -module and n is square free, $H_i^\pi(X; I\mathbf{Z}\pi) \otimes \mathbf{Z}_{(p)}$ is a \mathbf{Z}/p -vector space. Define

$$\beta_i = \dim H_i(X; \mathbf{Z}/p), \quad \gamma_i = \dim H_i^\pi(X; I\mathbf{Z}\pi) \otimes \mathbf{Z}_{(p)}.$$

Then

$$\gamma_i = \beta_i - \gamma_{i-1} = \beta_i - \beta_{i-1} + \cdots + (-1)^i \beta_0$$

and

$$\begin{aligned} \sum (-1)^j \gamma_j &= \sum (\beta_0 - \beta_1 + \cdots + (-1)^j \beta_j) \\ &= (m+1)\beta_0 - m\beta_1 + \cdots + (-1)^m \beta_m \end{aligned}$$

where m denotes the largest integer k with $\beta_k \neq 0$. Hence

$$\begin{aligned} \sum (-1)^j \gamma_j &= (-1)^m \frac{d}{dt} \left(t^{m+1} \chi_p \left(X, \frac{1}{t} \right) \right) \Big|_{t=-1} \\ &= (m+1) \chi_p(X, -1) + \chi'_p(X, -1). \end{aligned}$$

Since X is homologically nilpotent with non-trivial finite fundamental group, the Euler characteristic of X is 0 (cf. proof of Lemma 3.4) and hence $\chi_p(X, -1) = 0$. The above equation reduces therefore to

$$\sum (-1)^j \gamma_j = \chi'_p(X, -1) = e_p(X).$$

Hence $\text{card } H_{\text{ev}}^\pi(X; I\mathbf{Z}\pi) / \text{card } H_{\text{odd}}^\pi(X; I\mathbf{Z}\pi) = \prod_{p|n} p^{e_p(X)} = e(X)$ and therefore $\rho(X) = e(X)$.

In order to prove that for an H -space X the π -operation on $H_*(X; I\mathbf{Z}\pi)$ is trivial, we will need the following lemma.

LEMMA 4.2. *Let X be an H -space and $a \in \pi_1 X$. Then the induced covering transformation $a_*: \tilde{X} \rightarrow \tilde{X}$ is equivariantly homotopic to the identity.*

Proof. Without loss of generality we may assume that the H -structure $\mu: X \times X \rightarrow X$ has the base point as a strict identity. Equip \tilde{X} with the canonical H -structure $\tilde{\mu}$. Then $pr^{-1}(*)$ is a central subgroup of $(\tilde{X}, \tilde{\mu})$, naturally isomorphic to $\pi_1 X$. We may thus think of a as an element of $pr^{-1}(*)$ which acts on \tilde{X} by left multiplication. Choosing a path from $a \in pr^{-1}(*)$ to $*$ we get a homotopy of the

map a_* to Id , which is equivariant with respect to the $\pi_1 X = pr^{-1}(*)$ -action, because $pr^{-1}(*)$ is central.

COROLLARY 4.3. *If X is an H -space, then $H_*^\pi(X; I\mathbb{Z}\pi)$ is a trivial $\mathbb{Z}\pi/N$ -module.*

This is clear since by 4.2, the operation of $a \in \pi_1 X$ on $C_*\tilde{X}$ is chain homotopic to Id as map of $\pi_1 X$ -complexes, and therefore the induced action of a on $I\mathbb{Z}\pi \otimes_\pi C_*\tilde{X}$ is chain homotopic to Id .

Proof of Theorem II of the Introduction. If X is of rank 1, then X is equivalent to one of the spaces S^1 , S^3 , S^7 , $\mathbb{R}P^3$ or $\mathbb{R}P^7$ (cf. Browder [2]). Hence $wX = 0$ in these cases. If $\text{rank}(X) \geq 2$, then $\chi_p(X, t)$ contains a factor $(1 - t^{n_1})(1 - t^{n_2})$ with n_1 and n_2 odd. Therefore

$$e_p(X) = \chi'_p(X, -1) = 0$$

for all primes p . In particular we obtain $e(X) = 1$ and, since $H_i^\pi(X; I\mathbb{Z}\pi)$ is a trivial π -module for all i we infer from Theorem 4.1 that $\rho(X) = e(X)$. Hence $wX = T\rho(X) = 0$.

5. Appendix

If X denotes a homologically nilpotent space with finite fundamental group, then there is a simple criterion for deciding whether X is dominated by a finite complex.

THEOREM 5.1. *Let X be a homologically nilpotent complex with finite fundamental group. Then the following are equivalent.*

- (i) X is dominated by a finite complex
- (ii) $H_i(\tilde{X}; \mathbb{Z})$ and $H_i(X; \mathbb{Z})$ are finitely generated abelian groups for all i and zero for i sufficiently large.

Proof. Certainly (i) implies (ii). If (ii) is given, then from [3, Corollary 3.4] we infer that X has the homotopy type of a finite dimensional complex. Since $\mathbb{Z}\pi_1 X$ is noetherian and $H_i(\tilde{X}; \mathbb{Z})$ finitely generated for all i , Theorems B and F of [17] imply that X is dominated by a finite complex.

A more general result of this type in case X is nilpotent was proved in [11].

REFERENCES

- [1] BASS, H., *Algebraic K-Theory*, Benjamin 1968.
- [2] BROWDER, W., *Higher torsion in H-spaces*, Trans. Amer. Math. Soc. 108 (1963), 353–375.
- [3] BROWN, K. S. and KAHN, P. J., *Homotopy dimension and simple cohomological dimension of spaces*, preprint.
- [4] CARTAN, H. and EILENBERG, S., *Homological Algebra*, Princeton University Press 1956.
- [5] FRÖHLICH, A., *On the classgroup of integral groupings of finite abelian groups*, Mathematika 16 (1969), 143–152.
- [6] HILTON, P., MISLIN, G., and ROITBERG, J., *Localization of nilpotent groups and spaces*, Mathematics Studies 15, North-Holland, Amsterdam 1975.
- [7] KERVAIRE, M. A. and MURTHY, M. P., *On the projective class group of cyclic groups of prime power order p^n* , preprint.
- [8] LUNDMARK, R., Thesis, E.T.H.-Zürich 1976.
- [9] MILNOR, J., *Introduction to Algebraic K-Theory*, Princeton University Press, 1971.
- [10] MISLIN, G., *Wall's obstruction for nilpotent spaces*, Topology 14 (1975), 311–318.
- [11] —, *Finitely dominated nilpotent spaces*, Ann. of Math. 103 (1976), 547–556.
- [12] REINER, I. and ULLOM, S., *A Mayer–Vietoris sequence for class groups*, J. of Algebra 31 (1974), 305–342.
- [13] RIM, D. S., *Modules over finite groups*, Ann. of Math. 69 (3) 1959, 700–712.
- [14] SWAN, R. G., *Periodic resolutions for finite groups*, Ann. of Math. 72 (2) 1960, 267–291.
- [15] —, *The p -period of a finite group*, Illinois J. of Math. 4 (1960), 341–346.
- [16] ULLOM, S. V., *Nontrivial lower bounds for class groups of integral group rings*, to appear in Illinois J. of Math.
- [17] WALL, C. T. C., *Finiteness conditions for CW-complexes*, Ann. of Math. 81 (1965), 56–69.

E.T.H. Rämistr. 101,
8006 Zurich
Switzerland

Received October 11, 1976.

