

Hölder continuity of conformal mappings and non-quasiconformal Jordan curves.

Autor(en): **Becker, Jochen / Pommerenke, Christian**

Objekttyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **57 (1982)**

PDF erstellt am: **28.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-43884>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Hölder continuity of conformal mappings and non-quasiconformal Jordan curves

JOCHEN BECKER and CHRISTIAN POMMERENKE

Let $\Gamma \subset \mathbf{C}$ be a quasiconformal curve (quasicircle), that is the image of the unit circle under a quasiconformal mapping of the plane. Let f and f^* denote conformal mappings of the unit disk \mathbf{D} onto $\text{int } \Gamma$ and $\text{ext } \Gamma$ respectively.

It is well-known that f and f^* have quasiconformal extensions onto the plane [3], p. 98. Because of the Hölder continuity of quasiconformal mappings ([3], p. 71), f and f^* as well as their inverse mappings f^{-1} and f^{*-1} satisfy Hölder conditions:

$$\begin{aligned} |f(z_1) - f(z_2)| &\leq K |z_2 - z_1|^\alpha & (z_1, z_2 \in \mathbf{D}), \\ |f^{-1}(w_1) - f^{-1}(w_2)| &\leq L |w_1 - w_2|^\beta & (w_1, w_2 \in \text{int } \Gamma) \end{aligned} \quad (1)$$

and similarly for f^* where one has to use the spherical metric in $\text{ext } \Gamma$. The Hölder exponents α ($0 < \alpha \leq 1$) and β ($1 \leq \beta < 2$) depend only on Γ [4], [5], [8], p. 287, 289, 347.

By means of a simple geometrical characterization of (1) (Theorem 1), we construct a non-quasiconformal Jordan curve (Theorem 2) such that nevertheless f , f^* , and also f^{-1} , f^{*-1} remain Hölder continuous. This shows that Hölder continuity of all these conformal mappings is only a necessary, but not a sufficient condition for quasicircles.

Let $G \subsetneqq \mathbf{C}$ be a simply connected domain and let $h(w_1, w_2)$ denote the hyperbolic distance of the points $w_1, w_2 \in G$ defined by

$$h(w_1, w_2) = \log \frac{|1 - \bar{z}_1 z_2| + |z_2 - z_1|}{|1 - \bar{z}_1 z_2| - |z_2 - z_1|} \quad (w_i = f(z_i), i = 1, 2) \quad (2)$$

where f is a conformal mapping of \mathbf{D} onto G . Let $\delta(w) = \text{dist}(w, \partial G)$ denote the (Euclidean) boundary distance.

THEOREM 1. *If G is a bounded simply connected domain, $w_0 \in G$, and f a conformal mapping of \mathbf{D} onto G , then (1) is satisfied if and only if*

$$\limsup_{\delta(w) \rightarrow 0} [h(w_0, w) + \frac{1}{\alpha} \log \delta(w)] < +\infty. \quad (3)$$

Proof. It is well-known that (1) is equivalent to

$$|f'(z)| \leq M(1 - |z|^2)^{\alpha-1} \quad (|z| < 1) \quad (4)$$

(see for instance [2], p. 361–363). Introducing the non-Euclidean length element

$$\rho(w) |dw| = \frac{2 |dz|}{1 - |z|^2} \quad (w = f(z))$$

corresponding to (2) we see that (4) means

$$h(f(0), f(z)) \leq \frac{1}{\alpha} \log \left[\frac{M}{2} (1 + |z|)^{2\alpha} \rho(f(z)) \right].$$

Since G is bounded we have $\delta(f(z)) \rightarrow 0$ if and only if $|z| \rightarrow 1$. Thus we obtain as an equivalent condition

$$\limsup_{\delta(w) \rightarrow 0} \left[h(f(0), w) - \frac{1}{\alpha} \log \rho(w) \right] < +\infty. \quad (5)$$

Because of

$$\frac{1}{2} \leq \rho(w) \delta(w) \leq 2$$

[8], p. 22 it follows by the triangle inequality for $h(w_1, w_2)$ that (5) is the same as (3) which proves Theorem 1.

The following corollary gives a convenient sufficient condition for Hölder continuity.

COROLLARY. *Let f be a bounded univalent function defined in \mathbf{D} , $G = f(\mathbf{D})$, then f satisfies (1) if there are positive numbers M , δ_0 , δ_1 with the property that, for every $w \in G$ with $\delta(w) < \delta_0$, there exists $w_1 \in G$ with $\delta(w_1) \geq \delta_1$ and a connecting arc $C \subset G$, such that*

$$\int_C \frac{|d\omega|}{\delta(\omega)} \leq M + \frac{1}{2\alpha} \log \frac{1}{\delta(w)}. \quad (6)$$

Proof. We choose $w_0 = f(0)$. Then we have

$$\begin{aligned} h(w_0, w) &\leq h(w_0, w_1) + \int_C \rho(\omega) |d\omega| \\ &\leq \log \frac{1+|f^{-1}(w_1)|}{1-|f^{-1}(w_1)|} + 2 \int_C \frac{|d\omega|}{\delta(\omega)}. \end{aligned}$$

From (6) and $\delta(w_1) \geq \delta_1 > 0$ it follows that (3) is satisfied which implies (1) by Theorem 1.

We are now ready to construct a Jordan curve with the desired properties.

THEOREM 2. *There is a non-quasiconformal Jordan curve Γ such that the conformal mappings f and f^* of \mathbf{D} onto $\text{int } \Gamma$ and $\text{ext } \Gamma$ respectively, and also their inverse mappings are Hölder continuous.*

Proof. We set $Q = \{w = u + iv : |u| < 1, |v| < 1\}$ and $R_n = \{w = u + iv : 0 \leq u - 1 < a_n, |v - v_n| < \varepsilon_n\}$ where we choose $v_n = 1/n$, $\varepsilon_n = 2^{-n}$, $a_n = -\varepsilon_n \log \varepsilon_n$, $n = 2, 3, \dots$. We consider the domain $G = Q \cup \bigcup_{n=2}^{\infty} R_n$. Since $\Gamma = \partial G$ is a locally connected continuum without cut points it is a Jordan curve [8], p. 281. But it is not a quasicircle, because the Ahlfors criterion [1] is not satisfied. For we have

$$\frac{\text{diam } R_n}{2\varepsilon_n} \geq \frac{a_n}{2\varepsilon_n} \rightarrow +\infty (n \rightarrow \infty).$$

On the other hand, we shall show below that, for every $w \in G$, $\delta(w) < \frac{1}{2}$, there exists $w_1 \in G$, $\delta(w_1) \geq \frac{1}{2}$, and a connecting arc C such that

$$\int_C \frac{|d\omega|}{\delta(\omega)} \leq 2 \log \frac{1}{\delta(w)}. \quad (7)$$

Thus (6) is satisfied with $\alpha = \frac{1}{4}$ which shows that f is α -Hölder-continuous for some $\alpha \geq \frac{1}{4}$.

The Hölder continuity of f^{-1} , f^* , f^{*-1} follows easily from results by Näkki and Palka [6], [7]:

Let $d(w_1, w_2) = \inf_C \int_C |dw|$ denote the inner distance of $w_1, w_2 \in G$ where the infimum is taken over all connecting arcs $C \subset G$. Then, obviously, there exists a constant A such that

$$d(w_1, w_2) \leq A |w_2 - w_1| (w_1, w_2 \in G).$$

It follows from [6] that f^* and f^{-1} are Hölder continuous.

Considering the inner distance $d^*(w_1, w_2)$ with regard to $G^* = \text{ext } \Gamma$ one easily sees that, for every β ($0 < \beta < 1$), and $R > 0$, there exists a constant B such that

$$d^*(w_1, w_2) \leq B |w_2 - w_1|^\beta \quad (w_i \in G^*, |w_i| \leq R, i = 1, 2)$$

This implies, by another result of Näkki and Palka [6], [7], that f^{*-1} is also Hölder continuous.

Hence it remains to prove (7). We have to consider the following cases.

(a) $w = u + iv \in Q$, and, if $u \geq v \geq 0$, $|v - v_n| \geq \varepsilon_n$, $n = 2, 3, \dots$. Hence $\delta(w) = \min(1 - |u|, 1 - |v|)$. Choose $w_1 = 0$ and $C = [w_1, w_2]$ (segment with end points w_1, w_2). Then

$$\int_C \frac{|d\omega|}{\delta(\omega)} \leq \int_{\delta(w)}^1 \frac{\sqrt{2} dt}{t} = \sqrt{2} \log \frac{1}{\delta(w)}.$$

(b) $w = u + iv \in Q$, $u \geq v \geq 0$, $|v - v_n| < \varepsilon_n$ for some n (hence $\delta(w) > 1 - u$). Choose, for $\delta(w) < \frac{1}{2}$, $w_1 = \frac{1}{2} + iv$, $C = [w_1, w]$. Because of

$$\delta(w) = \sqrt{((1 - x^2) + (\varepsilon_n - |v - v_n|)^2)}, \quad \omega = x + iv \in C,$$

we obtain

$$\begin{aligned} \int_C \frac{|d\omega|}{\delta(\omega)} &= \log \frac{\frac{1}{2} + \sqrt{(\frac{1}{4} + (\varepsilon_n - |v - v_n|)^2)}}{1 - u + \sqrt{((1 - u)^2 + (\varepsilon_n - |v - v_n|)^2)}} \\ &\leq \log \frac{1 + \sqrt{(1 + 4\varepsilon_n^2)}}{2\delta(w)} < 2 \log \frac{1}{\delta(w)}. \end{aligned}$$

(c) $w = u + iv \in R_n$ for some n , $0 \leq u - 1 \leq a_n - \varepsilon_n$ hence $\delta(w) = \varepsilon_n - |v - v_n|$. Choose $w_1 = \frac{1}{2} + iv_n$ and $C = [w_1, u + iv_n] \cup [u + iv_n, w]$. Then

$$\begin{aligned} \int_C \frac{|d\omega|}{\delta(\omega)} &= \int_{1/2}^1 \frac{dx}{\sqrt{((1 - x)^2 + \varepsilon_n^2)}} + \int_1^u \frac{dx}{\varepsilon_n} + \int_0^{|v - v_n|} \frac{dy}{\varepsilon_n - y} \\ &< \frac{a_n}{\varepsilon_n} - \log(\varepsilon_n - |v - v_n|) \leq 2 \log \frac{1}{\delta(w)}. \end{aligned}$$

(d) $w = u + iv \in R_n$ for some n , $a_n - \varepsilon_n < u - 1$ hence $\delta(w) = \min(\varepsilon_n - |v - v_n|, 1 + a_n - u)$. Choose $w_1 = \frac{1}{2} + iv_n$, $C = [w_1, 1 + a_n - \varepsilon_n + iv_n] \cup [1 + a_n - \varepsilon_n + iv_n, w]$.

Then

$$\begin{aligned} \int_C \frac{|d\omega|}{\delta(\omega)} &\leq \int_{1/2}^1 \frac{dx}{\sqrt{(1-x)^2 + \varepsilon_n^2}} + \int_0^{a_n - \varepsilon_n} \frac{dx}{\varepsilon_n} + \int_{\delta(w)}^{\varepsilon_n} \frac{\sqrt{2} dt}{t} \\ &< -\log \varepsilon_n + \frac{a_n}{\varepsilon_n} + \sqrt{2} \log \frac{\varepsilon_n}{\delta(w)} \leq 2 \log \frac{1}{\delta(w)}. \end{aligned}$$

Thus we have proved that (7) holds in each case which completes the proof of Theorem 2.

Remark. The Hölder continuity of f^* could have also been shown by this method.

REFERENCES

- [1] L. V. AHLFORS, *Quasiconformal reflections*, Acta Math. 109 (1963), 291–301.
- [2] G. M. GOLUZIN, *Geometric Theory of Functions of a Complex Variable*, (Moscow 1952). German transl.: Deutscher Verlag der Wissenschaften, Berlin 1957.
- [3] O. LEHTO and K. I. VIRTANEN, *Quasiconformal Mappings in the Plane*, Springer, Berlin-Heidelberg-New York 1973.
- [4] F. D. LESLEY, *Hölder continuity of conformal mappings at the boundary via the strip method*, to appear.
- [5] R. NÄKKI and B. PALKA, *Quasiconformal circles and Lipschitz classes*, Comment. Math. Helvetici 55 (1980), 485–498.
- [6] R. NÄKKI and B. PALKA, *Lipschitz conditions, b-arcwise connectedness and conformal mappings*, to appear.
- [7] R. NÄKKI and B. PALKA, *Lipschitz classes and quasiconformal mappings*, Indiana J. Math., to appear.
- [8] CH. POMMERENKE, *Univalent Functions*, Vandenhoeck and Ruprecht, Göttingen 1975.

Technische Universität Berlin
 Fachbereich 3—Mathematik
 1000 Berlin 12
 Germany

Received January 21, 1982