

# On the cohomology of groups of p-length 1.

Autor(en): **Diethelm, Thomas**

Objekttyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **58 (1983)**

PDF erstellt am: **28.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-44605>

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# On the cohomology of groups of $p$ -length 1

THOMAS DIETHELM

## 1. Introduction

Let  $G$  be a finite group, whose order is divisible by the prime  $p$ , and let  $k$  denote the field of  $p$  elements. We consider the cohomology  $H^n(G, A)$ , where  $A$  is a simple  $kG$ -module. It is well known that  $H^n(G, A) \neq 0$  implies that  $A$  lies in the principal block of  $kG$ . We ask, if the converse is true, i.e. if to every simple  $kG$ -module  $A$  in the principal block there is an  $n \in \mathbb{N}$  with  $H^n(G, A) \neq 0$ .

Swan proved that this is true for the trivial module  $k$ . Therefore the above question has a positive answer for  $p$ -nilpotent groups ( $G = O_{p'p}G$ ). In this paper we show: (Theorem 5.3) if  $G = O_{p'pp'p}G$ , then there are infinitely many  $n \in \mathbb{N}$  with  $H^n(G, A) \neq 0$ .

In §3 we first consider the case where  $G$  is of  $p$ -length 1. In order to show the nontriviality of  $H^n(G, A)$  we analyze the action of the  $p'$ -group  $Q = G/O_{p'p}G$  on the cohomology ring  $H^*(O_{p'p}G/O_{p'}G, k)$  of the  $p$ -group  $P = O_{p'p}G/O_{p'}G$ . We prove the following result, which is of interest in its own right (Theorem 4.5):

If the  $p'$ -group  $Q$  acts faithfully on the  $p$ -group  $P$ , then every simple  $kQ$ -module  $A$  appears infinitely often in  $H^*(P, k)$  as a direct summand.

The proof of this result is by induction on the length of a central series of  $P$  with elementary abelian factors. With the aid of this result we can prove Theorem 4.6:

Let  $G$  be a group of  $p$ -length 1, and let  $A$  be a simple  $kG$ -module lying in the principal block of  $kG$ . Then  $H^n(G, A) \neq 0$  for infinitely many  $n \in \mathbb{N}$ .

In §5 we show how the result for groups  $G$  of  $p$ -length 1 can be used to treat the case where  $G = O_{p'pp'p}G$ . We do that by considering the extension

$$O_{p'pp'}G \rightarrowtail G \twoheadrightarrow G/O_{p'pp'}G.$$

Most of the results of this paper first appeared in the author's doctoral thesis (ETH, Zürich, Switzerland, 1981; adviser: U. Stammbach).

## 2. Techinal lemmas

As a preparation we state the following well known results:

**LEMMA 2.1.** *Let  $G$  be an extension of a  $p'$ -group  $N$  by a group  $H$ ,  $N \rightarrowtail G \twoheadrightarrow H$ . If  $V$  is an indecomposable  $kG$ -module lying in the principal block, then:*

$$H^n(G, V) \cong H^n(H, V); \quad n \geq 0.$$

*Proof.* Since  $N$  is a  $p'$ -group,  $V$  is centralised by  $N$  and the spectral sequence of the extension  $N \rightarrowtail G \twoheadrightarrow H$  collapses.

**LEMMA 2.2.** *Let  $G$  be an extension of a group  $N$  by a  $p'$ -group  $H$ . If  $V$  is a  $kG$ -module, then  $H^n(G, V) \cong H^n(N, V)^H$ ;  $n \geq 0$ .*

*Proof.* Since  $H$  is a  $p'$ -group, the spectral sequence of the extension  $N \rightarrowtail G \twoheadrightarrow H$  collapses.

**LEMMA 2.3.** *Let  $G$  be an extension of  $N$  by a group  $H$ , and let  $A$  be a  $kG$ -module with  $C_G(A) \supseteq N$ . Then:*

$$H^n(N, A)^H \cong \text{Hom}_{kH}(H_n(N, k), A); \quad n \geq 0.$$

*Proof.* Since  $A$  is a trivial  $kN$ -module, the universal coefficient theorem holds

$$H^n(N, A) \cong \text{Hom}_k(H_n(N, k), A).$$

The above isomorphism is natural and thus  $H$  acts diagonally on the right hand side. Hence

$$H^n(N, A)^H = \text{Hom}_{kH}(H_n(N, k), A).$$

## 3. The cohomology of groups of $p$ -length 1

Let  $G$  be a group of  $p$ -length 1 ( $G = O_{p'pp'}G$ ), and let  $A$  be a simple  $kG$ -module lying in the principal block of  $kG$ . Then  $O_{p'p}G \subseteq C_G(A)$ . ([6] p. 164.)

From Lemma 2.1 we obtain

$$H^i(G/O_{p'}G, A) \cong H^i(G, A)$$

and from Lemmas 2.1, 2.2

$$H^i(G, A) \cong H^i(O_{p'p}G/O_{p'}G, A)^{G/O_{p'p}G}.$$

Let  $Q$  denote the  $p'$ -group  $G/O_{p'p}G$ , and let  $P$  denote the  $p$ -group  $O_{p'p}G/O_{p'}G$ . Then Lemma 2.3 yields

$$H^n(G, A) \cong \text{Hom}_{kQ}(H_n(P, k), A).$$

This preparation allows the proof of the following result.

**THEOREM 3.1.** *Let  $G$  be a group of  $p$ -length 1, and let  $A$  be a simple  $kG$ -module lying in the principal block of  $kG$ . Then:*

$$H^n(G, A) \neq 0$$

*if and only if  $A$  is a direct summand of  $H_n(P, k)$ .*

*Proof.*

- “ $\Rightarrow$ ” If  $H^n(G, A)$  is nontrivial, then  $\text{Hom}_{kQ}(H_n(P, k), A)$  is nontrivial, and the simple  $kQ$ -module  $A$  is a direct summand of  $H_n(P, k)$ .
- “ $\Leftarrow$ ” By Maschke’s theorem  $H_n(P, k)$  is semi-simple. If  $A$  is a direct summand of  $H_n(P, k)$  the projection onto  $A$  is a nontrivial  $kQ$ -module homomorphism  $f: H_n(P, k) \rightarrow A$ . But the nontriviality of  $\text{Hom}_{kQ}(H_n(P, k), A)$  implies the nontriviality of  $H^n(G, A)$ .

**Note 3.1.** It follows from Theorem 3.1, that it is necessary to analyze the  $G/P$ -module structure of  $H_*(P, k)$  induced by conjugation of  $G$  in  $P$ . Since the cohomology  $H^*(P, k)$  is the dual of  $H_*(P, k)$ , this is equivalent to analyze the  $G/P$ -module structure of  $H^*(P, k)$ . The advantage of working in cohomology is, that we may use its algebra structure which is induced by the cup-product.

**Note 3.2.** Clearly the  $p'$ -group  $Q = G/O_{p'p}G$  acts faithfully on the  $p$ -group  $P = O_{p'p}G/O_{p'}G$ .

#### 4. The $kQ$ -module structure of $H^*(P, k)$

By Note 3.2, the  $p'$ -group  $Q$  acts faithfully on the  $p$ -group  $P$ . This action induces an action of  $Q$  on the cohomology ring  $H^*(P, k)$ .

Our problem is to determine these  $kQ$ -modules which are direct summands of  $H^*(P, k)$ .

**LEMMA 4.1.** *Let the  $p'$ -group  $Q$  act faithfully on the elementary abelian  $p$ -group  $E = C_p^{(1)} \times \cdots \times C_p^{(m)}$ . Then every simple  $kQ$ -module  $A$  is infinitely often a direct summand of  $H^*(E, k)$ .*

*Proof.* It is well known, that the cohomology ring  $H^*(E, k)$  contains the polynomial ring  $k[x_1, x_2, \dots, x_m]$ ;  $x_i \in H^2(C_p^{(i)}, k)$  as a subring. The generators  $x_1, x_2, \dots, x_m$  correspond to a basis of  $E$ , and  $Q$  acts faithfully on the subspace  $\langle x_1, x_2, \dots, x_m \rangle$  of  $H^2(E, k)$ . By the theorem of Steinberg [7], every simple  $kQ$ -module  $A$  is infinitely often a direct summand of  $k[x_1, x_2, \dots, x_m]$ , and  $k[x_1, x_2, \dots, x_m]$  is a direct summand of  $H^*(E, k)$ .

*Note 4.1.* The map  $\phi_s : k[x_1, \dots, x_m] \rightarrow k[x_1^{p^s}, \dots, x_m^{p^s}]$ ,  $f(x_1, \dots, x_m) \mapsto f(x_1, \dots, x_m)^{p^s} = f(x_1^{p^s}, \dots, x_m^{p^s})$ ;  $s = 0, 1, 2, \dots$  is a  $kQ$ -module isomorphism. Therefore  $k[x_1, x_2, \dots, x_m]$  contains infinitely many copies of itself.

**LEMMA 4.2.** *Let  $E = C_p^{(1)} \times C_p^{(2)} \times \cdots \times C_p^{(m)}$  be an elementary abelian central subgroup of the  $p$ -group  $P$ . Then for some  $s \in \mathbb{N}$  the polynomial ring  $k[x_1^{p^s}, x_2^{p^s}, \dots, x_m^{p^s}]$  lies in the image of the restriction map*

$$\text{res} : H^*(P, k) \rightarrow H^*(E, k).$$

*Proof.* We consider the spectral sequence  $E_2^{i,j} \cong H^i(P/E, H^j(E, k)) \Rightarrow H^{i+j}(P, k)$  of the extension  $E \rightarrowtail P \twoheadrightarrow P/E$ . Since  $E$  is a central subgroup, we get  $E_2^{0,j} = H^j(E, k)^{P/E} = H^j(E, k)$ .

There is a cup-product [4]

$$E_r^{i,j} \otimes E_r^{i',j'} \xrightarrow{\cup} E_r^{i+i',j+j'}$$

with the following rules

- (i)  $a \cdot b = (-1)^{ii'+jj'} b \cdot a$ ;
- (ii)  $d_r(a \cdot b) = d_r a \cdot b + (-1)^{i+j} a \cdot d_r b$   
 $a \in E_r^{i,j}; \quad b \in E_r^{i',j'}$ .

Suppose  $0 \neq x \in E_2^{0,2}$ . Since  $\text{char } k = p$ , one easily checks that  $d_2(x^p) = pd_2 x \cdot x^{p-1} = 0$ .

Now  $x^p$  is a nontrivial cocycle of  $E_3^{0,2p}$  and  $d_3(x^{p^2}) = p \cdot d_3 x^p \cdot x^{p(p-1)} = 0$ . Iteration of this process yields  $0 \neq x^{p^s} \in E_{s+2}^{0,2p^s}$ . By a theorem of Evens [1] the spectral sequence of a finite group extension stops, i.e. there is a  $t \in \mathbb{N}$  with  $E_t = E_\infty$ . Now  $s = t - 2$  yields  $0 \neq x^{p^s} \in E_\infty^{0,2p^s}$ , but  $x^{p^s}$  then lies in the image of the restriction map

$$\text{res} : H^{2p^s}(P, k) \rightarrow H^{2p^s}(E, k).$$

It follows that the polynomial ring  $k[x_1^{p^s}, \dots, x_m^{p^s}]$  lies in the image of the restriction map.

**Note 4.2.** It follows from the naturality of the *LHS*-spectral sequence, that, if the  $p'$ -group  $Q$  acts on the extension  $E \rightarrow P \twoheadrightarrow P/E$ , then the restriction map

$$\text{res}: H^*(P, k) \rightarrow H^*(E, k)$$

is a  $kQ$ -module homomorphism.

**LEMMA 4.3.** *Let the  $p'$ -group  $Q$  act on the central extension  $E \rightarrow P \twoheadrightarrow P/E$ . Let  $N$  denote the centraliser  $C_Q(E)$ . Then every simple  $k(Q/N)$ -module  $A$  is infinitely often a direct summand of  $H^*(P, k)$ .*

*Proof.* The group  $Q/N$  acts faithfully on  $E$ . By Lemma 4.1 and Note 4.1 every simple  $k(Q/N)$ -module  $A$  is infinitely often a direct summand of  $k[x_1^{p^s}, \dots, x_m^{p^s}]$ . By Lemma 4.2  $A$  is infinitely often a direct summand in the image of the restriction map

$$\text{res}: H^*(P, k) \rightarrow H^*(E, k),$$

and by Note 4.2  $A$  is infinitely often a direct summand of  $H^*(P, k)$ .

**THEOREM 4.4.** *Let the  $p'$ -group  $Q$  act on the central extension  $E \rightarrow P \twoheadrightarrow P/E$ .*

*If the simple  $kQ$ -module  $A$  is a direct summand of  $H^*(P/E, k)$ , then  $A$  is infinitely often a direct summand of  $H^*(P, k)$ .*

*Proof.* We consider the spectral sequence  $E_2^{i,j} \cong H^i(P/E, H^j(E, k)) \Rightarrow H^{i+j}(P, k)$  associated with the extension  $E \rightarrow P \twoheadrightarrow P/E$ . Let  $B_1, B_2, \dots, B_m$  be the simple direct summands of  $E_2^{0,*}$  and let  $A_1, A_2, \dots, A_n$  be the simple direct summands of  $E_2^{*,0}$ .

Since  $E$  is a central subgroup, we get

$$H^i(P/E, H^j(E, k)) \cong H^i(P/E, k) \bigotimes_k H^j(E, k) \cong E_2^{i,0} \otimes E_2^{0,j},$$

and  $E_2^{i,j}$  is a direct sum of tensorproducts  $A_a \otimes B_b$ . If we let the  $p'$ -group  $Q$  act diagonally on  $E_r^{i,0} \otimes E_r^{0,j}$ , then the map  $E_r^{i,0} \otimes E_r^{0,j} \xrightarrow{\cong} E_r^{i,j}$  is a  $kQ$ -module homomorphism.

First we prove that there is a simple  $kQ$ -module  $A_s$  depending on  $A$  such that  $A_s$  is a direct summand of  $E_\infty^{i',0}$ . Secondly we show that  $A$  is infinitely often a

direct summand in the image of the map

$$E_\infty^{i',0} \otimes E_\infty^{0,*} \xrightarrow{\cup} E_\infty^{i',*}.$$

(1) Let  $i'$  be the smallest  $i$  such that  $A$  is a direct summand in some tensorproduct  $A_s \otimes B_u$  with  $A_s \subseteq E_2^{i',0}$  and  $B_u \subseteq E_2^{0,*}$ . We show that  $A_s$  is a direct summand in  $E_\infty^{i',0}$ :

If  $A_s$  lies in the image of the differential  $d_r : E_r^{i'-r-1,r} \rightarrow E_r^{i',0}$ , then  $A_s$  is a direct summand in some tensorproduct  $A_t \otimes B_v$  with

$$A_t \subseteq E_2^{i'-r-1,0} \quad \text{and} \quad B_v \subseteq E_2^{0,r}.$$

The module  $A$  is then a direct summand in the tensorproduct  $(A_t \otimes B_v) \otimes B_u = A_t \otimes (B_v \otimes B_u)$ .

But  $B_v \otimes B_u = \bigoplus_w B_w$  and therefore  $A$  is a direct summand in  $A_t \otimes B_w$ . Since  $B_w$  is a direct summand of  $E_2^{0,*}$  it follows that  $A$  is a direct summand of  $E_2^{i'-r-1,*}$ . This contradicts the minimality of  $i'$ . Hence  $A_s$  is a direct summand of  $E_\infty^{i',0}$ .

(2) By Lemma 4.2 and Lemma 4.3  $B_u$  is infinitely often a direct summand in the image of the restriction map

$$\text{res} : H^*(P, k) \rightarrow H^*(E, k) \quad \text{i.e.}$$

$B_u$  is infinitely often a direct summand of  $E_\infty^{0,*}$ . Hence  $A$  is infinitely often a direct summand in  $E_\infty^{i',0} \otimes E_\infty^{0,*}$ .

If  $A$  is contained in the kernel of the map  $E_\infty^{i',0} \otimes E_\infty^{0,*} \xrightarrow{\cup} E_\infty^{i',*}$ , then  $A$  lies in the image of some differential  $d_r : E_r^{i'-r-1,*} \rightarrow E_r^{i',*}$ .

This contradicts the minimality of  $i'$ . It thus follows that  $A$  is infinitely often a direct summand of  $E_\infty^{i',*}$ .

**THEOREM 4.5.** *If the  $p'$ -group  $Q$  acts faithfully on the  $p$ -group  $P$ , then every simple  $kQ$ -module  $A$  is infinitely often a direct summand of  $H^*(P, k)$ .*

*Proof.* Let us consider the lower central series of  $P$

$$P = P^{(0)} \trianglerighteq P^{(1)} \trianglerighteq \dots \trianglerighteq P^{(m)} = e.$$

We obviously can refine this series to a central series

$$P = \tilde{P}^{(0)} \trianglerighteq \tilde{P}^{(1)} \trianglerighteq \dots \trianglerighteq \tilde{P}^{(n)} = e,$$

with elementary abelian factor groups  $\tilde{P}^{(i)}/\tilde{P}^{(i+1)}$  and  $\tilde{P}^{(0)}/\tilde{P}^{(1)} = P/\Phi(P)$ .

If  $Q$  acts faithfully on  $P$ ,  $Q$  acts faithfully on  $P/\Phi(P)$ , see for example [5] p. 102.

By Lemma 4.1 every simple  $kQ$ -module  $A$  is infinitely often a direct summand of  $H^*(P/\tilde{P}^{(1)}, k)$ . By Theorem 4.4  $A$  is infinitely often a direct summand of  $H^*(P/\tilde{P}^{(2)}, k)$ . Iterating this step for the factor groups  $P/\tilde{P}^{(i)}$  yields the result that  $A$  is infinitely often a direct summand of  $H^*(P, k)$ .

**THEOREM 4.6.** *Let  $G$  be a group of  $p$ -length 1, and let  $A$  be a simple  $kG$ -module lying in the principal block of  $kG$ . Then*

$$H^n(G, A) \neq 0 \quad \text{for infinitely many } n \in \mathbb{N}.$$

*Proof.* Let  $A^*$  denote the dual of  $A$ . By Theorem 4.5  $A^*$  is infinitely often a direct summand of  $H^*(P, k)$ . Dualisation yields the fact that  $A$  is infinitely often a direct summand of  $H_*(P, k)$ . By Theorem 3.1  $H^n(G, A)$  is nontrivial for infinitely many  $n \in \mathbb{N}$ .

## 5. The case $G = O_{p'pp'p}G$

**LEMMA 5.1.** *Let  $N \rightarrowtail G \twoheadrightarrow P$  be a group extension with  $|P| = p^a$ ;  $a \in \mathbb{N}$ , and let  $A$  be a  $kG$ -module. Then*

$$H^n(N, A) \neq 0 \Rightarrow H^n(G, A) \neq 0.$$

*Proof.* We consider the long exact sequence [3] p. 224

$$\rightarrow H^n(G, A) \rightarrow H^n(N, A) \rightarrow \text{Ext}_G^n(IP, A) \rightarrow$$

where  $IP$  denotes the augmentation ideal of the factor group  $P$ . Let  $IP^*$  denote the dual of  $IP$ ; then there is a natural isomorphism

$$\text{Ext}_G^n(IP, A) \cong H^n(G, IP^* \otimes_k A).$$

Since  $P$  is a  $p$ -group, all composition factors of  $IP^*$  are isomorphic to the trivial module  $k$ . A composition series of  $IP^*$  induces a composition series of  $IP^* \otimes_k A$ , of which all composition factors are isomorphic to  $A$ .

If  $H^n(G, IP^* \otimes_k A)$  is nontrivial, it follows by induction, that  $H^n(G, A)$  is nontrivial.

From  $H^n(N, A) \neq 0$  and from the above sequence we may conclude that  $H^n(G, A)$  or  $H^n(G, IP^* \otimes_k A)$  and hence again  $H^n(G, A)$  is nontrivial.

**LEMMA 5.2.** *Let  $G$  be a  $p$ -solvable group with normal subgroup  $N$ , and let  $A$  be a simple  $kG$ -module lying in the principal block of  $kG$ . Then:*

- (i)  $A = \bigoplus_{i=1}^m B_i$  as a  $kN$ -module and all  $B_i$  are simple  $kN$ -modules.
- (ii) The simple  $kN$ -modules  $B_i$  lie in the principal block of  $kN$ .

*Proof.* (i) is a consequence of Clifford's theorem.

(ii) For  $p$ -solvable groups the following holds [2] p. 279

$$C_G(A) \supseteq O_{p'p}G \Leftrightarrow A \text{ lies in the principal block of } kG$$

Since  $O_{p'p}G$  is the maximal  $p$ -nilpotent normal subgroup of  $G$ ,  $O_{p'p}N$  is a subgroup of  $O_{p'p}G$ . Therefore we get  $O_{p'p}N \subseteq O_{p'p}G \subseteq C_G(A)$ , and thus all  $B_i$  are simple  $kN$ -modules lying in the principal block of  $kN$ .

**THEOREM 5.3.** *Let  $G = O_{p'pp'p}G$ , and let  $A$  be a simple  $kG$ -module lying in the principal block of  $kG$ . Then*

$$H^n(G, A) \neq 0 \text{ for infinitely many } n \in \mathbb{N}.$$

*Proof.* We consider the extension

$$O_{p'pp'}G \rightarrowtail G \twoheadrightarrow G/O_{p'pp'}G.$$

The factor groups  $G/O_{p'pp'}G$  is a  $p$ -group, and the normal subgroup  $O_{p'pp'}G$  has  $p$ -length 1.

By Lemma 5.2  $A$  is a direct sum of simple  $k(O_{p'pp'}G)$ -modules  $B_i$  lying in the principal block of  $k(O_{p'pp'}G)$ . By Theorem 4.6  $H^n(O_{p'pp'}G, B_i)$  is nontrivial for infinitely many  $n \in \mathbb{N}$ , and Lemma 5.1 yields

$$H^n(G, A) \neq 0 \text{ for infinitely many } n \in \mathbb{N}.$$

## REFERENCES

- [1] EVENS, L., *The spectral sequence of a finite group extension stops*. Trans. amer. math. soc., vol. 210 (1975), p. 269–277.
- [2] FONG, P., *On the characters of  $p$ -solvable groups*. Trans. amer. math. soc., vol. 98, p. 263–284.
- [3] HILTON, P. J. and STAMMBACH, U., *A course in Homological Algebra*. Springer 1971.

- [4] HOCHSCHILD, G. P. and SERRE, J. P., *Cohomology of group extensions*. Trans. amer. math. soc., vol. 74 (1953), p. 110–135.
- [5] KURZWEIL, H., *Endliche Gruppen*. Springer 1977.
- [6] PUTTASWAMAIAH, B. M. and DIXON, J. D., *Modular Representations of Finite Groups*. Academic Press 1977.
- [7] STEINBERG, R., *Complete sets of representations of algebras*. Proc. amer. math. soc. Vol. 13, p. 746–747 (1962).

*Eidgenössische Technische Hochschule  
Abt. Mathematik  
Zentrum  
CH-8092 Zürich/Switzerland*

Received July 26, 1982