

On small eigenvalues of the Laplacian for ... (q) \... .

Autor(en): **Stopple, Jeffrey**

Objekttyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **65 (1990)**

PDF erstellt am: **28.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-49748>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

On small eigenvalues of the Laplacian for $\Gamma_0(q) \backslash \mathcal{H}$

JEFFREY STOPPLE

Let \mathcal{H} be the complex upper half plane, and $\Gamma_0(q)$ be the subgroup of matrices

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

in $SL(2, \mathbb{Z})$ with $c \equiv 0 \pmod{q}$. Suppose f is a Maass cusp form with eigenvalue λ ; i.e., a non-constant function $f: \mathcal{H} \rightarrow \mathbb{C}$ satisfying

$$f(\gamma z) = f(z) \quad \text{for } \gamma \text{ in } \Gamma_0(q), z \text{ in } \mathcal{H}$$

$$y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) f + \lambda f = 0$$

$$\int_{\mathcal{F}} |f(z)|^2 \frac{dx \, dy}{y^2} < \infty$$

where \mathcal{F} is a fundamental domain for $\Gamma_0(q)$. Selberg conjectured [8] that $\lambda \geq 1/4$, and showed that $\lambda \geq 3/16$. Iwaniec has a statistical result [5] that shows the rarity of small eigenvalues, similar to density theorems about real zeros of Dirichlet's L -series.

For an odd prime q , whether q is ramified, split, or inert in a real quadratic field $\mathbb{Q}(\sqrt{d})$ depends only on the Legendre symbol (d/q) , and so is periodic in $d \pmod{q}$. If we consider instead the set of all norm 1 units $\epsilon > 1$ in all real quadratic fields, ordered by the size of ϵ , we expect the behavior of q in $\mathbb{Q}(\epsilon)$ to be more or less random. We use the trace formula to show that if there is an eigenvalue λ less than $1/4$ then that expectation is wrong; instead q has a bias towards a certain behavior.

Specifically let $t \geq 3$ in \mathbb{Z} , and write $t^2 - 4 = u^2 d$ with d a discriminant of a real quadratic field. Then $\mathbb{Q}(\sqrt{d}) = \mathbb{Q}(\epsilon)$ for ϵ the larger root of $x^2 - tx + 1 = 0$. Let $h(d)$ be the narrow class number and ϵ_1 the fundamental

norm 1 unit. Define

$$\Theta(t) = \begin{cases} q & \text{if } t \equiv \pm 2 \pmod{q^2} \\ \left(\frac{\Delta}{q}\right) & \text{otherwise} \end{cases}$$

If λ is the smallest eigenvalue; $\lambda = 1/4 + r^2$ with $r = i\rho$ purely imaginary, $0 < \rho < 1/2$ then for $T \rightarrow \infty$ we have

THEOREM.

$$\frac{1}{\sqrt{\pi T}} \sum_{t \geq 3} \frac{2h(\Delta) \log(\epsilon_1)}{\sqrt{\Delta}} \sum_{m \mid u} \left(\frac{\Delta}{m}\right) \sigma(u/m) \mu(m) u^{-1} \Theta(t) \exp(-\log^2(t)/T) \sim \exp(\rho^2 T)$$

Here $\sigma(n) = \sum_{d \mid n} d$ and $\mu(n)$ is the Möbius function. In the course of proving this formula we will see that

$$\sum_{m \mid u} \left(\frac{\Delta}{m}\right) \sigma(u/m) \mu(m) > 0;$$

one can show that it is less than $\sigma(u)$. From [2], Theorem 322 we know that

$$\frac{\sigma(u)}{u} = O(u^\kappa) = O(t^\kappa)$$

for any $\kappa > 0$. By the Brauer–Siegel theorem and the fact that $\Delta \leq t^2 - 4$,

$$t^{-\kappa} \leq \frac{2h(\Delta) \log(\epsilon_1)}{\sqrt{\Delta}} \leq t^\kappa$$

for any $\kappa > 0$, for Δ sufficiently large. Thus if one expects that (Δ/q) is random for $\Delta = \Delta(t)$ as t increases, then there should be cancellation in the sum and it will not grow like $\exp(\rho^2 T)$, so $\lambda < 1/4$ will not occur.

The proof of the theorem depends, of course, on the trace formula. Let f be a Maass cusp form corresponding to the eigenvalue $\lambda < 1/4$. Then for γ in $\Gamma = SL(2, \mathbb{Z})$, $f(\gamma z)$ is also a Maass cusp form for the same eigenvalue, so Γ acts in this finite dimensional space. The principal congruence subgroup $\Gamma(q)$ acts trivially giving a representation π of the factor group $G = \Gamma/\Gamma(q)$, (isomorphic to $SL(2, \mathbb{F}_q)$). Let $B = \Gamma_0(q)/\Gamma(q)$, then since f is fixed by $\Gamma_0(q)$, the multiplicity of the trivial representation in the restriction of π to B is ≥ 1 . By Frobenius Reciprocity, the multiplicity of π in the induced representation $\text{Ind}_B(1)$ is also ≥ 1 . This is a $q + 1$ dimensional representation isomorphic to the space of functions

$$\{f : B \backslash G \rightarrow \mathbb{C}\}$$

where the group G acts by right translation

$$\text{Ind}_B(1)(g)f(Bx) = f(Bxg).$$

By Mackey's Irreducibility Criterion (see e.g. [9] p. 59) the induced representation has two irreducible components,

$$\text{Ind}_B(1) = 1 \oplus \theta.$$

Here 1 is the trivial representation of G and θ is realized in the q dimensional subspace orthogonal to the constant functions; i.e. in the space of functions

$$\{f : B \backslash G \rightarrow \mathbb{C} \mid \sum f(Bg) = 0\}.$$

Since f is not fixed by $\Gamma = SL(2, \mathbb{Z})$ ($\lambda > 1/4$ is known), the projection of f on the space isomorphic to that of θ is not 0. Thus there exist cusp forms which transform according to θ . Then by [4] ((16) on page 182), λ is an eigenvalue for the Laplacian acting in the space of vector valued Maass cusp forms for Γ with multiplier θ :

$$F : \mathcal{H} \rightarrow \mathbb{C}^q \quad \text{such that} \quad F(\gamma z) = \theta(\gamma)F(z).$$

We briefly recall the trace formula for such forms, as in Theorem 4.2 in [3], page 315. Let

$$g(u) = \exp(-u^2/4T)/\sqrt{4\pi T}, \quad \text{and} \quad h(r) = \exp(-r^2T)$$

its Fourier transform be our test functions. The eigenvalues of the Laplacian

$\lambda_n = 1/4 + r_n^2$ are related to the primitive hyperbolic conjugacy classes $\{P\}$ by

$$\begin{aligned} \sum_n h(r_n) &= \sum_{\{P\}} \sum_{k=1}^{\infty} \frac{\text{Trace } \theta(P^k) \log(NP)}{NP^{k/2} - NP^{-k/2}} g(\log(NP^k)) \\ &\quad + \int_{-\infty}^{\infty} h(r) \{\text{contribution from central class}\} dr \\ &\quad + \int_{-\infty}^{\infty} h(r) \{\text{contribution from elliptic classes}\} dr \\ &\quad + \int_{-\infty}^{\infty} h(r) \{\text{parabolic contribution to continuous spectrum}\} dr \\ &\quad + g(0) \{\text{parabolic contribution to discrete spectrum}\} \\ &\quad + h(0) \{\text{parabolic contribution to discrete spectrum}\}. \end{aligned}$$

All sums and integrals are absolutely convergent. Recall that $NP = \epsilon^2$ where ϵ is the larger root of the characteristic polynomial of P . By the Dominated Convergence Theorem, we have, as $T \rightarrow \infty$,

$$\int_{-\infty}^{\infty} h(r) \{*\} dr \rightarrow 0.$$

The terms with $h(0)$ and $g(0)$ are $O(1)$ as $T \rightarrow \infty$. We next consider the terms from the spectral side. For all but finitely many n , say $n > N$ we have $\lambda_n > 1/4$ so r_n is in \mathbb{R} . Thus

$$\sum_{n>N} h(r_n) \rightarrow 0$$

again by the Dominated Convergence Theorem. The finitely many eigenvalues less than $1/4$ have r_n purely imaginary. Note that the contribution of the smallest such eigenvalue dominates the others, and 0 does not occur as an eigenvalue as θ is a nontrivial representation. Thus as $T \rightarrow \infty$ we have

$$\frac{1}{\sqrt{4\pi T}} \sum_{\{P\}} \sum_{k=1}^{\infty} \frac{\text{Trace } \theta(P^k) \log(NP)}{NP^{k/2} - NP^{-k/2}} \exp(-\log^2(NP^k)/4T) \sim \exp(\rho^2 T). \quad (1)$$

We define the usual map ϕ from matrices to binary quadratic forms

$$\phi : \begin{bmatrix} a & b \\ c & d \end{bmatrix} \rightarrow \left[\begin{bmatrix} b & d-a \\ v & v \end{bmatrix}, \frac{-c}{v} \right]$$

where $v = \gcd(b, d - a, c)$. A form $[\alpha, \beta, \gamma]$ with discriminant D is the image of

$$\begin{bmatrix} \frac{t - u\beta}{2} & u\alpha \\ -u\gamma & \frac{t + u\beta}{2} \end{bmatrix}.$$

where $t^2 - u^2D = 4$ is the fundamental solution to Pell's equation for D . Sarnak shows in Proposition 1.4 of [7] that ϕ is a $2 - 1$ map and commutes with the action of the modular group giving a $2 - 1$ correspondence between primitive hyperbolic conjugacy classes and equivalence classes of indefinite binary quadratic forms. For a $\{P\}$ corresponding to a class $\{\phi\}$ of discriminant D , write $D = d^2\Delta$, with Δ the discriminant of a real quadratic field. We will often suppress Δ from the notation. Note that NP is the square of the larger root of the characteristic polynomial and so only depends on the discriminant D . Write

$$NP = \epsilon_d^2 \quad \text{for} \quad \epsilon_d = \frac{t_d + u_d \sqrt{d^2\Delta}}{2}.$$

We next analyze $\text{Trace } \theta$. We will show that $\text{Trace } \theta(P^k)$ depends only on the characteristic polynomial of P^k which is the minimum polynomial $x^2 - tx + 1$ of ϵ_d^k . First suppose

$$t \equiv \pm 1 \pmod{q^2}.$$

Sarnak ([7], Proposition 3.3) shows that

$$P^k \equiv \pm I \pmod{q} \Leftrightarrow q \mid u \Leftrightarrow q^2 \mid t^2 - 4 \Leftrightarrow t \equiv \pm 2 \pmod{q^2}.$$

Tables ([1] vol. IV, p. 1829) show this is the case when $\text{Trace } \theta$ is equal to q . Now suppose

$$t \not\equiv \pm 2 \pmod{q}.$$

From the tables $\text{Trace } \theta = 1$ if and only if the matrix element $P^k \pmod{q}$ is diagonalizable over \mathbb{F}_q but not central. This occurs if and only if the discriminant $t^2 - 4 = u^2\Delta$ is a square in \mathbb{F}_q^* ; i.e., $(\Delta/q) = 1$. Similarly $\text{Trace } \theta = -1$ if and only if the matrix element $P^k \pmod{q}$ is not diagonalizable over \mathbb{F}_q but is diagonalizable over \mathbb{F}_{q^2} . This occurs when the discriminant $t^2 - 4 = u^2\Delta$ is not a square in \mathbb{F}_q^* ; i.e.,

$(\Delta/q) = -1$. Finally if

$$t \equiv \pm 1 \pmod{q} \quad \text{but not } \pmod{q^2},$$

then

$$t^2 - 4 \equiv 0 \pmod{q} \quad \text{but not } \pmod{q^2}.$$

By the above, $q \nmid u$, so $q \mid \Delta$. Also, $x^2 - tx + 1$ has repeated roots \pmod{q} , but $P^k \pmod{q}$ is not central. Thus P^k is unipotent \pmod{q} and the table shows that $\text{Trace } \theta(t) = 0 = (\Delta/q)$. This shows that the Trace θ depends only on the characteristic polynomial and is equal to the function Θ defined above. (Since the determinant is always 1 only the x coefficient, yet another trace, matters.) It will be convenient to write $\Theta(\epsilon_d^k)$.

This gives the formula

$$\frac{1}{\sqrt{\pi T}} \sum_{D=d^2\Delta} \sum_{k=1}^{\infty} 2h(d^2\Delta) \log(\epsilon_d) \frac{\Theta(\epsilon_d^k)}{\epsilon_d^k - \epsilon_d^{-k}} \exp(-\log^2(\epsilon_d^k)/T) \sim \exp(\rho^2 T). \quad (2)$$

The class number $h(d^2\Delta)$ of forms is related to the ideal class number $h(\Delta)$ of $\mathbb{Q}(\sqrt{\Delta})$ by formula (see e.g. [6] p. 95)

$$h(d^2\Delta) = \frac{h(\Delta)d}{[O_\Delta^* : \mathbb{Z}[\epsilon_d]^*]} \prod_{l \mid d} \left(1 - \frac{\left(\frac{\Delta}{l}\right)}{l} \right).$$

And for ϵ_1 the fundamental norm 1 unit in $\mathbb{Q}(\sqrt{\Delta})$ we have

$$\epsilon_d = \epsilon_1^{[O_\Delta^* : \mathbb{Z}[\epsilon_d]^*]}.$$

In fact this follows the definition of ϵ_1 and $\mathbb{Z}[\epsilon_d]$. Substituting this in (2) gives

$$\begin{aligned} \frac{1}{\sqrt{\pi T}} \sum_{D=d^2\Delta} 2h(\Delta) \log(\epsilon_1) d \prod_{l \mid d} \left(1 - \frac{\left(\frac{\Delta}{l}\right)}{l} \right) \sum_{k=1}^{\infty} \frac{\Theta(\epsilon_d^k)}{\epsilon_d^k - \epsilon_d^{-k}} \exp(-\log^2(\epsilon_d^k)/T) \\ \sim \exp(\rho^2 T) \quad (3) \end{aligned}$$

Still viewing Δ as fixed we want to group all terms of the form $\epsilon_d^k = \epsilon_1^n$.

We have

$$\epsilon_d = \frac{t_d + u_d \sqrt{d^2 \Delta}}{2}$$

and suppose

$$\epsilon_1^n = \frac{t(n) + u(n) \sqrt{\Delta}}{2}$$

then

$$\epsilon_d^k - \epsilon_d^{-k} = \epsilon_1^n - \epsilon_1^{-n} = u(n) \sqrt{\Delta}.$$

We combine the infinite sum on d and k to sum on n and $d \mid u(n)$ to get

$$\begin{aligned} \frac{1}{\sqrt{\pi T}} \sum_{\Delta} \frac{2h(\Delta) \log(\epsilon_1)}{\sqrt{\Delta}} \sum_{n=1}^{\infty} \sum_{d \mid u(n)} \frac{d}{u(n)} \prod_{l \mid d} \left(1 - \frac{\left(\frac{\Delta}{l}\right)}{l}\right) \Theta(\epsilon_1^n) \exp(-\log^2(\epsilon_1^n)/T) \\ \sim \exp(\rho^2 T) \quad (4) \end{aligned}$$

Now

$$\begin{aligned} \sum_{d \mid u(n)} \frac{d}{u(n)} \prod_{l \mid d} \left(1 - \frac{\left(\frac{\Delta}{l}\right)}{l}\right) &= \sum_{d \mid u(n)} \frac{d}{u(n)} \sum_{m \mid d} \frac{\mu(m)}{m} \left(\frac{\Delta}{m}\right) \\ &= \frac{1}{u(n)} \sum_{dd' = u(n)} \sum_{mm' = d} m' \mu(m) \left(\frac{\Delta}{m}\right) \\ &= \frac{1}{u(n)} \sum_{m \mid u(n)} \mu(m) \left(\frac{\Delta}{m}\right) \sum_{m' \mid u(n)/m} m' \\ &= \frac{1}{u(n)} \sum_{m \mid u(n)} \mu(m) \left(\frac{\Delta}{m}\right) \sigma \left(\frac{u(n)}{m}\right). \end{aligned}$$

From this we get that (4) is equal

$$\begin{aligned} \frac{1}{\sqrt{\pi T}} \sum_{\Delta} \frac{2h(\Delta) \log(\epsilon_1)}{\sqrt{\Delta}} \sum_{n=1}^{\infty} \sum_{m \mid u(n)} \mu(m) \left(\frac{\Delta}{m}\right) \sigma \left(\frac{u(n)}{m}\right) \frac{\Theta(\epsilon_1^n)}{u(n)} \exp(-\log^2(\epsilon_1^n)/T) \\ \sim \exp(\rho^2 T). \quad (5) \end{aligned}$$

To get the theorem we now need to reorder the terms in the sum. We have a sum over all real quadratic fields, and over all positive powers of the fundamental norm 1 unit of that field. These units are in 1–1 correspondence with their minimum polynomial $x^2 - tx + 1$ ordered by their trace t . Note that Θ depends only on t above. As the units $\epsilon_1^n \rightarrow \infty$; we have $\epsilon_1^{-n} \rightarrow 0$, and since $t = \epsilon_1^n + \epsilon_1^{-n}$ we can replace ϵ_1^n with t in the statement of the theorem.

The authors would like to thank Morris Newman for correcting some mistakes in an earlier version of this paper.

REFERENCES

- [1] Mathematical Society of Japan, *Encyclopedic Dictionary of Mathematics*, MIT Press.
- [2] G. H. HARDY and E. M. WRIGHT, *An Introduction to the Theory of Numbers*, Oxford University Press.
- [3] D. HEJHAL, The Selberg Trace Formula for $\mathrm{PSL}(2, R)$, II, Lecture Notes in Mathematics 1001.
- [4] H. HUBER, Über die Darstellungen der Automorphismengruppe einer Riemannschen Fläche in die Eigenräume des Laplace-operators. Comment. Math. Helvetici 52, 177–184 (1977).
- [5] H. IWANIEC, Character sums and small eigenvalues for $\Gamma_0(p)$, Glasgow Mathematical Journal 27, 99–116.
- [6] S. LANG, *Elliptic Functions*, Springer Verlag.
- [7] P. SARNAK, Class number of indefinite binary quadratic forms, J. Number Theory 15, 229–247.
- [8] A. SELBERG, On the estimation of Fourier coefficients of modular forms, Proc. Sympos. Pure Math. 8, 1–15.
- [9] J. P. SERRE, *Linear Representations of Finite Groups*, Springer Verlag.

*Mathematics Department
University of California
Santa Barbara, CA 93106/USA*

Received January 30, 1990