

On a simplicial complex associated with tilting modules.

Autor(en): **Riedtmann, Ch., Schofield, Aidan**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **66 (1991)**

PDF erstellt am: **28.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-50391>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

On a simplicial complex associated with tilting modules

CHRISTINE RIEDTMANN AND AIDAN SCHOFIELD

Introduction

Let A be a finite-dimensional associative algebra over an algebraically closed field, and denote by $\text{mod } A$ the category of finite-dimensional A -modules. We fix the number of pairwise non-isomorphic simple A -modules to be $n + 1$.

Denote by \mathcal{E} a set of fixed representatives for the isomorphism classes of indecomposable A -modules T satisfying the following conditions:

- (i) The projective dimension of T is at most 1.
- (ii) T does not extend itself, i.e. $\text{Ext}_A^1(T, T) = 0$.

Following Ringel, we define a simplicial complex \mathcal{C}_A on the set \mathcal{E} of vertices: (T_0, \dots, T_r) is an r -simplex if $\text{Ext}_A^1(T_0 \oplus \dots \oplus T_r, T_0 \oplus \dots \oplus T_r) = 0$. Ringel told us that \mathcal{C}_A is a triangulated ball for certain hereditary algebras. Our goal is to prove the following result:

THEOREM. *If \mathcal{E} is finite, the geometric realization of \mathcal{C}_A is an n -dimensional ball.*

We wish to thank C. Ringel for drawing our attention to \mathcal{C}_A and N. A'Campo for discussing with us the topological aspects of the question.

1. The Bongartz completion

1.1. Recall from [3], [5] that a A -module T is a *tilting module* if it satisfies:

- (i) $\text{projdim}_A T \leq 1$,
- (ii) $\text{Ext}_A^1(T, T) = 0$,
- (iii) There is an exact sequence

$$0 \rightarrow A \rightarrow T' \rightarrow T'' \rightarrow 0,$$

with modules T', T'' that belong to the full subcategory $\text{add } T$ of $\text{mod } A$ whose objects are direct summands of T^N for some N .

The simplest example of a tilting module is A itself, and for some algebras, e.g. the selfinjective ones, there are no others (aside from those obtained by changing the multiplicities of the indecomposable direct summands). Bongartz proved in [2] that a module T satisfying (i) and (ii) is a tilting module if and only if the number of its pairwise non-isomorphic indecomposable direct summands equals the number $n + 1$ of isomorphism classes of simple modules. He also showed that any module T satisfying (i) and (ii) is a direct summand of a tilting module. We recall his construction: write $T = \bigoplus_{i=0}^r T_i^{\lambda_i}$ as a direct sum of pairwise non-isomorphic indecomposables T_0, \dots, T_r with multiplicities $\lambda_0, \dots, \lambda_r$. Choose an exact sequence

$$0 \rightarrow A \rightarrow X \rightarrow \bigoplus_{i=0}^r T_i^{\mu_i} \rightarrow 0$$

with the property that, for any $k = 0, \dots, r$, the induced map

$$\text{Hom}_A \left(T_k, \bigoplus_{i=0}^r T_i^{\mu_i} \right) \rightarrow \text{Ext}_A^1(T_k, A), \quad (*)$$

is surjective. Then $T \oplus X$ is the desired tilting module.

Of course the condition $(*)$ does not determine X uniquely. But it is easy to see that possible choices for X only differ by direct summands in $\text{add } T$, up to isomorphism. Hence T determines a multiplicity-free tilting module $\tilde{T} = \bigoplus_{i=0}^n T_i$, which is unique up to isomorphism. We call $T_B = T_{r+1} \oplus \dots \oplus T_n$ the *Bongartz completion* of T .

1.2. Let T_0, \dots, T_n be pairwise non-isomorphic indecomposables, and suppose that $\bigoplus_{i=0}^n T_i$ is a tilting module.

PROPOSITION. *The following statements are equivalent:*

- (a) $\bigoplus_{i=r+1}^n T_i$ is the Bongartz completion of $\bigoplus_{i=0}^r T_i$.
- (b) For $j = r + 1, \dots, n$, there is no surjection from any module in $\text{add } (T_0 \oplus \dots \oplus T_{j-1} \oplus T_{j+1} \oplus \dots \oplus T_n)$ to T_j .

Proof. Let $\bigoplus_{i=r+1}^n T_i$ be the Bongartz completion of $\bigoplus_{i=0}^r T_i$, and suppose there is a surjection $f: \bigoplus_{i \neq j} T_i^{\nu_i} \rightarrow T_j$ for some $j > r$. Consider the following commutative diagram:

$$\begin{array}{ccccccc}
0 \longrightarrow \Lambda & \longrightarrow & T_j^{\rho_j} \oplus \bigoplus_{i \neq j} T_i^{\rho_i} & \longrightarrow & \bigoplus_{i=0}^r T_i^{\mu_i} & \longrightarrow & 0 \\
\parallel & & \uparrow \begin{bmatrix} f & 0 \\ 0 & f \end{bmatrix} & & \parallel & & \uparrow \\
0 \longrightarrow \Lambda & \xrightarrow{g} & \left(\bigoplus_{i \neq j} T_i^{\nu_i} \right)^{\rho_j} \oplus \bigoplus_{i \neq j} T_i^{\rho_i} & \longrightarrow & X & \longrightarrow & 0.
\end{array}$$

The first row is an exact sequence used to construct the Bongartz completion, and the existence of g follows from the projectivity of Λ . The square on the right yields another exact sequence:

$$0 \rightarrow \bigoplus_{i \neq j} T_i^{\nu_i \rho_j + \rho_i} \rightarrow \bigoplus_{i=0}^n T_i^{\rho_i} \oplus X \rightarrow \bigoplus_{i=0}^r T_i^{\mu_i} \rightarrow 0,$$

which must split. But then T_j is isomorphic to some T_i for $i \neq j$, and this is impossible.

As to the converse, we choose an exact sequence

$$0 \rightarrow \Lambda \rightarrow \bigoplus_{i=0}^n T_i^{\alpha_i} \xrightarrow{h} \bigoplus_{i=0}^n T_i^{\beta_i} \rightarrow 0.$$

For any $j > r$ with $\beta_j > 0$, the composition of h with the canonical projection from $\bigoplus_{i=0}^n T_i^{\beta_i}$ to $T_j^{\beta_j}$ must be retraction by (b). So we can choose another such sequence with $\beta_j = 0$ for $j > r$. As our sequence then satisfies (*), $\bigoplus_{i=r+1}^n T_i$ must be the Bongartz completion of $\bigoplus_{i=0}^r T_i$.

Remark. The same arguments show that $T = \bigoplus_{i=0}^n T_i$ is a projective tilting module if and only if there is no surjection from any modules in $\text{add}(T_0 \oplus \cdots \oplus T_{j-1} \oplus T_{j+1} \oplus \cdots \oplus T_n)$ to T_j , for $j = 0, \dots, n$.

1.3. Let T_0, \dots, T_{n-1} be pairwise non-isomorphic indecomposables of projective dimension 1 at most, and assume that $\text{Ext}_{\Lambda}^1(T, T) = 0$ for $T = \bigoplus_{i=0}^{n-1} T_i$. Denote by T_n the Bongartz completion of T .

The following result has been obtained independently by Happel in [4]. In case Λ is hereditary, it was proved in [7] and later in [6].

PROPOSITION. *There is at most one indecomposable T'_n not isomorphic to T_n such that $T \oplus T'_n$ is a tilting module. If such a T'_n exists, there is an exact sequence*

$$0 \rightarrow T_n \rightarrow \bigoplus_{i=0}^{n-1} T_i^{\lambda_i} \rightarrow T'_n \rightarrow 0.$$

We first have to recall the definitions of a source map and a sink map used in [7]. Closely related concepts have been introduced in [1]. Let X_1, \dots, X_r be pairwise non-isomorphic indecomposables and let Y be a module not having any direct summands in $\text{add } X$, where $X = \bigoplus_{i=1}^r X_i$.

A map $f: Y \rightarrow \bigoplus_{i=1}^r X_i^{\lambda_i}$ is a *source map* from Y to $\text{add } X$ if

- (i) for any X' in $\text{add } X$, any map from Y to X' factors through f , and
- (ii) f is minimal with respect to property (i); i.e. if $\alpha \circ f$ still has property (i) for an endomorphism α of $\bigoplus_{i=1}^r X_i^{\lambda_i}$, then α is an automorphism.

Source maps exist and are unique up to isomorphism. If a map $g: Y \rightarrow \bigoplus_{i=1}^r X_i^{\mu_i}$ has property (i), it is isomorphic to $\begin{bmatrix} f \\ 0 \end{bmatrix}: Y \rightarrow \bigoplus_{i=1}^r X_i^{\lambda_i} \oplus X'$ for any source map f , where X' lies in $\text{add } X$.

Sink maps from $\text{add } X$ to Y are defined by dualizing the definition of source maps.

Proof of the proposition. Let T'_n be an indecomposable not isomorphic to T_n such that $T \oplus T'_n$ is a tilting module. By the preceding proposition, there is a surjection from some module in $\text{add } T$ to T'_n . In particular, any sink map

$$g: \bigoplus_{i=0}^{n-1} T_i^{\lambda_i} \rightarrow T'_n,$$

from $\text{add } T$ to T'_n is surjective. Consider the exact sequence

$$0 \rightarrow Z \xrightarrow{f} \bigoplus_{i=0}^{n-1} T_i^{\lambda_i} \xrightarrow{g} T'_n \rightarrow 0,$$

where $Z = \ker g$.

Since g is a sink map, f lies in the radical of $\text{mod } \Lambda$; i.e., its restriction to any indecomposable direct summand of Z is never a section. Moreover, any map from Z to T_j factors through f , since we have $\text{Ext}^1(T'_n, T_j) = 0$, for $j = 0, \dots, n-1$. Therefore Z has no direct summand that belongs to $\text{add } T$. As g lies in the radical of $\text{mod } \Lambda$, f is a source map from Z to $\text{add } T$.

Obviously the projective dimension of Z is 1 at most, and by construction we have $\text{Ext}_\Lambda^1(T_j, Z) = 0$, for $j = 0, \dots, n-1$. Considering maps from our sequence to Z and T_j , respectively, and using that $\text{projdim}_\Lambda T'_n \leq 1$, we find that $\text{Ext}_\Lambda^1(Z, Z) = 0$ and $\text{Ext}_\Lambda^1(T_j, Z) = 0$, for $j = 0, \dots, n-1$. As Z does not belong to $\text{add } T$, $T \oplus Z$ is a tilting module.

If there were a surjection from some T' in $\text{add } T$ to Z , it would induce a surjection from $\text{Ext}_\Lambda^1(T'_n, T')$ to $\text{Ext}_\Lambda^1(T'_n, Z)$, since $\text{projdim}_\Lambda T'_n \leq 1$. But this is impossible, as the first group is zero and our sequence does not split. By the preceding proposition, we know that Z is isomorphic to T_n^λ for some $\lambda \geq 1$, and we may suppose $Z = T_n^\lambda$.

We now want to show that $\lambda = 1$. Let $h : T_n \rightarrow T'$ be a source map from T_n to add T . The map

$$\begin{bmatrix} h & 0 \\ 0 & h \end{bmatrix} : T_n^\lambda \rightarrow T'^\lambda,$$

still has the first property of a source map, and it is therefore isomorphic to

$$\begin{bmatrix} f \\ 0 \end{bmatrix} : T_n^\lambda \rightarrow \bigoplus_{i=0}^{n-1} T_i^{\lambda_i} \oplus T'',$$

for some T'' in add T . Comparing cokernels, we find that $(\text{coker } h)^\lambda$ is isomorphic to $T'' \oplus T'_n$, which implies $\lambda = 1$, by Krull–Schmidt.

Finally, since $f : T_n \rightarrow \bigoplus_{i=0}^{n-1} T_i^{\lambda_i}$ is a source map, its cokernel T'_n is determined uniquely, up to isomorphism, by T_n . Our proposition is proved.

Remark. There exist modules T as in the proposition whose only completion is the Bongartz completion T_n . Indeed, if $\bigoplus_{i=0}^n P_i$ is a projective tilting module, at least one of the modules $\bigoplus_{i \neq j} P_i$ has this property, since chains of injections in the radical of mod Λ between projectives have bounded length.

2. Proof of the theorem

2.1. We associate a quiver K with the complex \mathcal{C}_Λ defined in the introduction in the following way: the vertices of K are the n -simplices of \mathcal{C}_Λ . For each $(n-1)$ -simplex (T_0, \dots, T_{n-1}) which is face of two n -simplices, K contains an arrow $\sigma = (T_0, \dots, T_n) \rightarrow \sigma' = (T_0, \dots, T_{n-1}, T'_n)$, where T_n is the Bongartz completion of $\bigoplus_{i=0}^{n-1} T_i$. For any simplex τ of \mathcal{C}_Λ , we let K_τ denote the full subquiver of K whose vertices are then n -simplices of \mathcal{C}_Λ containing τ .

LEMMA. *Let τ be a simplex of \mathcal{C}_Λ . If there is a path $\sigma_1 \rightarrow \sigma_2 \rightarrow \dots \rightarrow \sigma_s$ in K with σ_1, σ_s in K_τ , then the whole path lies in K_τ .*

Proof. Recall that, for a tilting module T , the category $\mathcal{T}(T)$ of torsion modules with respect to T is the full subcategory of mod Λ whose objects are quotients of T^N for some N . Set $\mathcal{T}(\sigma) = \mathcal{T}(\bigoplus_{i=0}^n T_i)$ for $\sigma = (T_0, \dots, T_n)$.

If K contains an arrow $\sigma = (T_0, \dots, T_n) \rightarrow \sigma' = (T_0, \dots, T_{n-1}, T'_n)$, there is an exact sequence

$$0 \rightarrow T_n \rightarrow \bigoplus_{i=0}^{n-1} T_i^{\lambda_i} \rightarrow T'_n \rightarrow 0,$$

by 1.3, and therefore any module in $\mathcal{T}(\sigma')$ belongs to $\mathcal{T}(\sigma)$. However by 1.2, T_n does not lie in $\mathcal{T}(\sigma')$. Moreover, for any path $\sigma \rightarrow \sigma' \rightarrow \cdots \rightarrow \sigma''$ in K , $\mathcal{T}(\sigma'')$ lies in $\mathcal{T}(\sigma')$ and thus does not contain T_n .

The lemma follows by applying these considerations to $\sigma = \sigma_k \rightarrow \sigma' = \sigma_{k+1} \rightarrow \cdots \rightarrow \sigma'' = \sigma_s$ in case $\sigma_1 \rightarrow \cdots \rightarrow \sigma_s$ does not lie in K_τ , where k is the maximal index for which $\sigma_1 \rightarrow \cdots \rightarrow \sigma_k$ is in K_τ . Then τ contains T_n , by the choice of k , but σ_s cannot.

2.2. Applying the lemma to an n -simplex we find:

PROPOSITION. *K does not contain oriented cycles.*

This allows us to define an *order relation* for the n -simplices of \mathcal{C}_A : $\sigma \leq \sigma'$ if there is an oriented path $\sigma = \sigma_1 \rightarrow \sigma_2 \rightarrow \cdots \rightarrow \sigma_s = \sigma'$ in K .

Remarks. (a) The *Hasse diagram* of this order relation is the quiver whose vertices are the n -simplices of \mathcal{C}_A and which contains an arrow $\sigma \rightarrow \sigma'$ if $\sigma \leq \sigma'$, $\sigma \neq \sigma'$ and $\sigma \leq \sigma'' \leq \sigma'$ implies either $\sigma'' = \sigma$ or $\sigma'' = \sigma'$. Applying the lemma to an $(n-1)$ -simplex which is face of two n -simplices, it is easy to see that the Hasse diagram coincides with K .

(b) Our order relation is in general distinct from the one defined by: $\sigma \leq \sigma'$ if $\mathcal{T}(\sigma) \supseteq \mathcal{T}(\sigma')$. The projective and the injective tilting module of a hereditary algebra of infinite representation type furnish an example. We don't know, however, whether the Hasse diagrams coincide.

2.3. Suppose now that \mathcal{E} is finite. Number the n -simplices $\sigma_1, \sigma_2, \dots, \sigma_M$ of \mathcal{C}_A in such a way that $\sigma_i \leq \sigma_j$ implies $i \leq j$. For $N \leq M$, let \mathcal{B}_N be the union of $\sigma_1, \sigma_2, \dots, \sigma_N$.

The following proposition implies our theorem.

PROPOSITION. *The geometric realization of \mathcal{B}_N is an n -ball, for all N .*

Proof. The result is true for $n = 0$, as a local algebra admits no modules of projective dimension 1.

For $n > 0$, we proceed by induction on N , the case $N = 1$ being obvious. Suppose that the geometric realization of \mathcal{B}_{N-1} is an n -ball for some $N \geq 2$. Our goal is to show that the intersection $\sigma_N \cap \mathcal{B}_{N-1}$, which lies in the boundary of \mathcal{B}_{N-1} , is a union of $(n-1)$ -faces of σ_N . Then the geometric realization of \mathcal{B}_N is either an n -sphere or an n -ball, according as $\sigma_N \cap \mathcal{B}_{N-1}$ is the whole boundary of σ_N or not. The case of a sphere can be ruled out, as we know that \mathcal{B}_N has a non-empty boundary by the remark in 1.3.

The intersection $\sigma_N \cap \mathcal{B}_{N-1}$ contains at least one $(n-1)$ -face of σ_N , and hence \mathcal{B}_N is connected. Indeed, σ_N is distinct from the unique minimal n -simplex of \mathcal{C}_A , whose vertices are the indecomposable projectives (remark 1.2). Any predecessor of σ_N in K , and in particular the tail of any arrow in K whose head is σ_N , belongs to \mathcal{B}_{N-1} .

Now let $\tau = (T_0, \dots, T_r)$ be a simplex in $\sigma_N \cap \mathcal{B}_{N-1}$, and let $\bigoplus_{i=r+1}^n T_i$ be the Bongartz completion of $\bigoplus_{i=0}^r T_i$. By proposition 1.2, the n -simplex $\sigma = (T_0, \dots, T_n)$ is the unique minimal vertex of K_τ . Note that σ_N is a vertex of K_τ . As any path in K from σ to σ_N lies in K_τ by lemma 2.1, and since any predecessor of σ_N belongs to \mathcal{B}_{N-1} , there is an $(n-1)$ -simplex in $\sigma_N \cap \mathcal{B}_{N-1}$ containing τ .

Remark. If \mathcal{C}_A is infinite, the same argument shows that the geometric realization of a union $\sigma_1 \cup \dots \cup \sigma_M$ is an n -ball, provided that the full subquiver of K whose vertices are $\sigma_1, \dots, \sigma_M$ is closed under predecessors in K .

3. Examples

3.1. Let Q be the quiver $\cdot \rightrightarrows \cdot$ and A its quiver algebra. Denote by P_m and I_m the preprojective and preinjective indecomposables, respectively, given by

$$\begin{array}{ccc}
 \begin{array}{c} \begin{bmatrix} 1 & & 0 \\ & \ddots & \\ 0 & & 1 \\ 0 & \cdots & 0 \end{bmatrix} \\ P_m = k^m \longrightarrow k^{m+1} \end{array} & & \begin{array}{c} \begin{bmatrix} 1 & 0 & 0 \\ & \ddots & \vdots \\ 0 & 1 & 0 \end{bmatrix} \\ I_m = k^{m+1} \longrightarrow k_m \end{array} \\
 \begin{array}{c} \longrightarrow \\ \begin{bmatrix} 0 & 0 \\ 1 & \\ & \ddots & \\ & & 1 \end{bmatrix} \end{array} & & \begin{array}{c} \longrightarrow \\ \begin{bmatrix} 0 & 1 & 0 \\ \vdots & \ddots & \\ 0 & 0 & 1 \end{bmatrix} \end{array}
 \end{array}$$

for $m \geq 0$. These are the only indecomposables that do not extend themselves. As \mathcal{C} is infinite, our theorem does not apply. In fact, the complex \mathcal{C}_A has two connected components:

$$P_0 - P_1 - P_2 - \cdots$$

$$\cdots I_2 - I_1 - I_0.$$

The arrows of K are:


$$(P_m, P_{m+1}) \rightarrow (P_{m+1}, P_{m+2})$$

and


$$(I_{m+2}, I_{m+1}) \rightarrow (I_{m+1}, I_m),$$

for $m \geq 0$. They all correspond to almost split sequences.

3.2. Let Λ be the quiver algebra of $Q = 1 \rightarrow 2 \rightarrow 3 \leftarrow 4$, and denote by \overline{ij} a representative of the indecomposable whose support are the vertices $i, i+1, \dots, j$, for $1 \leq i \leq j \leq 4$. We only draw K as it contains all information necessary to build \mathcal{C}_Λ .




3.3. Consider the quiver $Q = \cdot \xrightarrow{\alpha} \cdot \circlearrowleft \beta$, let I be the two-sided ideal in the quiver-algebra kQ generated by β^3 , and set $\Lambda = kQ/I$. Then C_Λ is an interval:




To picture representations, we represent each basis vector by a dot. The linear map $V(\gamma) : V(i) \rightarrow V(j)$ corresponding to an arrow $\gamma : i \rightarrow j$ sends a dot in $V(i)$ to the sum of the heads of all arrows of type γ starting at the dot, and to zero if there is no such arrow.

3.4. Finally, we give an example of an algebra A of infinite representation type and for which the complex \mathcal{C}_A is finite. Let Q be the quiver



and I the two-sided ideal in kQ generated by $\alpha\beta$ and $\gamma\delta$. The complex \mathcal{C}_A for the algebra $A = kQ/I$ is the following:



REFERENCES

- [1] AUSLANDER M. and SMALØ S., *Preprojective modules over Artin algebras*. J. of Algebra, 66, 61–122 (1980).
- [2] BONGARTZ K., *Tilted algebras*. In *Representations of algebras*. Lecture Notes in Math. Springer, 903, 26–39 (1981).
- [3] BRENNER S. and BUTLER M. C. R., *Generalizations of the Bernstein-Gelfand-Ponomarev reflection-functors*. In *Representations theory II*. Lecture Notes in Math. Springer, 832, 103–170 (1979).
- [4] HAPPEL D., *Partial tilting modules and recollements*. Preprint 89-016, Bielefeld.
- [5] HAPPEL D. and RINGEL C. M., *Tilted algebras*. TAMS, 274, 399–443 (1982).
- [6] HAPPEL D. and UNGER L., *Almost complete tilting modules*. Proc. AMS, 107, 603–610 (1989).
- [7] RIEDTMANN CH. and SCHOFIELD A., *On open orbits and their complements*. J. of Algebra, 130, 388–411 (1990).

Christine Riedtmann
 Institut Fourier
 Université de Grenoble I
 BP74
 38402 St Martin D'Hères
 (France)

Aidan Schofield
 School of Mathematics
 University of Bristol
 University Walk
 Bristol BS 81 TW
 (Great Britain)

Received October 18, 1989