

On the triple points of singular maps

Autor(en): **Ekholm, Tobias / Szücs, András**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **77 (2002)**

PDF erstellt am: **27.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-57934>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

On the triple points of singular maps

Tobias Ekholm and András Szűcs

Abstract. The number of triple points (mod 2) of a self-transverse immersion of a closed $2n$ -manifold M into $3n$ -space are known to equal one of the Stiefel–Whitney numbers of M . This result is generalized to the case of generic (i.e. stable) maps with singularities. Besides triple points and Stiefel–Whitney numbers, a certain linking number of the manifold of singular values with the rest of the image is involved in the generalized equation which corrects an erroneous formula in [9].

If n is even and the closed manifold is oriented then the equations mentioned above make sense over the integers. Together, the integer- and mod 2 generalized equations imply that a certain Stiefel–Whitney number of closed oriented $4k$ -manifolds vanishes. This Stiefel–Whitney number is in fact the first in a family which vanish on such manifolds.

Mathematics Subject Classification (2000). 57R20, 57R45, 58K30.

Keywords. Stable map, linking number, triple point, Stiefel–Whitney number, orientable $4k$ -manifold.

1. Introduction

In his classical paper [10] of 1946, Whitney showed that the number of double points of a self-transverse immersion of an n -manifold into $2n$ -space is related to the Euler number of its normal bundle. Since then many results of a similar nature have been found. This paper deals with a generalization of one of these results, the Herbert–Ronga formula [5] which expresses the number of triple points of a self-transverse immersion of a closed $2n$ -manifold into $3n$ -space in terms of one of its characteristic numbers. More precisely, the Herbert–Ronga formula is extended to singular generic (i.e. stable) maps of $2n$ -manifolds into $3n$ -space. (In this paper all manifolds and maps are assumed to be C^∞ -smooth, unless otherwise explicitly stated.) To state the formula, some notation is needed:

Let M be a closed $2n$ -manifold and let $f: M \rightarrow \mathbb{R}^{3n}$ be a generic map. If $\Delta(f) \subset \mathbb{R}^{3n}$ denotes the set of double points of f then $\Delta(f)$ is an immersed n -dimensional submanifold with boundary. The self-intersection points of $\Delta(f)$ are the triple points of f . The boundary of $\Delta(f)$ is $\Sigma(f)$, the set of singular values of f .

Define $t_2(f) \in \mathbb{Z}_2$ as the mod 2-number of triple points of f . Let $\Sigma'(f)$ denote the $(n-1)$ -dimensional submanifold of \mathbb{R}^{3n} which is obtained by shifting $\Sigma(f)$ slightly along its outward normal vector field in $\Delta(f)$. Then $\Sigma'(f) \cap f(M) = \emptyset$. Define $l_2(f) \in \mathbb{Z}_2$ as the mod 2-linking number of the cycles $f(M)$ and $\Sigma'(f)$ in \mathbb{R}^{3n} . If $i_1 + \dots + i_m = 2n$ then let $\bar{w}_{i_1} \dots \bar{w}_{i_m}[M] \in \mathbb{Z}_2$ denote the product of the normal Stiefel–Whitney classes of M in dimensions i_1, \dots, i_m evaluated on the fundamental homology class of M .

Theorem 1. *Let M be a closed manifold of dimension $2n$ and let $f: M \rightarrow \mathbb{R}^{3n}$ be a generic map. Then*

$$t_2(f) + l_2(f) = \bar{w}_n^2[M] + \bar{w}_{n+1}\bar{w}_{n-1}[M] \quad (1)$$

Theorem 1 is proved in Section 2. It corrects the erroneous theorem on the second page of [9], in which the second term in the right hand side of Equation (1) is missing.

For closed oriented $4k$ -manifolds Equation (1) can be lifted to an integer equation: If $n = 2k$ is even and M is oriented then there is an induced orientation on $\Delta(f)$ as well as on the triple points of f . Define $t(f) \in \mathbb{Z}$ as the algebraic number of triple points of f . The orientation of $\Delta(f)$ induces an orientation of its boundary $\Sigma(f)$ which in turn induces an orientation of $\Sigma'(f)$. Define $l(f) \in \mathbb{Z}$ as the linking number of the oriented cycles $f(M)$ and $\Sigma'(f)$ in \mathbb{R}^{6k} . Let $\bar{p}_k[M^{4k}]$ denote the k^{th} normal Pontryagin number of M . The following theorem is Lemma 4 in [1].

Theorem 2. *Let M be a closed oriented manifold of dimension $4k$ and let $f: M \rightarrow \mathbb{R}^{6k}$ be a generic map. Then*

$$3t(f) - 3l(f) = \bar{p}_k[M]. \quad (2)$$

Equation (2) turned out to be very useful: It is used in the derivation of a geometric formula for Smale invariants of immersions of spheres, see [1] and [2], and in the study of geometric features of the regular homotopy classification of immersions of 3-manifolds in 5-space, see [7].

If M is a closed oriented $4k$ -manifold then the mod 2-reduction of $\bar{p}_k[M]$ equals $\bar{w}_{2k}^2[M]$. Hence Theorems 1 and 2 together imply that

$$\bar{w}_{2k+1}\bar{w}_{2k-1}[M] = 0 \quad (3)$$

for any closed oriented $4k$ -manifold M . In fact, $\bar{w}_{2k+1}\bar{w}_{2k-1}[M]$ is the first in a sequence of Stiefel–Whitney numbers which vanish on closed oriented $4k$ -manifolds. More precisely,

Theorem 3. (Stong). *If M is an oriented $4k$ -manifold and $(2k_1 + 1) + \dots + (2k_r + 1) = 4k$ then*

$$\bar{w}_{2k_1+1} \dots \bar{w}_{2k_r+1}[M] = 0.$$

This theorem was communicated by R. Stong to the second author together with a proof of the first case (3). A proof of Theorem 3 is presented in Section 3.

2. Proof of Theorem 1

Fix a generic map $f: M \rightarrow \mathbb{R}^{3n}$ of a closed $2n$ -manifold. Let $\tilde{\Sigma} \subset M$ denote the $(n-1)$ -dimensional submanifold of singular points of f and let $\Sigma = f(\tilde{\Sigma})$. Then f maps $\tilde{\Sigma}$ diffeomorphically to Σ .

Let $\tilde{\Delta} \subset M$ denote the closure of the preimages of multiple points of f . Then $\tilde{\Delta}$ is an immersed closed n -dimensional manifold with transverse double points at the preimages of triple points of f . Let $\tilde{\Delta}_{\text{res}}$ denote the resolution of $\tilde{\Delta}$ and let $\tilde{\iota}: \tilde{\Delta}_{\text{res}} \rightarrow M$ denote the natural immersion with image $\tilde{\Delta} \subset M$.

There is a natural involution $T: \tilde{\Delta}_{\text{res}} \rightarrow \tilde{\Delta}_{\text{res}}$ such that $f \circ \tilde{\iota} \circ T = f \circ \tilde{\iota}$. Since no triple point of f is singular we have a natural embedding $\tilde{\Sigma} \subset \tilde{\Delta}_{\text{res}}$ and $\tilde{\Sigma}$ is the fix point set of T .

Let $\nu(\tilde{\iota})$ denote the normal bundle of the immersion $\tilde{\iota}$ and let ν denote its restriction to $\tilde{\Sigma}$. Since ν is an n -dimensional vector bundle over an $(n-1)$ -manifold there exists a non-zero section. Let \tilde{s} be such a section.

A standard transversality argument allows us to extend \tilde{s} to a section \tilde{S} of $\nu(\tilde{\iota})$ which is transverse to the 0-section and which satisfies the following two conditions:

- If x is a double point of $\tilde{\iota}$ then $\tilde{S}(x) \neq 0$.
- If $\tilde{S}(x) = 0$ then $\tilde{S}(T(x)) \neq 0$.

Let $\Delta \subset \mathbb{R}^{3n}$ denote the closure of the double points of f . Then Δ is an immersed submanifold with boundary Σ and Δ has triple points at the triple points of f . Let Δ_{res} denote the resolution of Δ and let $\iota: \Delta_{\text{res}} \rightarrow \mathbb{R}^{3n}$ denote the natural immersion with image Δ . Let $\nu(\iota)$ denote the normal bundle of the immersion ι . Note that there is a natural map $\Pi: \tilde{\Delta}_{\text{res}} \rightarrow \Delta_{\text{res}}$ which is a double cover of $\Delta_{\text{res}} - \Sigma$ when restricted to $\tilde{\Delta}_{\text{res}} - \tilde{\Sigma}$, and which maps $\tilde{\Sigma}$ diffeomorphically onto Σ .

Define the section S of $\nu(\iota)$ as follows:

$$S(y) = \begin{cases} df(\tilde{S}(y_1)) + df(\tilde{S}(y_2)) & \text{if } y \in \Delta_{\text{res}} - \Sigma, \text{ where } y_1 \neq y_2, \Pi(y_1) = \Pi(y_2) = y, \\ 2df(\tilde{S}(y_1)) & \text{if } y \in \Sigma, \text{ where } \Pi(y_1) = y. \end{cases}$$

Let $C(\Sigma) \subset \Delta_{\text{res}}$ be a small open collar on the boundary Σ of Δ_{res} . Let Δ'' denote the image of the immersion $y \mapsto \iota(y) + \epsilon S(y)$, $y \in \Delta_{\text{res}} - C(\Sigma)$ for some small $\epsilon > 0$. Then, if ϵ and the collar $C(\Sigma)$ are small enough, Δ'' is a chain with boundary $\partial\Delta'' = \Sigma''$ satisfying $\Sigma'' \cap f(M) = \emptyset$. If lk_2 denotes the mod 2-linking number, \bullet denotes the mod 2-intersection number, and $\sharp(F)$ denotes the mod 2-number of elements in the finite set F , then

$$\text{lk}_2(\Sigma'', f(M)) = \Delta'' \bullet f(M) = \sharp(\tilde{S}^{-1}(0)) + t_2(f), \quad (4)$$




Figure 1. A piece of $f(M)$ (represented by a 2-sphere and a piece of a plane) with the double point set Δ (fat lines), its normal field S , and singularity set Σ (dots) with its outward normal field V in Δ .

since near each zero z of \tilde{S} there is a unique intersection point of Δ'' and $f(M)$ near $f(z)$, and near each triple point of f there are exactly three such intersection points.

The homology class of the cycle $\tilde{\Delta}$ in M is Poincaré dual to n^{th} normal Stiefel–Whitney class \bar{w}_n of M , see [6]. Thus

$$\bar{w}_n^2[M] = \tilde{\Delta} \bullet \tilde{\Delta} = \sharp(\tilde{S}^{-1}(0)), \quad (5)$$

since the image of a slight shift of the immersion $\tilde{\iota}$ along \tilde{S} intersects $\tilde{\Delta}$ near each zero of \tilde{S} and in *two* points near each double point of $\tilde{\iota}$.

Equations (4) and (5) imply

$$\text{lk}_2(\Sigma'', f(M)) = \bar{w}_n^2[M] + t_2(f). \quad (6)$$

Recall that $\Sigma' \subset \mathbb{R}^{3n}$ is the submanifold which results when Σ is shifted slightly along its unit outward normal vector field V in Δ , and that $\Sigma' \cap f(M) = \emptyset$. We compare the linking numbers $\text{lk}_2(\Sigma'', f(M))$ and $\text{lk}_2(\Sigma', f(M))$:

Let $\tilde{\Sigma}_0 \subset M$ be the submanifold which results when $\tilde{\Sigma}$ is shifted a small distance along \tilde{S} . Let $\Sigma_0 = f(\tilde{\Sigma}_0)$ and for $p \in \Sigma$, let $p_0 = f(\tilde{p}_0)$ where \tilde{p}_0 is the point in $\tilde{\Sigma}_0$ corresponding to $\tilde{p} \in \tilde{\Sigma}$ with $f(\tilde{p}) = p$.

For small $\epsilon > 0$ and $p \in \Sigma$ let $l_p(\epsilon)$ be the segment of the straight line through $p + \epsilon V(p)$ and p_0 of length 2ϵ and centered at p_0 . For $\epsilon > 0$ and the shifting of $\tilde{\Sigma}$ in M small enough,

$$\Gamma = \bigcup_{p \in \Sigma} l_p(\epsilon)$$

is a submanifold of \mathbb{R}^{3n} . If the collar $C(\Sigma)$ is chosen small enough and if the

the fundamental homology class of the manifold V and PD denotes the Poincaré duality operator,

$$\begin{aligned} \langle w_{n-1}(\xi), F_{\Sigma_0} \rangle &= \langle i_0^* \bar{w}(M), F_{\Sigma_0} \rangle = \langle \bar{w}(M), i_{0*}(F_{\Sigma_0}) \rangle = \langle \bar{w}(M), \text{PD } \bar{w}_{n+1}(M) \rangle \\ &= \langle \bar{w}(M) \cup \bar{w}_{n+1}(M), F_M \rangle = \bar{w}_{n-1} \bar{w}_{n+1}[M]. \end{aligned} \quad (10)$$

Here, the third equality follows from the well-known formula $\text{PD } \bar{w}_{n+1}(M) = i_* F_{\tilde{\Sigma}}$, where $i: \tilde{\Sigma} \rightarrow M$ denotes the inclusion, together with $i_* F_{\tilde{\Sigma}} = i_{0*} F_{\Sigma_0}$. Equations (6), (7), and (10) prove the theorem. \square

3. Proof of Theorem 3

Let \mathfrak{N}_* , Ω_* , and Ω_*^U denote the cobordism ring, the oriented cobordism ring, and the complex cobordism ring, respectively. Note that there are natural forgetting homomorphisms

$$\Omega_*^U \longrightarrow \Omega_* \longrightarrow \mathfrak{N}_*.$$

For a manifold M , let $[M]$ denote its cobordism class.

Using some facts from cobordism theory which can all be found in Chapter 4 of Stong's book [8], we show that it is enough to prove the theorem for oriented $4k$ -manifolds M such that either

- (a) $[M] \in \Omega_{4k}$ maps to a square $[N \times N] \in \mathfrak{N}_{4k}$, or
- (b) $[M]$ is a torsion element of Ω_{4k} (in fact, $[M]$ torsion implies $2 \cdot [M] = 0$):

Let $\text{Tors}(\Omega_*)$ denote the torsion subgroup of Ω_* . The homomorphism $\Omega_*^U \rightarrow \Omega_*$ induces an epimorphism

$$\Omega_*^U \longrightarrow \Omega_* / \text{Tors}(\Omega_*).$$

and the image $\Omega_*^U \rightarrow \mathfrak{N}_*$ consists of squares of elements in \mathfrak{N}_* .

Hence, if M is any oriented $4k$ -manifold then there exists some oriented $4k$ -manifold V such that $[V]$ is torsion in Ω_{4k} and $[M] + [V] = [N \times N]$ in \mathfrak{N}_{4k} . This implies that the theorem follows once it is proved for manifolds satisfying (a) or (b) above.

First consider (a): let $M = N \times N$. Then $\bar{w}(M) = \bar{w}(N) \times \bar{w}(N)$ and hence

$$\bar{w}_{2k+1}(M) = \sum_{i+j=2k+1} \bar{w}_i(N) \times \bar{w}_j(N).$$

Thus

$$\begin{aligned} \langle \bar{w}_{2k_1+1}(M) \dots \bar{w}_{2k_r+1}(M), F_M \rangle &= \\ &= \sum \langle \bar{w}_{i_1}(N) \dots \bar{w}_{i_r}(N), F_N \rangle \cdot \langle \bar{w}_{j_1}(N) \dots \bar{w}_{j_r}(N), F_N \rangle. \end{aligned} \quad (11)$$

Since $i_s + j_s$ is odd for all i_s, j_s there is a fixed point free involution T acting on

the set of the terms in the sum in (11) such that $i_1 + \dots + i_r = 2k = j_1 + \dots + j_r$:

$$\begin{aligned} T: & \langle \bar{w}_{i_1}(N) \dots \bar{w}_{i_r}(N), F_N \rangle \cdot \langle \bar{w}_{j_1}(N) \dots \bar{w}_{j_r}(N), F_N \rangle \\ & \mapsto \langle \bar{w}_{j_1}(N) \dots \bar{w}_{j_r}(N), F_N \rangle \cdot \langle \bar{w}_{i_1}(N) \dots \bar{w}_{i_r}(N), F_N \rangle. \end{aligned}$$

Thus the terms in the left hand side of (11) which does not vanish for dimensional reasons cancel in pairs and hence $\bar{w}_{2k_1+1} \dots \bar{w}_{2k_r+1}[M] = 0$.

Next consider (b): let $u: \mathfrak{N}_{4k} \rightarrow \mathbb{Z}_2$ denote the homomorphism induced by the product of odd-dimensional normal Stiefel–Whitney classes $\bar{w}_{2k_1+1} \dots \bar{w}_{2k_r+1}$, $\sum 2k_j + 1 = 4k$. Odd-dimensional Stiefel–Whitney classes are mod 2-reductions of twisted integer classes, see [3], p. 140. Hence, a product of an even number of such classes is an integer class so the map

$$\Omega_{4k} \xrightarrow{\pi} \mathfrak{N}_{4k} \xrightarrow{u} \mathbb{Z}_2$$

lifts to a homomorphism

$$\Omega_{4k} \xrightarrow{U} \mathbb{Z}.$$

Thus U and therefore $u \circ \pi$ is zero on any torsion element of Ω_{4k} . \square

References

- [1] T. Ekholm and A. Szűcs, Geometric formulas for Smale invariants of codimension two immersions, *Topology*, to appear.
- [2] T. Ekholm and A. Szűcs, The group of immersions of homotopy $4k - 1$ -spheres, preprint.
- [3] J. Milnor and J. Stasheff, *Characteristic classes*, Ann. of Math. Stud. **76**, Princeton Univ. Press, 1974.
- [4] D. Quillen, Elementary proof of some results of cobordism theory using Steenrod operations, *Adv. in Math.* **7** (1971), 29–56.
- [5] F. Ronga, On multiple points of smooth immersions *Comment. Math. Helv.* **55** (1980), 521–527.
- [6] F. Ronga, La classe duale aux points doubles d’une application *Composito Math.* **27** (1973), 223–232.
- [7] O. Saeki, A. Szűcs and M. Takase, Regular homotopy classes of immersions of 3-manifolds into 5-space, *Manuscripta Math.*, to appear.
- [8] R. Stong, *Notes on cobordism theory*, Princeton Univ. Press, 1968.
- [9] A. Szűcs, The linking number of singular maps, *Comment. Math. Helv.* **61** (1986), 360–369.
- [10] H. Whitney, The self-intersections of a smooth n -manifold in $2n$ -space, *Ann. of Math.* **45** (1944), 220–293.

Tobias Ekholm
Uppsala University
Department of Mathematics
P.O. Box 480
SE-751 06 Uppsala
Sweden
e-mail: tobias@math.uu.se

A. Szűcs
Eötvös Loránd University
Department of Analysis
Pázmány Péter sétány 1/C
H-1117 Budapest
Hungary
e-mail: szucs@cs.elte.hu

(Received: October 12, 2001)