

Homology torsion growth and Mahler measure

Autor(en): **Le, Thang**

Objektyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **89 (2014)**

PDF erstellt am: **27.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-515684>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Homology torsion growth and Mahler measure

Thang Le*

Abstract. We prove a conjecture of K. Schmidt in algebraic dynamical system theory on the growth of the number of components of fixed point sets. We also generalize a result of Silver and Williams on the growth of homology torsions of finite abelian covering of link complements. In both cases, the growth is expressed by the Mahler measure of the first non-zero Alexander polynomial of the corresponding modules. We use the notion of pseudo-isomorphism, and also tools from commutative algebra and algebraic geometry, to reduce the conjectures to the case of torsion modules. We also describe concrete sequences which give the expected values of the limits in both cases. For this part we utilize a result of Bombieri and Zannier (conjectured before by A. Schinzel) and a result of Lawton (conjectured before by D. Boyd).

Mathematics Subject Classification (2010). 57M10, 57M25, 57Q10, 37B50, 37B10.

Keywords. Torsion growth, Mahler measure, Alexander polynomials, algebraic dynamical system, entropy, pseudo-isomorphism.

Introduction

A conjecture of K. Schmidt. Suppose M is a finitely generated module over the commutative ring $\mathfrak{R} := \mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$. Let \mathbb{S} be the unit circle in the complex plane \mathbb{C} . There is a natural action of \mathbb{Z}^n on the compact abelian group $\hat{M} = \text{Hom}(M, \mathbb{S})$, the Pontryagin dual of M . For details on dynamical systems of this type the reader is referred to the remarkable book [Sch]. The entropy of this action, denoted by $h(M)$, can be defined in a standard manner. Lind, Schmidt, and Ward [LSW] (see also [EW]) proved that if M is a *torsion module*, then

$$h(M) = \mathbb{M}(\Delta_0(M)), \quad (1)$$

where $\Delta_0(M)$ is the 0-th Alexander polynomial of M (also known as the order of M), and $\mathbb{M}(f)$ is the additive Mahler measure of the polynomial f . We will recall the definitions of these notions in Section 1.

For a subgroup $\Gamma \subset \mathbb{Z}^n$ of finite index let $\text{Fix}_\Gamma(\hat{M})$ be the set of elements of \hat{M} fixed by actions of elements of Γ . Then $\text{Fix}_\Gamma(\hat{M})$ is a compact subgroup of \hat{M} and

*The author was supported in part by National Science Foundation.

has a finite number $P_\Gamma(\hat{M})$ of connected components. The following theorem was conjectured by K. Schmidt [Sch], based on results in the torsion module case.

Theorem 1. *For any finitely generated \mathfrak{R} -module M one has*

$$\limsup_{\langle \Gamma \rangle \rightarrow \infty} \frac{\log P_\Gamma(\hat{M})}{|\mathbb{Z}^n / \Gamma|} = h(\text{tor}(M)).$$

If $n = 1$ then one can replace the \limsup by the ordinary \lim .

Here $\text{tor}(M)$ is the torsion submodule of M , and

$$\langle \Gamma \rangle = \min\{|x|, x \in \Gamma \setminus \{0\}\},$$

where $|x| = \sqrt{\sum_i |x_i|^2}$ for $x = (x_1, \dots, x_n) \in \mathbb{Z}^n$.

The theorem had been proved for the case when M is a torsion module by Schmidt, see [Sch], Theorem 21.1, and we will make substantial use of this case.

A conjecture of Silver and Williams. Suppose L is an oriented link with n ordered components in an oriented integral homology 3-sphere Z , with the complement $X = Z \setminus L$. There is a natural identification $H_1(X, \mathbb{Z}) = \mathbb{Z}^n$. For a subgroup $\Gamma \subset \mathbb{Z}^n$ of finite index let X_Γ be the corresponding abelian covering of X , and X_Γ^{br} the corresponding branched covering of Z . There are defined the Alexander polynomials $\Delta_i(L) \in \mathfrak{R} = \mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$, $i = 0, 1, 2, \dots$. We will recall the definition of $\Delta_i(L)$ in Section 4.

Let $\Delta(L) = \Delta_j(L)$, where j is the smallest index such that $\Delta_j(L) \neq 0$. For an abelian group G , denote $\text{tor}_{\mathbb{Z}}(G)$ the \mathbb{Z} -torsion subgroup of G .

Theorem 2. *Notations as above. One has*

$$\limsup_{\langle \Gamma \rangle \rightarrow \infty} \frac{\log |\text{tor}_{\mathbb{Z}}(H_1(X_\Gamma, \mathbb{Z}))|}{|\mathbb{Z}^n / \Gamma|} = \mathbb{M}(\Delta(L)), \quad (\text{a})$$

$$\limsup_{\langle \Gamma \rangle \rightarrow \infty} \frac{\log |\text{tor}_{\mathbb{Z}}(H_1(X_\Gamma^{\text{br}}, \mathbb{Z}))|}{|\mathbb{Z}^n / \Gamma|} = \mathbb{M}(\Delta(L)). \quad (\text{b})$$

If $n = 1$ then one can replace the \limsup by the ordinary \lim .

For the special case when $\Delta(L) = \Delta_0(L)$, part (b) was proved by Silver and Williams [SW], who, based on that result, formulated part (b), with the upper limit replaced by the ordinary limit, as a conjecture. The proof in [SW] (for the case $\Delta(L) = \Delta_0(L)$), written for $Z = S^3$ and for branched covering only, can be modified for the case of general homology 3-spheres and non-branched coverings. Hence the real new content of Theorem 2 is the case when $\Delta(L) \neq \Delta_0(L)$. The proof

in [SW] is based on the torsion module case of Theorem 1. It is not surprising that if one can get Theorem 1, then one can generalize the result of Silver and Williams to the case $\Delta(L) \neq \Delta_0(L)$.

The investigation of the growth of homology torsions of finite covering of knots has a long history, with an interesting conjecture posed by Gordon [Go]. The conjecture was proved by Riley [Ri] and Gonzalez-Acuna and Short [GS] using Gelfond–Baker results in number theory. Silver and Williams’ result mentioned above and Theorem 2 are generalizations of Riley and Gonzalez-Acuna and H. Short from the knot to the link case. The surprising appearance of the Mahler measure can be explained from the perspective of L^2 -torsion theory [Lü2]: The L^2 -torsion of the maximal abelian covering of a link complement, at least when $\Delta_0(L) \neq 0$, is the Mahler measure of $\Delta_0(L)$. Theorem 2 more or less says that the L^2 -torsion in this case can be approximated by its finite-dimensional counterparts.

In the non-abelian covering case the L^2 -torsion is given by the hyperbolic volume. So one should expect the similar limit of the left-hand side of Equation (a) would give the hyperbolic volume of the link complements. In [Le2], we will show that, if L is a non-split link in S^3 , then

$$\limsup_{\langle \Gamma \rangle \rightarrow \infty} \frac{\log |\operatorname{tor}_{\mathbb{Z}}(H_1(X_{\Gamma}, \mathbb{Z}))|}{|\mathbb{Z}^n / \Gamma|} \leq \frac{\operatorname{vol}(L)}{6\pi}. \quad (2)$$

where $\operatorname{vol}(L)$ is the sum of the hyperbolic volumes of the hyperbolic pieces in the Jaco–Shalen–Johansson decomposition of $S^3 \setminus L$. Here Γ runs the set of all subgroups of $\pi_1(L)$ of finite index, and $\langle \Gamma \rangle$ is the minimal word length of $\Gamma \setminus \{1\}$, measured using a fixed finite generator set of π_1 . In particular, if $\operatorname{vol}(L) = 0$, we have the equality in (2). For example, if L is a torus knot, then one has equality in (2). For works in this direction see also [Le1], [Mü], [BV]. It is expected that the non-abelian case is much more complicated than the abelian case.

An algebraic version of Theorem 1. It is not difficult to reformulate Theorem 1 entirely in terms of module M , without going through the Pontryagin dual \hat{M} . We will show that Theorem 1 is equivalent to the following.

Theorem 3. *For any finitely generated \mathfrak{R} -module M one has*

$$\limsup_{\langle \Gamma \rangle \rightarrow \infty} \frac{\log |\operatorname{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[\mathbb{Z}^n / \Gamma])|}{|\mathbb{Z}^n / \Gamma|} = \mathbb{M}(\Delta(\operatorname{tor} M)).$$

If $n = 1$ then one can replace the \limsup by the ordinary \lim .

Theorem 3 is a special case of the following.

Theorem 4. Suppose \mathcal{C} is a chain complex of finitely generated free \mathfrak{R} -modules. Then for every $i \geq 0$,

$$\limsup_{\langle \Gamma \rangle \rightarrow \infty} \frac{\log |\operatorname{tor}_{\mathbb{Z}}(H_i(\mathcal{C} \otimes \mathbb{Z}[\mathbb{Z}^n/\Gamma]))|}{|\mathbb{Z}^n/\Gamma|} = \mathbb{M}(\Delta(H_i(\mathcal{C}))).$$

If $n = 1$ then one can replace the \limsup by the ordinary \lim .

In this paper we will prove Theorem 3, and from there deduce Theorem 2 and 4.

Application: abelian covering of CW-complex. Suppose X is a finite CW-complex, equipped with a surjection $\rho: H_1(X, \mathbb{Z}) \rightarrow \mathbb{Z}^n$. Let \tilde{X} be the abelian covering of X corresponding ρ . The CW-structure of X lifts to a CW-structure of \tilde{X} . The group \mathbb{Z}^n acts as deck transformations on the covering \tilde{X} , making the cellular complex $\mathcal{C}(\tilde{X})$ of \tilde{X} a free finitely generated \mathfrak{R} -complex.

For every subgroup $\Gamma \subset \mathbb{Z}^n$ of finite index let $\rho_{\Gamma}: \pi_1(X) \rightarrow A_{\Gamma} = \mathbb{Z}^n/\Gamma$ be the composition $\pi_1(X) \rightarrow H_1(X, \mathbb{Z}) \xrightarrow{\rho} \mathbb{Z}^n \rightarrow A_{\Gamma}$, where the first map is the abelianization map. Let X_{Γ} be the finite regular covering corresponding to ρ_{Γ} . Apply Theorem 4 to the complex $\mathcal{C}(\tilde{X})$ we get the following.

Theorem 5. Notations as above. Then

$$\limsup_{\langle \Gamma \rangle \rightarrow \infty} \frac{|\operatorname{tor}_{\mathbb{Z}}(H_i(X_{\Gamma}, \mathbb{Z}))|}{|\mathbb{Z}^n/\Gamma|} = \mathbb{M}(\Delta(\operatorname{tor}_{\mathfrak{R}} H_i(\tilde{X}, \mathbb{Z}))).$$

If $n = 1$ then \limsup can be replaced by the ordinary \lim .

Ideas of proofs. To prove Theorem 3 (and the equivalent Theorem 1) we will reduce it to the case when M is a torsion module, which had been proved in [Sch], and the case when M is torsion-free, i.e. when $\operatorname{tor}(M) = 0$. Although the fact that M is isomorphic to $\operatorname{tor}(M) \oplus (M/\operatorname{tor}(M))$ is not true in general, it would hold true if we replace isomorphism by *pseudo-isomorphism*, a notion introduced by Bourbaki [Bou]. The notion of pseudo-isomorphism is important for us, and we will review it in Section 1. The following will be one of the main technical results used in the proof of Theorem 3.

Theorem 6. Suppose M_1 and M_2 are two pseudo-isomorphic finitely generated \mathfrak{R} -modules. Then $|\operatorname{tor}_{\mathbb{Z}}(M_1 \otimes \mathbb{Z}[\mathbb{Z}^n/\Gamma])|$ and $|\operatorname{tor}_{\mathbb{Z}}(M_2 \otimes \mathbb{Z}[\mathbb{Z}^n/\Gamma])|$ have the same growth rate in the sense that

$$\lim_{\langle \Gamma \rangle \rightarrow \infty} \left(\frac{\log |\operatorname{tor}_{\mathbb{Z}}(M_1 \otimes \mathbb{Z}[\mathbb{Z}^n/\Gamma])|}{|\mathbb{Z}^n/\Gamma|} - \frac{\log |\operatorname{tor}_{\mathbb{Z}}(M_2 \otimes \mathbb{Z}[\mathbb{Z}^n/\Gamma])|}{|\mathbb{Z}^n/\Gamma|} \right) = 0.$$

Note that the limit in Theorem 6 is the ordinary limit, not the upper limit.

In general, the direct calculation of $|\mathrm{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[\mathbb{Z}^n/\Gamma])|$ (resp. the exact value of $|\mathrm{tor}_{\mathbb{Z}}(H_1(X_{\Gamma}^{\mathrm{br}}, \mathbb{Z}))|$) is very difficult, especially in the case when M is not a torsion module (resp. the 0-th Alexander polynomial is 0). The only known formula for $\mathrm{tor}_{\mathbb{Z}}(X_{\Gamma}^{\mathrm{br}}, \mathbb{Z})$, due to Mayberry and Murasugi [MM], applies only to the case when $H_1((X_{\Gamma}^{\mathrm{br}}, \mathbb{Z}))$ itself is a torsion \mathbb{Z} -group. When the 0-th Alexander polynomial is 0, which is the case concerned in this paper, the only known result is that of Hillman and Sakuma [HS] who calculated part of the torsion $\mathrm{tor}_{\mathbb{Z}}(H_1(X_{\Gamma}^{\mathrm{br}}, \mathbb{Z}))$. The other not-yet-calculated part is related to the more difficult theory of modular representations of finite groups.

To circumvent this problem, we use an approximation $\beta(\Gamma)$ of $\mathbb{Z}[\mathbb{Z}^n/\Gamma]$, for which the calculation of $|\mathrm{tor}_{\mathbb{Z}}(M_1 \otimes \beta)|$ is easier. Here $\beta(\Gamma)$ depends on Γ and other data, and it approximates $\mathbb{Z}[\mathbb{Z}^n/\Gamma]$ in the sense that $|\mathrm{tor}_{\mathbb{Z}}(M_1 \otimes \mathbb{Z}[\mathbb{Z}^n/\Gamma])|$ and $|\mathrm{tor}_{\mathbb{Z}}(M_1 \otimes \beta(\Gamma))|$ have the same growth rate. The construction of $\beta(\Gamma)$ is based on the theory of torsion points on algebraic varieties. Needless to say, we have to use tools in commutative and homological algebra to get the desired estimates.

Sequence of converging subgroups. Theorem 3 guarantees there is a sequence of subgroups $\Gamma_s \subset \mathbb{Z}^n$ of finite index such that

$$\lim_{s \rightarrow \infty} \frac{\log |\mathrm{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[\mathbb{Z}^n/\Gamma_s])|}{|\mathbb{Z}^n/\Gamma_s|} = \mathbb{M}(\Delta(\mathrm{tor} M)).$$

In the case when M is a torsion module, half of the proof of Theorem 1 in [Sch] is to construct such a sequence. The construction is long and difficult. In Section 5 (see Theorem 7) we give new sequences Γ_s that work for both torsion and non-torsion modules. The proof is probably simpler, because we are able to use a result of Bombieri and Zannier [SZ], [BMZ] on irreducibility of lacunary polynomials which was conjectured before by Schinzel, and a result of Lawton on approximation of Mahler measure which was a conjecture of Boyd. The methods and results of Section 5 are independent of the other parts and give an independent proof of “half” of Theorem 3 (or Theorem 1), namely that the left-hand side of the identity of Theorem 3 is greater than or equal to the right-hand side.

While writing this paper I was informed by Raimbault [Ra] that he gets an independent result similar to Theorem 7 of Section 5 by modifying the sequences in [Sch].

Structure of the paper. Section 1 contains notations, basic facts (with some enhancements) about torsion points on algebraic varieties, pseudo-isomorphism, order of modules, lattices in Hermitian spaces, the integral group ring of finite abelian groups. It also contains a proof that Theorems 1 and 3 are equivalent. In Section 2, the main technical section, we present the construction the approximation β of $\mathbb{Z}[\mathbb{Z}^n/\Gamma]$. Section 3 contains proofs of Theorems 6, 3, and 4. Section 4 gives a proof

of Theorem 2. The last section contains the construction of converging sequences of lattices and Theorem 7.

Acknowledgements. I would like to thank M. Baker, Hailong Dao, J. Hillman, K. Murasugi, S. Sakuma, K. Schmidt, D. Silver, and S. Williams for helpful conversations. I am particularly indebted to U. Zannier who has patiently explained to me over the course of two years many things from diophantine approximations to his (joint with Bombieri and Masser) remarkable work on anomalous subvariety structures, including a proof of a conjecture of Schinzel.

I lectured on parts of this work at conferences in Fukuoka (March 2009), Trieste (May 2009), Columbia University (June 2009), Osaka City University (November 2009), and would like to thank the organizers for the chance to give talks there.

The paper grew out of my attempt to prove a topological volume conjecture [Le1], [Le2]. This was part of a program aiming at understanding the question “Under what conditions L^2 -torsions can be approximated by finite group counter parts?”. I was attracted to this program while trying to develop an approach to attack the volume conjecture in quantum topology, and by the beautiful work of Lück work [Lü1] on approximation of L^2 -Betti number.

1. Notations and preliminaries

1.1. Modules over $\mathfrak{R} = \mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$. Fix a free abelian group \mathbb{Z}^n . Let $\mathfrak{R} = \mathbb{Z}[\mathbb{Z}^n]$, which we identify with $\mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ by sending $\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{Z}^n$ to $t^{\mathbf{k}} = \prod_{i=1}^n t_i^{k_i}$. The ring \mathfrak{R} is a unique factorization Noetherian domain. In this paper \mathfrak{R} -modules are supposed to be finitely generated, and tensor products are assumed over \mathfrak{R} unless otherwise indicated.

For a module M over a commutative domain R , the torsion submodule $\text{tor}_R(M)$ is defined by

$$\text{tor}_R(M) = \{x \in M \mid ax = 0 \text{ for some } 0 \neq a \in R\}.$$

An R -module M is a *torsion module* if $M = \text{tor}_R M$. If $\text{tor}_R M = 0$, we call M *torsion-free*. If $R = \mathfrak{R}$ we usually drop the subscript \mathfrak{R} in the tor notation.

For a subgroup $\Gamma \subset \mathbb{Z}^n$ let $A_\Gamma := \mathbb{Z}^n / \Gamma$ and $I(\Gamma)$ the ideal of $\mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ generated by $\{1 - t_1^{k_1} \dots t_n^{k_n}, (k_1, \dots, k_n) \in \Gamma\}$. Then we have the following exact sequence

$$0 \rightarrow I(\Gamma) \rightarrow \mathfrak{R} \xrightarrow{\text{pr}} \mathbb{Z}[A_\Gamma] \rightarrow 0.$$

Hence for every \mathfrak{R} -module M ,

$$M/I(\Gamma)M \cong M \otimes_{\mathfrak{R}} \mathbb{Z}[A_\Gamma].$$

Suppose f, g are functions with positive real values on the set of subgroups $\Gamma \subset \mathbb{Z}^n$ of finite index. We say $f(\Gamma)$ has *negligible growth rate* if

$$\lim_{|\Gamma| \rightarrow \infty} f(\Gamma)^{1/|\mathbb{Z}^n/\Gamma|} = 1.$$

We say f and g have *the same growth rate*, and write $f \sim g$ if f/g has negligible growth rate. Note that we do not require the individual limit $\lim_{|\Gamma| \rightarrow \infty} f(\Gamma)^{1/|\mathbb{Z}^n/\Gamma|}$ exists in this case.

We say two \mathfrak{R} -modules M_1 and M_2 have *the same torsion growth*, and write $M_1 \sim M_2$, if

$$|\mathrm{tor}_{\mathbb{Z}}(M_1 \otimes \mathbb{Z}[A_\Gamma])| \sim |\mathrm{tor}_{\mathbb{Z}}(M_2 \otimes \mathbb{Z}[A_\Gamma])|.$$

1.2. Alexander polynomials. All definitions and facts here are standard and can be found in [Tu], [Hi].

Every finitely generated \mathfrak{R} -module M has a presentation by an exact sequence

$$\mathfrak{R}^{m_1} \xrightarrow{\partial_1} \mathfrak{R}^{m_0} \rightarrow M \rightarrow 0,$$

where ∂_1 , given by a matrix of size $m_1 \times m_0$ with entries in \mathfrak{R} , is known as a *presentation matrix* of M . A k -minor of ∂_1 is the determinant of any sub-matrix of size $k \times k$ of ∂_1 . For $j \geq 0$, the j -th *Alexander polynomial* $\Delta_j(M)$ is the greatest common divisor of all the $(m_0 - j)$ -minor of ∂_1 .

It is known that $\Delta_j(M)$ depends only on M , but not on any particular presentation matrix. Each $\Delta_j(M)$ is defined up to units in \mathfrak{R} , so identity involving $\Delta_j(M)$ should be understood “up to units”.

The 0-th polynomial $\Delta_0(M)$ is known as the *order* of M , which is non-zero if and only M is a torsion module. Besides, $\Delta_j(M)$ divides $\Delta_{j-1}(M)$ for every $j \geq 1$.

The *rank* of a module M over \mathfrak{R} is the dimension of the vector space $M \otimes F(\mathfrak{R})$ over the fractional field $F(\mathfrak{R})$ of \mathfrak{R} . If M has rank r , then $\Delta_j(M) = 0$ if $j < r$, and

$$\Delta_{j-r}(M) = \Delta_j(\mathrm{tor} M).$$

For any finitely generated \mathfrak{R} -module of rank r , define

$$\Delta(M) := \Delta_r(M) = \Delta_0(\mathrm{tor}(M)).$$

If $M = \mathfrak{R}/I$, where $I = (f_1, \dots, f_l)$ is the ideal generated by f_1, \dots, f_l , then $\Delta_0(M) = \mathrm{gcd}(f_1, \dots, f_l)$, the greatest common divisor of the elements f_1, \dots, f_l .

1.3. Pseudo-isomorphism. Reference for this part is [Bou], [Hi].

An \mathfrak{R} -module N is *pseudo-zero* if for every prime ideal \mathcal{P} of height 1, the localization $N_{\mathcal{P}}$ is 0. It is known that submodules and quotient modules of a pseudo-zero module are pseudo-zero.

In \mathfrak{R} , a prime ideal is of height 1 if and only if it is principal and generated by an irreducible polynomial.

An \mathfrak{R} -morphism $M_1 \rightarrow M_2$ is a *pseudo-isomorphism* if the kernel and co-kernel are pseudo-zero.

Two finitely generated torsion \mathfrak{R} -modules M_1, M_2 are pseudo-isomorphic if and only if $\Delta_j(M_1) = \Delta_j(M_2)$ for every $j \geq 0$; in particular, a finitely generated torsion \mathfrak{R} -module is pseudo-zero if and only if $\Delta_0(M) = 1$, see [Hi], Theorem 3.5.

Let us formulate some well-known facts in the form that will be useful for us.

Lemma 1.1. *Suppose $I \subset \mathfrak{R}$ is a prime ideal, $I \neq \mathfrak{R}$.*

- a) \mathfrak{R}/I is pseudo-zero if and only if I is not principal.
- b) If \mathfrak{R}/I is pseudo-zero and $0 \neq p \in I$, then there exists $q \in I$ such that $\gcd(p, q) = 1$.

Proof. a) Since $I \neq \mathfrak{R}$ and I is prime, $I = (p_1, \dots, p_l)$, where the p_i 's are irreducible, non-unit, and distinct. One has

$$\begin{aligned} \mathfrak{R}/I \text{ is pseudo-zero} &\iff \Delta_0(I) = \gcd(p_1, \dots, p_l) \text{ is } 1 \\ &\iff l \geq 2 \\ &\iff I \text{ is not principal.} \end{aligned}$$

b) Suppose q_1, \dots, q_l are all irreducible factors of p . Suppose the contrary that every $q \in I$ is not co-prime with p , i.e. every $q \in I$ is divisible by one of the q_i 's. Then $I \subset \bigcup_{i=1}^l (q_i)$. Since each ideal (q_i) is prime, there is an index i such that $I \subset (q_i)$. Because (q_i) has height 1 and I is prime, this means $I = (p_i)$, which is principal. This contradicts the fact that \mathfrak{R}/I is pseudo-zero. \square

The following is the main fact about pseudo-isomorphism which we will use.

Theorem 1.2 ([Bou], Theorem VII.4.5). *Any finitely generated module \mathfrak{R} -module M is pseudo-isomorphic to $\text{tor}(M) \oplus M/\text{tor}(M)$.*

Remark 1.1. It follows from (1) that if M_1, M_2 are pseudo-isomorphic, then they have the same entropy, $h(M_1) = h(M_2)$. In particular, if M is pseudo-zero, then $h(M) = 0$.

1.4. Equivalence of Theorem 1 and Theorem 3. Recall that \mathbb{S} is the unit circle in \mathbb{C} . With the usual multiplication \mathbb{S} is an abelian Lie group. For an abelian group G , the Pontryagin dual $\hat{G} = \text{Hom}(G, \mathbb{S})$ is a compact group. If $G \cong \mathbb{Z}^k$, then $\hat{G} \cong \mathbb{S}^k$. On the other hand, if $|G| < \infty$, then $G \cong \hat{\hat{G}}$. If $G \cong \mathbb{Z}^k \oplus \text{tor}_{\mathbb{Z}}(G)$, then $\hat{G} \cong \mathbb{S}^k \times \widehat{\text{tor}_{\mathbb{Z}}(G)}$. In particular, the cardinality $|\text{tor}_{\mathbb{Z}} G|$ of the \mathbb{Z} -torsion G is the number of connected components of the compact group \hat{G} .

Suppose M is a finitely generated \mathfrak{R} -module, and $\Gamma \subset \mathbb{Z}^n$ a subgroup of finite index. By definition

$$\begin{aligned} \text{Fix}_\Gamma(\widehat{M}) &= \{x \in \widehat{M} \mid \gamma \cdot x = x \text{ for all } \gamma \in \Gamma\} \\ &= \{x \in \widehat{M} = \text{Hom}(M, \mathbb{S}) \mid x(y) = x(\gamma(y)) \text{ for all } \gamma \in \Gamma, y \in M\} \\ &= \{x \in \text{Hom}(M, \mathbb{S}) \mid x((1 - \gamma)(y)) = 1 \text{ for all } \gamma \in \Gamma, y \in M\} \\ &= \{x \in \text{Hom}(M, \mathbb{S}) \mid x(I(\Gamma)M) = 1\}. \end{aligned}$$

It follows that

$$\text{Fix}_\Gamma(\widehat{M}) \cong (M/I(\Gamma)M)^\wedge \cong (M \otimes \mathbb{Z}[A_\Gamma])^\wedge.$$

We can conclude that $P_\Gamma(\widehat{M})$, the number of connected components of $\text{Fix}_\Gamma(\widehat{M})$, is

$$P_\Gamma(\widehat{M}) = |\text{tor}_\mathbb{Z}(M \otimes \mathbb{Z}[A_\Gamma])|. \quad (3)$$

From (1) and (3) we see that Theorem 1 and Theorem 3 are equivalent.

1.5. Theorem 3, the case when M is a torsion module. As explained in the introduction, Theorem 3, in the case when M is a torsion module, has been proved (see [Sch], Theorem 21.1). We will use this result for the case N is pseudo-zero. Since $\Delta_0(N) = 1$ if N is pseudo-zero, we have the following result.

Proposition 1.3. *Suppose that N is pseudo-zero. Then $N \sim 0$, i.e.,*

$$|\text{tor}_\mathbb{Z}(N \otimes \mathbb{Z}[A_\Gamma])| \sim 1.$$

1.6. Lattices in Hermitian spaces and \mathbb{Z} -torsion. Suppose W is a finite-dimensional based Hermitian space, i.e. a \mathbb{C} -vector space equipped with an Hermitian product (\cdot, \cdot) and a preferred orthonormal basis. The \mathbb{Z} -submodule $\Lambda \subset W$ spanned by the basis is called the *fundamental lattice*.

For a \mathbb{Z} -submodule (also called a lattice) $\Theta \subset \Lambda$ with \mathbb{Z} -basis v_1, \dots, v_l define

$$\text{vol}(\Theta) = \left| \det((v_i, v_j)_{i,j=1}^l) \right|^{1/2}.$$

It is clear that $\text{vol}(\Theta) \geq 1$.

For a lattice $\Theta \subset \Lambda$ define its orthogonal complement in Λ by

$$\Theta^\perp = \{x \in \Lambda \mid (x, y) = 0 \text{ for all } y \in \Theta\}.$$

It is clear that $\Theta \subset \Theta^{\perp\perp}$. A lattice Θ is *primitive* if $\Theta = \Theta^{\perp\perp}$. It is known that Θ is primitive if and only if it is cut out by a subspace, i.e. $\Theta = (\Theta \otimes_\mathbb{Z} \mathbb{Q}) \cap \Lambda$; and if Θ is primitive, then (see e.g. [Ber])

$$\text{vol}(\Theta)^2 = |\Lambda/(\Theta \oplus \Theta^\perp)|. \quad (4)$$

Lemma 1.4. *For $i = 1, 2$ let W_i be a finite-dimensional based Hermitian space with fundamental lattice Λ_i . Suppose $f : \Theta_1 \rightarrow \Theta_2$ is a \mathbb{Z} -linear map, where $\Theta_i \subset \Lambda_i$ is a lattice of maximal rank. Then*

$$|\mathrm{tor}_{\mathbb{Z}}(\mathrm{coker} f)| \leq \|f\|^{\mathrm{rk} \Theta_2} \mathrm{vol}(\Theta_1).$$

Here $\|f\|$ is the norm of the linear extension of f to a \mathbb{C} -linear operator from W_1 to W_2 .

Proof. Let $\overline{f(\Theta_1)} = (f(\Theta_1) \otimes_{\mathbb{Z}} \mathbb{Q}) \cap \Lambda_2$. Then

$$\mathrm{tor}_{\mathbb{Z}}(\mathrm{coker} f) = \overline{f(\Theta_1)} / f(\Theta_1).$$

Hence

$$\begin{aligned} |\mathrm{tor}_{\mathbb{Z}}(\mathrm{coker} f)| &= |\overline{f(\Theta_1)} / f(\Theta_1)| \\ &= \mathrm{vol}(f(\Theta_1) / \overline{f(\Theta_1)}). \end{aligned} \quad (5)$$

It is known that

$$\det'(f) \mathrm{vol}(\Theta_1) = \mathrm{vol}(\ker f) \mathrm{vol}(f(\Theta_1)), \quad (6)$$

where $\det'(f)$ is the product of all non-zero singular values of f , i.e. the square root of the product of all non-zero eigenvalues of $f^* f$. From (6) and (5) we have

$$|\mathrm{tor}_{\mathbb{Z}}(\mathrm{coker} f)| = \frac{\det'(f) \mathrm{vol}(\Theta_1)}{\mathrm{vol}(\ker f) \mathrm{vol}(\overline{f(\Theta_1)})} \leq \det'(f) \mathrm{vol}(\Theta_1). \quad (7)$$

The maximal singular value of f is equal to $\|f\|$. The number of non-zero singular values is less than or equal to the rank of f . Hence $\det' f$, being the product of the non-zero singular value, is $\leq \|f\|^{\mathrm{rk} L^2}$. Now from (7) we get the lemma. \square

1.7. Decomposition of the group ring of a finite abelian group

1.7.1. Decomposition over \mathbb{C} . Suppose A is a finite abelian group. The group ring $\mathbb{C}[A]$ is a \mathbb{C} -vector space of dimension $|A|$. Equip $\mathbb{C}[A]$ with a Hermitian product so that A is an orthonormal basis. Then the integral group ring $\mathbb{Z}[A]$ is the corresponding fundamental lattice.

The theory of representations of A over \mathbb{C} is easy: $\mathbb{C}[A]$ decomposes as a direct sum of mutually orthogonal one-dimensional A -modules:

$$\mathbb{C}[A] = \bigoplus_{\chi \in \hat{A}} \mathbb{C} e_{\chi}, \quad (8)$$

where e_χ is the idempotent

$$e_\chi = \frac{1}{|A|} \sum_{a \in A} \chi(a^{-1})a.$$

The vector subspaces $\mathbb{C}e_\chi$ are not only orthogonal with respect to the Hermitian structure, but also orthogonal with respect to the ring structure in the sense that $e_\chi e_{\chi'} = 0$ if $\chi \neq \chi'$. Each $\mathbb{C}e_\chi$ is an ideal of the ring $\mathbb{Z}[A]$.

For a \mathbb{Z} -submodule $X \subset \mathbb{Z}[A]$ let $X_{\mathbb{C}}$ the \mathbb{C} -vector space spanned by X .

1.7.2. Decomposition corresponding to a subgroup. The integral group ring $\mathbb{Z}[A]$ does not have as nice a decomposition as (8). Given a subgroup $B \subset A$, we decompose a subring of $\mathbb{Z}[A]$ as follows.

The natural projection $A \rightarrow A/B$ gives rise to the exact sequence

$$0 \rightarrow \beta(B) \rightarrow \mathbb{Z}[A] \rightarrow \mathbb{Z}[A/B] \rightarrow 0, \quad (9)$$

where $\beta(B)$ is the ideal of $\mathbb{Z}[A]$ generated by $1 - b$, $b \in B$. As a lattice of $\mathbb{Z}[A]$, $\beta(B)$ is primitive.

Let $\alpha(B)$ be the annihilator of $\beta(B)$:

$$\alpha(B) = \{x \in \mathbb{Z}[A] \mid xy = 0 \text{ for all } y \in \beta(B)\}.$$

Then $\alpha(B)$ is also the orthogonal complement of $\beta(B)$ in $\mathbb{Z}[A]$. It is known that $\alpha(B)$ is the principal ideal generated by $u = u_B := \sum_{b \in B} b$, see e.g. [BM].

The complexification $\alpha_{\mathbb{C}}(B)$ and $\beta_{\mathbb{C}}(B)$ are easy to describe. Tensoring (9) with \mathbb{C} ,

$$0 \rightarrow \beta_{\mathbb{C}}(B) \rightarrow \mathbb{C}[A] \rightarrow \mathbb{C}[A/B] \rightarrow 0.$$

As a $\mathbb{C}[A]$ -module, $\mathbb{C}[A/B]$ is isomorphic to $\alpha_{\mathbb{C}}(B) = \beta_{\mathbb{C}}(B)^{\perp}$, and

$$\alpha_{\mathbb{C}}(B) = \beta_{\mathbb{C}}(B)^{\perp} = \bigoplus_{\chi \in \hat{A}, \chi|_B = 1} \mathbb{C}e_\chi, \quad (10)$$

$$\text{rk}_{\mathbb{Z}}(\alpha(B)) = \dim_{\mathbb{C}}(\alpha_{\mathbb{C}}(B)) = |A|/|B|. \quad (11)$$

Proposition 1.5. *The finite group $\mathbb{Z}[A]/(\alpha(B) \oplus \beta(B))$ has order $|B|^{|A|/|B|}$.*

Proof. Let $y_1, \dots, y_\ell \in A$ be representatives of cosets of B in A . Then $\ell = |A/B|$, and the elements $y_1 u_B, \dots, y_\ell u_B$ form a \mathbb{Z} -basis of $\alpha(B)$. It is easy to see that $(y_i u_B, y_j u_B) = 0$ if $i \neq j$. The length of each vector $y_j u_B$ is $(\sum_{b \in B} \|y_j b\|^2)^{1/2} = |B|^{1/2}$. It follows that

$$\text{vol}(\alpha(B)) = |B|^{\ell/2}.$$

From (4) we have $|\mathbb{Z}[A]/(\alpha(B) \oplus \beta(B))| = |B|^\ell = |B|^{|A|/|B|}$. \square

1.7.3. Decomposition corresponding to a collection of subgroups. Suppose that B_1, \dots, B_k are subgroups of a finite abelian group A . Let

$$\alpha(B_1, \dots, B_k) = \sum_{j=1}^k \alpha(B_j), \quad \beta(B_1, \dots, B_k) = \bigcap_{j=1}^k \beta(B_j).$$

Then $\alpha(B_1, \dots, B_k)$ and $\beta(B_1, \dots, B_k)$ are primitive lattices of $\mathbb{Z}[A]$, and they are orthogonal complement of each other in $\mathbb{Z}[A]$. In addition, both $\alpha(B_1, \dots, B_k)$ and $\beta(B_1, \dots, B_k)$ are ideals of $\mathbb{Z}[A]$, and they are the annihilator of each other.

Proposition 1.6. *The finite group $\mathbb{Z}[A]/(\beta(B_1, \dots, B_k) \oplus \alpha(B_1, \dots, B_k))$ has order less than or equal to $\prod_{j=1}^k |B_j|^{|A|/|B_j|}$. Equivalently,*

$$\text{vol}(\alpha(B_1, \dots, B_k)) \leq \left(\prod_{j=1}^k |B_j|^{|A|/|B_j|} \right)^{1/2}.$$

Proof. We write $\alpha = \alpha(B_1, \dots, B_k)$ and $\beta = \beta(B_1, \dots, B_k)$. Recall that $\beta = \bigcap_{j=1}^k \beta(B_j)$. We have

$$\mathbb{Z}[A]/(\alpha + \beta) \cong (\mathbb{Z}[A]/\alpha)/\beta = (\mathbb{Z}[A]/\alpha)/\left(\bigcap_{j=1}^k \beta(B_j)\right).$$

Since $(\mathbb{Z}[A]/\alpha)/\left(\bigcap_{j=1}^k \beta(B_j)\right)$ injects in

$$\prod_{j=1}^k (\mathbb{Z}[A]/\alpha)/\beta(B_j) = \prod_{j=1}^k (\mathbb{Z}[A]/\beta(B_j))/\alpha,$$

we have

$$|\mathbb{Z}[A]/(\alpha + \beta)| \leq \prod_{j=1}^k |(\mathbb{Z}[A]/\beta(B_j))/\alpha|. \quad (12)$$

Since $\alpha(B_j) \subset \alpha$, $(\mathbb{Z}[A]/\beta(B_j))/\alpha(B_j)$ surjects onto $(\mathbb{Z}[A]/\beta(B_j))/\alpha$, hence

$$|(\mathbb{Z}[A]/\beta(B_j))/\alpha| \leq |(\mathbb{Z}[A]/\beta(B_j))/\alpha(B_j)| = |\mathbb{Z}[A]/(\alpha(B_j) + \beta(B_j))|. \quad (13)$$

Inequalities (12) and (13), together with Proposition 1.5, show that

$$|\mathbb{Z}[A]/(\alpha \oplus \beta)| \leq \prod_{j=1}^k |B_j|^{|A|/|B_j|}.$$

The equivalence between the two statements follows from (4). \square

1.8. Torsion points in algebraic varieties. We recall well-known facts about algebraic subgroups of $(\mathbb{C}^*)^n$.

1.8.1. Algebraic subgroups of $(\mathbb{C}^*)^n$. With respect to the usual multiplication $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$ is an abelian group, and so is $(\mathbb{C}^*)^n$. An *algebraic subgroup* of $(\mathbb{C}^*)^n$ is a subgroup which is closed in the Zariski topology.

For a lattice, i.e. a subgroup, Λ of \mathbb{Z}^n , not necessarily of maximal rank, recall that $I(\Lambda)$ is the ideal of \mathfrak{R} generated by $1 - t^k$, $k \in \Lambda$. Let $G(\Lambda) = V_{I(\Lambda)}$, the zero-set of $I(\Lambda)$, i.e. the set of all $z \in \mathbb{C}^n$ such that $z^k - 1 = 0$ for every $k \in \Lambda$. Here for $k = (k_1, \dots, k_n) \in \mathbb{Z}^n$ and $z = (z_1, \dots, z_n) \in (\mathbb{C}^*)^n$ we set $t^k = \prod_i t_i^{k_i}$ and $z^k = \prod_i z_i^{k_i}$.

It is easy to see that $G(\Lambda)$ is an algebraic subgroup. The converse holds true: Every algebraic subgroup is equal to $G(\Lambda)$ for some lattice Λ , see [Sch-W].

Every element $z \in G(\Lambda)$ defines a character χ_z of the quotient group $A_\Lambda := \mathbb{Z}^n / \Lambda$ via

$$\chi_z(t^k) = z^k,$$

and conversely, every character of A_Λ arises in this way. Thus one can identify $G(\Lambda)$ with $\text{Hom}(A_\Lambda, \mathbb{C}^*)$ via $z \rightarrow \chi_z$. We will write e_z for the idempotent χ_{χ_z} , and the decomposition (8), with Λ having maximal rank, now becomes

$$\mathbb{C}[A_\Lambda] = \bigoplus_{z \in G(\Lambda)} \mathbb{C} e_z. \quad (14)$$

1.8.2. Torsion points. A point $z \in (\mathbb{C}^*)^n$ is a *torsion point* if it is a torsion element of the multiplicative group $(\mathbb{C}^*)^n$. Let \mathbb{U} denote the set of all roots of unity in \mathbb{C}^* . Then the set of torsion points of $(\mathbb{C}^*)^n$ is \mathbb{U}^n . For example, if $\Gamma \subset \mathbb{Z}^n$ is a lattice of maximal rank, then $G(\Gamma) \subset \mathbb{U}^n$.

The following fact is well known in the theory of torsion points on algebraic varieties.

Proposition 1.7. *Suppose $X \neq \mathbb{C}^n$ is an algebraic subset of \mathbb{C}^n defined over \mathbb{Q} . There exist a finite number of non-zero lattices $\Lambda_1, \dots, \Lambda_k$ in \mathbb{Z}^n , such that $\mathbb{U}^n \cap X \subset \mathbb{U}^n \cap (\bigcup_{j=1}^k G(\Lambda_j))$, i.e. any torsion point in X belongs to $\bigcup_{j=1}^k G(\Lambda_j)$.*

Proof. A *torsion coset* is a coset uG , where u is torsion point and G is an algebraic subgroup of $(\mathbb{C}^*)^n$. It is well known that there is a finite number of torsion cosets $u_j G_j \subset X$ such that every torsion point in X belongs to $\bigcup_j u_j G_j$, see [Lau], [Sch-W]. Since $u_j G_j \subset X$, the dimension of G_j is at most $n - 1$. Let U_j be the finite cyclic group generated by u_j . Then $U_j G_j$ is also an algebraic group of dimension $\leq n - 1$. Hence $U_j G_j = G(\Lambda_j)$, with Λ_j a non-zero lattice. Since $u_j G_j \subset U_j G_j$, it is clear that every torsion point in X belongs to $\bigcup_j G(\Lambda_j)$. \square

1.9. Elementary bounds from exact sequences

Lemma 1.8. *Suppose M is a finitely generated \mathfrak{R} -module with a free resolution*

$$\dots \rightarrow \mathfrak{R}^{m_2} \xrightarrow{\partial_2} \mathfrak{R}^{m_1} \xrightarrow{\partial_1} \mathfrak{R}^{m_0} \rightarrow M \rightarrow 0,$$

and Q is a \mathfrak{R} -module with $|Q| < \infty$. Then $|\mathrm{Tor}_i^{\mathfrak{R}}(M, Q)| \leq |Q|^{m_i}$ for every $i = 0, 1, \dots$

Proof. By definition, $\mathrm{Tor}_i^{\mathfrak{R}}(M, Q)$ is the homology groups of the complex

$$\dots \rightarrow \mathfrak{R}^{m_2} \otimes Q \rightarrow \mathfrak{R}^{m_1} \otimes Q \rightarrow \mathfrak{R}^{m_0} \otimes Q \rightarrow 0.$$

Since the i -th term of the this complex is $\mathfrak{R}^{m_i} \otimes Q \cong Q^{m_i}$, a finite group of order $|Q|^{m_i}$, its i -th homology group has $\leq |Q|^{m_i}$ elements. \square

Lemma 1.9. *Suppose in an exact sequence of abelian groups*

$$\dots \rightarrow N_1(\Gamma) \rightarrow M_1(\Gamma) \rightarrow M_2(\Gamma) \rightarrow N_2(\Gamma) \rightarrow \dots$$

each $M_i(\Gamma)$, $N_i(\Gamma)$ is an abelian group depending on subgroups $\Gamma \subset \mathbb{Z}^n$ of finite index. Assume further that $N_1(\Gamma)$ and $N_2(\Gamma)$ are finite, and

$$|N_1(\Gamma)| \sim 1 \sim |N_2(\Gamma)|.$$

Then

$$|\mathrm{tor}_{\mathbb{Z}}(M_1(\Gamma))| \sim |\mathrm{tor}_{\mathbb{Z}}(M_2(\Gamma))|.$$

Proof. Replacing N_1 by an appropriate quotient and N_2 by an appropriate subgroup, we may assume that

$$0 \rightarrow N_1(\Gamma) \rightarrow M_1(\Gamma) \rightarrow M_2(\Gamma) \rightarrow N_2(\Gamma) \rightarrow 0$$

is exact. We then have

$$M_1/N_1 \hookrightarrow M_2 \twoheadrightarrow N_2. \tag{15}$$

The inclusion in (15) shows that

$$|\mathrm{tor}_{\mathbb{Z}}(M_1)|/|N_1| \leq |\mathrm{tor}_{\mathbb{Z}}(M_2)|,$$

and the surjection in (15) shows that

$$\mathrm{tor}_{\mathbb{Z}}(M_2) \leq |N_2| |\mathrm{tor}_{\mathbb{Z}}(M_1/N_1)| = |N_2| |\mathrm{tor}_{\mathbb{Z}}(M_1)|/|N_1|.$$

From there we get the conclusion of the lemma. \square

Lemma 1.10. Suppose Θ is a free abelian group of finite rank, and $p, q: \Theta \rightarrow \Theta$ are \mathbb{Z} -linear operators with p injective. Let \mathcal{C} be the complex

$$0 \rightarrow \Theta \xrightarrow{\partial_2} \Theta^2 \xrightarrow{\partial_1} \Theta \rightarrow 0,$$

where $\partial_2(a) = (-q(a), p(a))$, $\partial_1(a, b) = p(a) + q(b)$. Then the homology groups of \mathcal{C} are finite, and

$$|H_1(\mathcal{C})| = |H_0(\mathcal{C})|.$$

Proof. The complex \mathcal{C} is the middle row of the commutative diagram with exact columns:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & 0 & \longrightarrow & \Theta & \xrightarrow{p} & \Theta & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow i_1 & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \Theta & \xrightarrow{\partial_2} & \Theta \oplus \Theta & \xrightarrow{\partial_1} & \Theta & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \Theta & \xrightarrow{p} & \Theta & \longrightarrow & 0 & \longrightarrow & 0. \end{array}$$

Here $i_1(a) = (a, 0)$, $i_2(a, b) = b$. Let \mathcal{C}_1 be the first row and \mathcal{C}_2 the last row. Then $0 \rightarrow \mathcal{C}_1 \rightarrow \mathcal{C} \rightarrow \mathcal{C}_2 \rightarrow 0$ is exact. The long exact sequence, together with $H_1(\mathcal{C}_1) = H_0(\mathcal{C}_2)$, gives us the following exact sequence

$$0 \rightarrow H_1(\mathcal{C}) \rightarrow H_1(\mathcal{C}_2) \rightarrow H_0(\mathcal{C}_1) \rightarrow H_0(\mathcal{C}) \rightarrow 0.$$

Note that $|H_1(\mathcal{C}_2)| = |H_0(\mathcal{C}_1)| = |\operatorname{coker} p|$, which is finite since p is injective. It follows that $H_1(\mathcal{C})$ and $H_0(\mathcal{C})$ are finite. In an exact sequence of finite abelian groups, the alternating product of the cardinalities is 1. Hence, with the two middle terms having $|H_1(\mathcal{C}_2)| = |H_0(\mathcal{C}_1)|$, we must have $|H_1(\mathcal{C})| = |H_0(\mathcal{C})|$. \square

2. Approximation of $\mathbb{Z}[\mathbb{Z}^n/\Gamma]$

2.1. Approximation of $\mathbb{Z}[A_\Gamma]$: formulation of results. As mentioned in the introduction, we search for a good approximation of $\mathbb{Z}[\mathbb{Z}^n/\Gamma]$ as $\langle \Gamma \rangle \rightarrow \infty$. The approximation depends on some extra choice, namely, a non-zero element $p \in \mathfrak{R}$.

Fix a non-zero Laurent polynomial $p \in \mathfrak{R} = \mathbb{Z}[t^{\pm 1}, \dots, t^{\pm n}]$. For each subgroup $\Gamma \subset \mathbb{Z}^n$ of rank n we will construct an \mathfrak{R} -module $\beta(p; \Gamma)$ with the following properties.

Proposition 2.1. (i) For every finitely generated \mathfrak{R} -module M one has

$$|\operatorname{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[A_\Gamma])| \sim |\operatorname{tor}_{\mathbb{Z}}(M \otimes \beta(p; \Gamma))|.$$

(ii) Suppose $p \in I$, where $I \neq \mathfrak{R}$ is ideal of \mathfrak{R} such that \mathfrak{R}/I is pseudo-zero. Then for each $i = 0, 1$ the module $\text{Tor}_i^{\mathfrak{R}}(\beta(p; \Gamma), \mathfrak{R}/I)$ is finite, and

$$|\text{Tor}_i^{\mathfrak{R}}(\beta(p; \Gamma), \mathfrak{R}/I)| \sim 1.$$

The remaining part of this section is devoted to the construction of $\beta(p; \Gamma)$ and the proof of Proposition 2.1.

2.2. Heuristics. There is no rigorous mathematics in this subsection. Logically the reader can skip this subsection.

In the estimate of $\text{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[A_{\Gamma}])$ using exact sequences, finiteness is very helpful. We will try to decompose $\mathbb{Z}[A]$ as a sum of two submodules, one is negligible, and the other is finite if tensoring with pre-given modules.

We have the decomposition (14) of $\mathbb{C}[A_{\Gamma}]$ into irreducible components

$$\mathbb{C}[A_{\Gamma}] = \bigoplus_{z \in G(\Gamma)} \mathbb{C} e_z.$$

The module $M \otimes \mathbb{C}[A_{\Gamma}]$ will decompose accordingly. Albeit over \mathbb{C} , this gives us hint that some $z \in G(\Gamma)$ are “good” and some are “bad”. Here a good z must satisfy some non-degeneracy property, and if a point is good, all its Galois conjugates are good. Combining all good points together one should get some “integral” sub-module of $\mathbb{Z}[A_{\Gamma}]$ for which non-degeneracy conditions imply some kind of finiteness. If S is the set of all bad points, and S^{\perp} be its complement in $G(\Gamma)$, then one has

$$\mathbb{C}[A_{\Gamma}] = \left(\bigoplus_{z \in S} \mathbb{C} e_z \right) \oplus \left(\bigoplus_{z \in S^{\perp}} \mathbb{C} e_z \right).$$

The module β would be the “integral spine” of the second part.

The set of bad points will consists of those in $G(\Gamma)$ which are zeros p . For good points z , $p(z) \neq 0$, and this will give us the non-degeneracy condition. We control the set of bad points, which is the intersection $G(\Gamma) \cap V_p$, by using theory of torsion points on V_p , see Proposition 1.7.

2.3. Definition of $\beta(p; \Gamma)$. The zero set

$$V_p := \{(z_1, \dots, z_n) \in (\mathbb{C}^*)^n \mid p(z_1, \dots, z_n) = 0\}$$

is an algebraic subset of $(\mathbb{C}^*)^n$ of dimension $\leq n-1$. Let $\Lambda_1, \dots, \Lambda_k$ be the non-zero subgroups of \mathbb{Z}^n described in Proposition 1.7 with $X = V_p$. By construction,

$$\text{if a torsion point } z \text{ does not belong to } \bigcup_{j=1}^k G(\Lambda_j), \text{ then } p(z) \neq 0. \quad (16)$$

Suppose $A = A_\Gamma := \mathbb{Z}^n / \Gamma$, where $\Gamma \subset \mathbb{Z}^n$ is a subgroup of maximal rank n . The abelian groups $B_j = (\Lambda_j + \Gamma) / \Gamma$ are subgroups of $A = \mathbb{Z}^n / \Gamma$. Let

$$\alpha(p; \Gamma) = \alpha(B_1, \dots, B_k), \quad \beta(p; \Gamma) = \beta(B_1, \dots, B_k),$$

where $\alpha(B_1, \dots, B_k)$ and $\beta(B_1, \dots, B_k)$ are ideals of $\mathbb{Z}[A]$ defined as in Section 1.7.3.

We partition $G(\Gamma) = \hat{A}$ into two disjoint subsets S, S^\perp by

$$S = G(\Gamma) \cap \left(\bigcup_j G(\Lambda_j) \right), \quad S^\perp = G(\Gamma) \setminus S. \quad (17)$$

We will see that as $\langle \Gamma \rangle \rightarrow \infty$, S is small compared to its complement S^\perp . Note that χ_z , with $z \in G(\Gamma)$, takes value 1 on B_j exactly when $z \in G(\Lambda_j)$. Hence from (10) we have

$$\alpha_{\mathbb{C}}(B_j) = \bigoplus_{z \in G(\Gamma) \cap G(\Lambda_j)} \mathbb{C}_z,$$

and hence

$$\beta_{\mathbb{C}}(p; \Gamma) = \bigoplus_{z \in S^\perp} \mathbb{C}_z, \quad \alpha_{\mathbb{C}}(p; \Gamma) = \bigoplus_{z \in S} \mathbb{C}_z. \quad (18)$$

We will write $\alpha = \alpha(p; \Gamma), \beta = \beta(p; \Gamma)$. Let $\text{pr}: \mathfrak{R} \rightarrow \mathbb{Z}[A_\Gamma]$ be the canonical projection. Note that $\text{pr}^{-1}(0)$ is the ideal of all polynomials taking values 0 at every point of $G(\Gamma)$. Similarly, $\tilde{\alpha} = \text{pr}^{-1}(\alpha)$ is the ideal of all polynomials taking values 0 at every point of S^\perp . Over \mathbb{C} , $\tilde{\alpha}_{\mathbb{C}}$ is the reduced ideal of $\mathfrak{R}_{\mathbb{C}} = \mathbb{C}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ whose zero set is S^\perp , $V_{\tilde{\alpha}_{\mathbb{C}}} = S^\perp$. In addition,

$$\mathfrak{R}_{\mathbb{C}} / \tilde{\alpha}_{\mathbb{C}} \cong \mathbb{C}[A] / \alpha_{\mathbb{C}} \cong \beta_{\mathbb{C}}. \quad (19)$$

The important facts concerning $\alpha(p; \Gamma)$ and $\beta(p; \Gamma)$ are the following.

Lemma 2.2. a) $S^\perp = V_{\tilde{\alpha}_{\mathbb{C}}}$ does not intersect V_p . It follows that the ideal of $\mathfrak{R}_{\mathbb{C}}$ generated by p and $\tilde{\alpha}_{\mathbb{C}}$ is the whole $\mathfrak{R}_{\mathbb{C}}$.

b) The multiplication map $p: \beta \rightarrow \beta, x \rightarrow p \cdot x$, is injective. It follows that $\text{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/(p), \beta) = 0$.

c) The quotient group $Q(p; \Gamma) := \mathbb{Z}[A_\Gamma] / (\alpha(p; \Gamma) \oplus \beta(p; \Gamma))$ is finite and its order is negligible, $|Q(p; \Gamma)| \sim 1$.

d) $|S| = \text{rk}_{\mathbb{Z}} \alpha(p; \Gamma)$ is small compared to the $\text{rk}_{\mathbb{Z}} \mathbb{Z}[A_\Gamma] = |A_\Gamma|$ in the sense that

$$\lim_{\langle \Gamma \rangle \rightarrow \infty} \frac{\text{rk}_{\mathbb{Z}} \alpha(p; \Gamma)}{|A_\Gamma|} = 0.$$

e) One has $\text{vol}(\alpha) \sim 1$.

Proof. a) Suppose $\mathbf{z} \in S^\perp$. By definition (17), \mathbf{z} is a torsion point not belonging to $\bigcup_{j=1}^k G(\Lambda_j)$. By (16), $p(\mathbf{z}) \neq 0$. In other words, $V_p \cap V_{\tilde{\alpha}_\mathbb{C}} = \emptyset$. By the Nullstellensatz, the ideal generated by p and $\tilde{\alpha}_\mathbb{C}$ is the whole $\mathfrak{R}_\mathbb{C}$.

b) Note that $\mathbb{C} e_z$ is a $\mathfrak{R}_\mathbb{C}$ -module by the action $f \cdot e_z = f(\mathbf{z}) e_z$. If $\mathbf{z} \in S^\perp$, then $p(\mathbf{z}) \neq 0$, hence $p: \mathbb{C} e_z \rightarrow \mathbb{C} e_z$ is an isomorphism. Since $\beta_\mathbb{C} = \bigoplus_{\mathbf{z} \in S^\perp} \mathbb{C} e_z$, the map $p: \beta_\mathbb{C} \rightarrow \beta_\mathbb{C}$ is also an isomorphism. It follows that $p: \beta \rightarrow \beta$ is injective.

One has $\text{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/(p), \beta) = \ker(p: \beta \rightarrow \beta, x \rightarrow p \cdot x) = 0$.

c) We will first show that for each $j = 1, \dots, k$,

$$\lim_{\langle \Gamma \rangle \rightarrow \infty} |B_j| = \infty. \quad (20)$$

By definition, $B_j = (\Lambda_j + \Gamma)/\Gamma$. Fix an element $x \in \Lambda_j$, $x \neq 0$, and look at the degree of x in $B_j = (\Lambda_j + \Gamma)/\Gamma$. If $m|x| < \langle \Gamma \rangle$, then $m|x|$ does not belong to Γ by the definition of $\langle \Gamma \rangle$, and hence mx is not 0 in $B_j = (\Lambda_j + \Gamma)/\Gamma$. This means the cyclic subgroup of B_j generated by x has order at least $\langle \Gamma \rangle/|x|$. It follows that $|B_j| \geq \langle \Gamma \rangle/|x|$. Hence $\lim_{\langle \Gamma \rangle \rightarrow \infty} |B_j| = \infty$.

From Proposition 1.6,

$$|Q(p; \Gamma)|^{1/|A|} \leq \prod_{j=1}^k |B_j|^{1/|B_j|},$$

from which together with (20) we get $|Q(p; \Gamma)| \sim 1$.

d) By (11) one has $\text{rk}(\alpha(B_j)) = |A|/|B_j|$. Since $\alpha = \sum \alpha(B_j)$, one gets

$$\text{rk}(\alpha)/|A| \leq \sum_{j=1}^k \text{rk} \alpha(B_j)/|A| = \sum_{j=1}^k (1/|B_j|), \quad (21)$$

which, with (20), shows that $\lim_{\langle \Gamma \rangle \rightarrow \infty} \frac{\text{rk}_\mathbb{Z} \alpha(p; \Gamma)}{|A_\Gamma|} = 0$.

e) This follows immediately from (4) and part (c). \square

2.4. Contribution from $\alpha(p; \Gamma)$ is negligible. The ideals α and β , being $\mathbb{Z}[A_\Gamma]$ -module, can be naturally considered as \mathfrak{R} -modules.

Lemma 2.3. *Suppose M is a finitely generated \mathfrak{R} -module. Then*

$$|\text{tor}_\mathbb{Z}(M \otimes \alpha(p; \Gamma))| \sim 1.$$

Proof. Tensoring the presentation

$$\mathfrak{R}^{m_1} \xrightarrow{\partial_1} \mathfrak{R}^{m_0} \rightarrow M \rightarrow 0$$

with $\mathbb{Z}[A]$ and α respectively, one gets

$$\begin{aligned} (\mathbb{Z}[A_\Gamma])^{m_1} &\xrightarrow{\partial_{1,\Gamma}} (\mathbb{Z}[A_\Gamma])^{m_0} \rightarrow M \otimes \mathbb{Z}[A_\Gamma] \rightarrow 0, \\ \alpha^{m_1} &\xrightarrow{\partial_{1,\alpha}} \alpha^{m_0} \rightarrow M \otimes \alpha \rightarrow 0, \end{aligned} \quad (22)$$

with $\partial_{1,\alpha}$ the restriction of $\partial_{1,\Gamma}$.

Recall that we have a Hermitian structure on $\mathbb{C}[A]$. It is not difficult to find an upper bound, not depending on Γ , for all the operator $\partial_{1,\Gamma}$. In fact, by Lemma 2.5 of [Lü1],

$$\|\mathcal{O}_\Gamma\| \leq D := m_1 m_0 \max_{i,j} \{|\mathcal{O}_{ij}|_1\},$$

where for a Laurent polynomial $a \in \mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ the norm $|a|_1$ is the sum of the absolute values of its coefficients.

Because $\alpha_{\mathbb{C}}$ is an invariant subspace of $\mathbb{C}[A]$, we also have

$$\|\mathcal{O}_\alpha\| \leq D.$$

Applying Lemma 1.4 to the sequence (22) we get

$$|\mathrm{tor}_{\mathbb{Z}}(M \otimes \alpha)| \leq D^{\mathrm{rk}_{\mathbb{Z}} \alpha} \mathrm{vol}(\alpha).$$

The right-hand side has negligible growth, by Lemma 2.2 d) and e). \square

2.5. Proof of Proposition 2.1 part (i)

Proof. Recall that $A = A_\Gamma := \mathbb{Z}^n / \Gamma$. We have an exact sequence

$$0 \rightarrow (\alpha \oplus \beta) \rightarrow \mathbb{Z}[A] \rightarrow Q \rightarrow 0, \quad (23)$$

with $Q = |Q(p; \Gamma)| \sim 1$ by Lemma 2.2 a). Tensoring (23) with M ,

$$\cdots \rightarrow \mathrm{Tor}_1^{\mathfrak{R}}(M, Q) \rightarrow ((M \otimes \alpha) \oplus (M \otimes \beta)) \rightarrow M \otimes \mathbb{Z}[A] \rightarrow \mathrm{Tor}_0^{\mathfrak{R}}(M, Q) \rightarrow 0. \quad (24)$$

Lemma 1.8 shows that $|\mathrm{Tor}_i^{\mathfrak{R}}(M, Q)| < |Q|^{m_i}$ for some constant m_i depending on M only. Since $Q \sim 1$, we also have

$$|\mathrm{Tor}_i^{\mathfrak{R}}(M, Q)| \sim 1.$$

Applying Lemma 1.9 to the sequence (24), we get

$$|\mathrm{tor}_{\mathbb{Z}}(M \otimes \alpha) \oplus \mathrm{tor}_{\mathbb{Z}}(M \otimes \beta)| \sim |\mathrm{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[A])|.$$

Since $|\mathrm{tor}_{\mathbb{Z}}(M \otimes \alpha)| \sim 1$ by Lemma 2.3, we have $|\mathrm{tor}_{\mathbb{Z}}(M \otimes \beta)| \sim |\mathrm{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[A])|$. \square

2.6. The intermediate ideal $J = (p, q)$. To prepare for the proof of Proposition 2.1 part (ii), we first study the ideal $J = (p, q)$, where $q \in \mathfrak{R}$ is co-prime with p . The reason is \mathfrak{R}/J has a simple free resolution, and hence $\text{Tor}_i^{\mathfrak{R}}(\mathfrak{R}/J, \beta(p; \Gamma))$ is easy to study.

Lemma 2.4. *Both modules $\text{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/J, \beta(p; \Gamma))$ and $\text{Tor}_0^{\mathfrak{R}}(\mathfrak{R}/J, \beta(p; \Gamma))$ are finite, have the same cardinality, and have negligible growth, i.e.*

$$|\text{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/J, \beta(p; \Gamma))| = |\text{Tor}_0^{\mathfrak{R}}(\mathfrak{R}/J, \beta(p; \Gamma))| \sim 1. \quad (25)$$

Proof. We have the following free resolution of \mathfrak{R}/J :

$$0 \rightarrow \mathfrak{R} \xrightarrow{d_2} \mathfrak{R} \oplus \mathfrak{R} \xrightarrow{d_1} \mathfrak{R} \rightarrow \mathfrak{R}/J \rightarrow 0, \quad (26)$$

where $d_2(a) = (-qa, pa)$ and $d_1(a, b) = pa + qb$. This can be directly checked easily, or can be deduced from the theory of Koszul complex as follows. Since p, q are co-prime, the sequence (p, q) is a regular sequence of \mathfrak{R} (see Exercise 5 of page 102 of [Ka]). Hence the Koszul complex of (p, q) , which is (26), is a free resolution of \mathfrak{R}/J .

From the free resolution (26), $\text{Tor}_i^{\mathfrak{R}}(\mathfrak{R}/J, \beta)$ is the i -th homology of the complex

$$0 \rightarrow \beta \xrightarrow{(-q, p)} \beta \oplus \beta \xrightarrow{\begin{pmatrix} p \\ q \end{pmatrix}} \beta \rightarrow 0. \quad (27)$$

The module β is a free \mathbb{Z} -module of finite rank, and the map $p: \beta \rightarrow \beta$ is injective, by Lemma 2.2 b). From Lemma 1.10 we see that both $\text{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/J, \beta)$ and $\text{Tor}_0^{\mathfrak{R}}(\mathfrak{R}/J, \beta) = (\mathfrak{R}/J) \otimes \beta$ are finite, and

$$|\text{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/J, \beta)| = |(\mathfrak{R}/J) \otimes \beta|.$$

By Lemma 1.1, \mathfrak{R}/J is pseudo-zero since p and q are co-prime. We have

$$\begin{aligned} |(\mathfrak{R}/J) \otimes \beta| &\sim |(\mathfrak{R}/J) \otimes \mathbb{Z}[A_\Gamma]| && \text{by Proposition 2.1 (i)} \\ &\sim 1 && \text{by Proposition 1.3.} \end{aligned}$$

This completes the proof of the lemma. \square

2.7. Complexification of tor modules. Recall that $\mathfrak{R}_{\mathbb{C}} = \mathbb{C}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$. Observe that

$$\mathfrak{R}_{\mathbb{C}} \cong \mathfrak{R} \otimes_{\mathbb{Z}} \mathbb{C}, \quad \text{and} \quad \mathfrak{R}_{\mathbb{C}} \cong \mathfrak{R} \otimes_{\mathfrak{R}} \mathfrak{R}_{\mathbb{C}}.$$

Let $I_{\mathbb{C}}$ be the \mathbb{C} -span of I in $\mathfrak{R}_{\mathbb{C}}$. Then $I_{\mathbb{C}}$ is also the extension of I from \mathfrak{R} to $\mathfrak{R}_{\mathbb{C}}$.

Lemma 2.5. *As \mathbb{Z} -modules, for every i ,*

$$(\mathrm{Tor}_i^{\mathfrak{R}}(\mathfrak{R}/I, \beta)) \otimes_{\mathbb{Z}} \mathbb{C} \cong \mathrm{Tor}_i^{\mathfrak{R}_{\mathbb{C}}}(\mathfrak{R}_{\mathbb{C}}/I_{\mathbb{C}}, \beta_{\mathbb{C}}).$$

Proof. Since \mathbb{C} is flat over \mathbb{Z} , we have $(\mathfrak{R}/I) \otimes_{\mathbb{Z}} \mathbb{C} \cong \mathfrak{R}_{\mathbb{C}}/I_{\mathbb{C}}$.

Since $\mathfrak{R}_{\mathbb{C}}$ is flat over \mathfrak{R} , we have $(\mathfrak{R}/I) \otimes_{\mathfrak{R}} \mathfrak{R}_{\mathbb{C}} \cong \mathfrak{R}_{\mathbb{C}}/I_{\mathbb{C}}$. It follows that

$$(\mathfrak{R}/I) \otimes_{\mathbb{Z}} \mathbb{C} \cong (\mathfrak{R}/I) \otimes_{\mathfrak{R}} \mathfrak{R}_{\mathbb{C}}. \quad (28)$$

Suppose $\mathcal{C} \rightarrow \beta$ is a free resolution of β . By definition,

$$\mathrm{Tor}_i^{\mathfrak{R}}(\mathfrak{R}/I, \beta) = H_i(\mathcal{C} \otimes_{\mathfrak{R}} (\mathfrak{R}/I)). \quad (29)$$

Tensoring (29) with $\mathfrak{R}_{\mathbb{C}}$, a flat \mathfrak{R} -module, we get

$$\mathrm{Tor}_i^{\mathfrak{R}_{\mathbb{C}}}(\mathfrak{R}_{\mathbb{C}}/I_{\mathbb{C}}, \beta_{\mathbb{C}}) = H_i(\mathcal{C} \otimes_{\mathfrak{R}} (\mathfrak{R}/I) \otimes_{\mathfrak{R}} \mathfrak{R}_{\mathbb{C}}). \quad (30)$$

Tensoring (29) over \mathbb{Z} with \mathbb{C} , a flat \mathbb{Z} -module, we get

$$(\mathrm{Tor}_i^{\mathfrak{R}}(\mathfrak{R}/I, \beta)) \otimes_{\mathbb{Z}} \mathbb{C} \cong H_i((\mathcal{C} \otimes_{\mathfrak{R}} (\mathfrak{R}/I)) \otimes_{\mathbb{Z}} \mathbb{C}). \quad (31)$$

Since \mathcal{C} is free, each term of \mathcal{C} is a direct of several \mathfrak{R} . It follows from (28) that the right-hand sides of (30) and (31) are isomorphic as \mathbb{Z} -modules, whence the lemma. \square

2.8. Proof of Proposition 2.1 (ii)

Proof. a) The case $i = 0$. Recall that $\mathrm{Tor}_0^{\mathfrak{R}}(\mathfrak{R}/I, \beta(p; \Gamma)) = (\mathfrak{R}/I) \otimes \beta$.

Since $(p) \subset I$, we have a natural surjection $\mathfrak{R}/(p) \twoheadrightarrow \mathfrak{R}/I$. Tensoring with $\beta = \beta(p; \Gamma)$ we get a surjective map

$$\beta \otimes (\mathfrak{R}/(p)) \twoheadrightarrow \beta \otimes (\mathfrak{R}/I).$$

Now $\beta \otimes (\mathfrak{R}/(p)) \cong \beta/p$, which is finite since p acts on the finite-rank free abelian group β by an injection, see Lemma 2.2 b). It follows that $\beta \otimes \mathfrak{R}/I$ is finite.

Since \mathfrak{R}/I is pseudo-zero, by Proposition 2.1 (i) and Proposition 1.3,

$$|\beta \otimes_{\mathfrak{R}} (\mathfrak{R}/I)| \sim |\mathbb{Z}[A_{\Gamma}] \otimes_{\mathfrak{R}} (\mathfrak{R}/I)| \sim 1.$$

b) The case $i = 1$. First we show that $\mathrm{Tor}_1^{\mathfrak{R}}(\beta, \mathfrak{R}/I)$ is finite.

By Lemma 2.2 a), the $\mathfrak{R}_{\mathbb{C}}$ -ideal generated by $\tilde{\alpha}_{\mathbb{C}}$ and p is $\mathfrak{R}_{\mathbb{C}}$, hence $\tilde{\alpha}_{\mathbb{C}} + I_{\mathbb{C}} = \mathfrak{R}_{\mathbb{C}}$ because $p \in I$. It is well known then (see e.g. [AM], Chapter 1)

$$\tilde{\alpha}_{\mathbb{C}} \cap I_{\mathbb{C}} = \tilde{\alpha}_{\mathbb{C}} I_{\mathbb{C}}. \quad (32)$$

In a commutative ring R , it is known that $\mathrm{Tor}_1^R(R/I_1, R/I_2) \cong I_1 \cap I_2 / I_1 I_2$ for two ideals I_1, I_2 . Hence from (32) we have

$$\mathrm{Tor}_1^{\mathfrak{R}\mathbb{C}}(\mathfrak{R}_{\mathbb{C}}/\tilde{\alpha}_{\mathbb{C}}, \mathfrak{R}_{\mathbb{C}}/I_{\mathbb{C}}) = 0.$$

Since $\beta_{\mathbb{C}} = \mathfrak{R}_{\mathbb{C}}/\tilde{\alpha}_{\mathbb{C}}$, this can be rewritten as

$$\mathrm{Tor}_1^{\mathfrak{R}\mathbb{C}}(\beta_{\mathbb{C}}, \mathfrak{R}_{\mathbb{C}}/I_{\mathbb{C}}) = 0,$$

which, by Lemma 2.5, implies that

$$(\mathrm{Tor}_1^{\mathfrak{R}}(\beta, \mathfrak{R}/I)) \otimes_{\mathbb{Z}} \mathbb{C} = 0. \quad (33)$$

Since $\mathrm{Tor}_1^{\mathfrak{R}}(\beta, \mathfrak{R}/I)$ is a finitely generated abelian group, (33) is equivalent to the fact $\mathrm{Tor}_1^{\mathfrak{R}}(\beta, \mathfrak{R}/I)$ is finite.

Now we show that $|\mathrm{Tor}_1^{\mathfrak{R}}(\beta, \mathfrak{R}/I)| \sim 1$. Since \mathfrak{R}/I is pseudo-zero, $I \neq (p)$. This means there is $q \in I$ such that q is not divisible by p . Since p is irreducible, p and q are co-prime. Let $J = (p, q)$. Then $(p) \subset J \subset I$.

Tensoring β with the exact sequence

$$0 \rightarrow I/J \rightarrow \mathfrak{R}/J \rightarrow \mathfrak{R}/I \rightarrow 0$$

we get the exact sequence

$$\cdots \rightarrow \mathrm{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/J, \beta) \rightarrow \mathrm{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/I, \beta) \rightarrow ((I/J) \otimes \beta) \rightarrow \cdots \quad (34)$$

The module I/J , being a submodule of the pseudo-zero module \mathfrak{R}/J , is also pseudo-zero. Hence by Proposition 2.1 (i) and Proposition 1.3,

$$|\mathrm{tor}_{\mathbb{Z}}((I/J) \otimes \beta)| \sim 1. \quad (35)$$

By Lemma 2.4, $\mathrm{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/J, \beta)$ is finite and has negligible growth,

$$|\mathrm{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/J, \beta)| \sim 1. \quad (36)$$

The middle term of (34), being finite, must satisfy

$$|\mathrm{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/I, \beta)| \leq |\mathrm{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/J, \beta)| |\mathrm{tor}_{\mathbb{Z}}((I/J) \otimes \beta)|$$

and hence by (34) and (36) is negligible, $|\mathrm{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/I, \beta)| \sim 1$. \square

3. Proof of Theorems 6, 3, and 4

3.1. Pseudo-zero kernel

Lemma 3.1. *Suppose M_1, M_2 are finitely generated \mathfrak{R} -module, $I \subset \mathfrak{R}$ is a prime ideal such that \mathfrak{R}/I is pseudo-zero, and*

$$0 \rightarrow \mathfrak{R}/I \rightarrow M_1 \rightarrow M_2 \rightarrow 0 \quad (37)$$

is exact. Then $M_1 \sim M_2$.

Proof. Choose a non-zero irreducible $p \in I$ and let $\beta = \beta(p; \Gamma)$. Tensoring (37) with β , we get the exact sequence

$$\cdots \rightarrow (\mathfrak{R}/I) \otimes \beta \rightarrow M_1 \otimes \beta \rightarrow M_2 \otimes \beta \rightarrow 0. \quad (38)$$

By Proposition 2.1 (ii), $(\mathfrak{R}/I) \otimes \beta$ is finite and $|(\mathfrak{R}/I) \otimes \beta| \sim 1$.

Applying Lemma 1.9 to the sequence (38), we get

$$|\mathrm{tor}_{\mathbb{Z}}(M_1 \otimes \beta)| \sim |\mathrm{tor}_{\mathbb{Z}}(M_2 \otimes \beta)|.$$

By Proposition 2.1 (i), $|\mathrm{tor}_{\mathbb{Z}}(M_i \otimes \beta)| \sim |\mathrm{tor}_{\mathbb{Z}}(M_i \otimes \mathbb{Z}[A_{\Gamma}])|$. Hence we can conclude that

$$|\mathrm{tor}_{\mathbb{Z}}(M_1 \otimes \mathbb{Z}[A_{\Gamma}])| \sim |\mathrm{tor}_{\mathbb{Z}}(M_2 \otimes \mathbb{Z}[A_{\Gamma}])|.$$

This means $M_1 \sim M_2$. □

Lemma 3.2. *Suppose N , M_1 and M_2 are finitely generated \mathfrak{R} -modules, and N is pseudo-zero. If*

$$0 \rightarrow N \rightarrow M_1 \rightarrow M_2 \rightarrow 0 \quad (39)$$

is exact, then $M_1 \sim M_2$.

Proof. It is well known that there is a composition series

$$N = N_s \supset N_{s-1} \supset \cdots \supset N_1 \supset N_0 = 0 \quad (40)$$

such that for each i , $N_{i+1}/N_i \cong \mathfrak{R}/I_i$ for some prime ideal I_i , see e.g. [Bou], Theorem IV.4.1. We use induction on s . The case $s = 1$ has been proved, see Lemma 3.1.

Let $M'_1 = M_1/N_{s-1}$ and $N' := N/N_{s-1} \cong \mathfrak{R}/I$, with $I = I_{s-1}$. From (39) we have

$$0 \rightarrow N' \rightarrow M'_1 \rightarrow M_2 \rightarrow 0. \quad (41)$$

From $M'_1 = M_1/N_{s-1}$, we have

$$0 \rightarrow N_{s-1} \rightarrow M_1 \rightarrow M'_1 \rightarrow 0. \quad (42)$$

Note that N' and N_{s-1} , being either a quotient or a submodule of the pseudo-zero module N , are pseudo-zero. By induction and the case $s = 1$, from the exact sequences (41) and (42), we have

$$M'_1 \sim M_2, \quad M_1 \sim M'_1.$$

Hence $M_1 \sim M_2$. □

3.2. Pseudo-zero quotient

Lemma 3.3. *Suppose N , M_1 and M_2 are finitely generated \mathfrak{R} -modules, and N is pseudo-zero. If*

$$0 \rightarrow M_1 \rightarrow M_2 \rightarrow N \rightarrow 0 \quad (43)$$

is exact, then $M_1 \sim M_2$.

Proof. Again using induction on the length of the composition series (40) like in the proof of Lemma 3.2 we can assume that $N = \mathfrak{R}/I$, where $I \neq \mathfrak{R}$ is a prime module. Choose a non-zero irreducible $p \in I$ and let $\beta = \beta(p; \Gamma)$.

Tensoring (43) with β , we have

$$\cdots \rightarrow \text{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/I, \beta) \rightarrow M_1 \otimes \beta \rightarrow M_2 \otimes \beta \rightarrow \text{Tor}_0^{\mathfrak{R}}(\mathfrak{R}/I, \beta) \rightarrow 0. \quad (44)$$

By Proposition 2.1 (ii), $\text{Tor}_i^{\mathfrak{R}}(\mathfrak{R}/I, \beta)$ is finite and $|\text{Tor}_i^{\mathfrak{R}}(\mathfrak{R}/I, \beta)| \sim 1$ for $i = 0, 1$.

Applying Lemma 1.9 to the sequence (40), we get

$$|\text{tor}_{\mathbb{Z}}(M_1 \otimes \beta)| \sim |\text{tor}_{\mathbb{Z}}(M_2 \otimes \beta)|.$$

Using Proposition 2.1 (i), we get

$$|\text{tor}_{\mathbb{Z}}(M_1 \otimes \mathbb{Z}[A_{\Gamma}])| \sim |\text{tor}_{\mathbb{Z}}(M_2 \otimes \mathbb{Z}[A_{\Gamma}])|,$$

which means $M_1 \sim M_2$. □

3.3. Proof of Theorem 6

Proof. Since M_1 and M_2 are pseudo-isomorphic, there are pseudo-zero N_1 and N_2 such that

$$0 \rightarrow N_1 \rightarrow M_1 \rightarrow M_2 \rightarrow N_2 \rightarrow 0$$

is exact. Then we have the following exact sequences:

$$0 \rightarrow M_1/N_1 \rightarrow M_2 \rightarrow N_2 \rightarrow 0, \quad (45)$$

$$0 \rightarrow N_1 \rightarrow M_1 \rightarrow M_1/N_1 \rightarrow 0. \quad (46)$$

From (45) and Lemma 3.3 we have $M_1/N_1 \sim M_2$, while from (46) and Lemma 3.2 we have $M_1/N_1 \sim M_1$. It follows that $M_1 \sim M_2$, which is equivalent to the statement of Theorem 6. □

3.4. The case when M is torsion-free

Proposition 3.4. *Suppose M is a torsion-free finitely generated \mathfrak{R} -module. Then*

$$\lim_{\langle \Gamma \rangle \rightarrow \infty} \frac{\log |\operatorname{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[\mathbb{Z}^n / \Gamma])|}{|\mathbb{Z}^n / \Gamma|} = 0.$$

We first prove the following lemma.

Lemma 3.5. *Suppose N , M_1 and M_2 are finitely generated \mathfrak{R} -modules and*

$$0 \rightarrow M_1 \rightarrow M_2 \rightarrow N \rightarrow 0 \quad (47)$$

is exact. If $M_2 \sim 0$, then $M_1 \sim 0$.

Proof. Using induction on the length of a composition series (40) of N we can assume that $N = \mathfrak{R}/I$, where $I \subset \mathfrak{R}$ is a prime ideal.

If \mathfrak{R}/I is pseudo-zero, then by Lemma 3.3, $M_1 \sim M_2 \sim 0$.

We will consider the remaining case, when \mathfrak{R}/I is not pseudo-zero. Then I is principal, $I = (p)$, where $p \in \mathfrak{R}$.

If $p = 0$, then $N = \mathfrak{R}$ is free, and the sequence (47) is split, $M_2 \cong M_1 \oplus \mathfrak{R}$. One clearly has $\operatorname{tor}_{\mathbb{Z}}(M_1 \otimes \mathbb{Z}[A_{\Gamma}]) = \operatorname{tor}_{\mathbb{Z}}(M_2 \otimes \mathbb{Z}[A_{\Gamma}])$, and the statement follows.

Suppose now $p \neq 0$. Let $\beta = \beta(p; \Gamma)$. Tensoring (47) with β , the following is exact

$$\cdots \rightarrow \operatorname{Tor}_1^{\mathfrak{R}}(\mathfrak{R}/(p), \beta) \rightarrow M_1 \otimes \beta \rightarrow M_2 \otimes \beta \rightarrow \mathfrak{R}/(p) \otimes \beta \rightarrow 0.$$

By Lemma 2.2 b), the first term is 0. It follows that $M_1 \otimes \beta$ is a subgroup of $M_2 \otimes \beta$, and hence

$$|\operatorname{tor}_{\mathbb{Z}}(M_1 \otimes \beta)| \leq |\operatorname{tor}_{\mathbb{Z}}(M_2 \otimes \beta)|.$$

By Proposition 2.1 (i),

$$|\operatorname{tor}_{\mathbb{Z}}(M_i \otimes \beta)| \sim |\operatorname{tor}_{\mathbb{Z}}(M_i \otimes \mathbb{Z}[A_{\Gamma}])|,$$

and since $|\operatorname{tor}_{\mathbb{Z}}(M_2 \otimes \mathbb{Z}[A_{\Gamma}])| \sim 1$, we can conclude that $|\operatorname{tor}_{\mathbb{Z}}(M_1 \otimes \mathbb{Z}[A_{\Gamma}])| \sim 1$, or $M_1 \sim 0$. \square

Proof of Proposition 3.4. Since M is torsion-free, the canonical map $M \rightarrow V := M \otimes_{\mathfrak{R}} \mathcal{F}$, where \mathcal{F} is the fractional field of \mathfrak{R} , is an embedding. This means M is a lattice of V with respect to \mathfrak{R} , and hence there is a free \mathfrak{R} -module F such that M embeds into F , see Chapter 7 of [Bou]. One has an exact sequence of finitely generated \mathfrak{R} -modules

$$0 \rightarrow M \rightarrow F \rightarrow N \rightarrow 0.$$

We have $F \sim 0$ since F is a free \mathfrak{R} -module. From Lemma 3.5 we conclude that $M \sim 0$. \square

3.5. Proof of Theorem 3

Proof. By Theorem 1.2, M is and $\text{tor}(M) \oplus M/\text{tor}(M)$ are pseudo-isomorphic. Hence by Theorem 6,

$$M \sim (\text{tor}(M) \oplus M/\text{tor}(M)).$$

Since $M/\text{tor}(M)$ is torsion-free, by Theorem 3.4, $M/\text{tor}(M) \sim 0$. Hence we have

$$M \sim \text{tor}(M). \quad (48)$$

The proof is thus reduced to the case when M is a torsion module, which had been proved by K. Schmidt, see [Sch], Theorem 21.1. \square

3.6. Proof of Theorem 4

Proof. Suppose \mathcal{D} is a chain complex of free finitely generated modules over a domain R ,

$$\dots \rightarrow R^{m_{i+1}} \xrightarrow{\partial_{i+1}} R^{m_i} \xrightarrow{\partial_i} \dots.$$

For our application either $R = \mathfrak{R}$ or $R = \mathbb{Z}$.

In the exact sequence

$$0 \rightarrow (\ker \partial_i / \text{Im } \partial_{i+1}) \rightarrow (R^{m_i} / \text{Im } \partial_{i+1}) \rightarrow (R^{m_i} / \ker \partial_i) \rightarrow 0$$

the first module is $H_i(\mathcal{D})$, the second $\text{coker } \partial_{i+1}$. Since the third is a torsion-free R -module, one has

$$\text{tor}_R(H_i(\mathcal{D})) = \text{tor}_R(\text{coker } \partial_{i+1}). \quad (49)$$

Suppose now \mathcal{C} is a chain complex of free finitely generated \mathfrak{R} -modules of the form

$$\dots \rightarrow \mathfrak{R}^{m_{i+1}} \xrightarrow{\partial_{i+1}} \mathfrak{R}^{m_i} \xrightarrow{\partial_i} \dots.$$

Applying (49) to the above chain complex, we have

$$\text{tor}(H_i(\mathcal{C})) = \text{tor}(M), \quad (50)$$

where $M = \text{coker } \partial_{i+1}$ which has a presentation

$$\mathfrak{R}^{m_{i+1}} \xrightarrow{\partial_{i+1}} \mathfrak{R}^{m_i} \rightarrow M \rightarrow 0. \quad (51)$$

Tensoring (51) with $\mathbb{Z}[A]$, where $A = \mathbb{Z}^n / \Gamma$, we get the exact sequence

$$\mathbb{Z}[A]^{m_{i+1}} \xrightarrow{\partial_{i+1, \Gamma}} \mathbb{Z}[A]^{m_i} \rightarrow M \otimes \mathbb{Z}[A] \rightarrow 0,$$

from which it follows that

$$M \otimes \mathbb{Z}[A] = \operatorname{coker}(\partial_{i+1, \Gamma}). \quad (52)$$

The complex $\mathcal{C} \otimes \mathbb{Z}[A]$ is

$$\dots \rightarrow \mathbb{Z}[A]^{m_{i+1}} \xrightarrow{\partial_{i+1, \Gamma}} \mathbb{Z}[A]^{m_i} \xrightarrow{\partial_{i, \Gamma}} \dots. \quad (53)$$

Apply (49) to the chain complex (53), considered as complex over \mathbb{Z} , we get

$$\operatorname{tor}_{\mathbb{Z}}(H_i(\mathcal{C} \otimes \mathbb{Z}[A])) = \operatorname{tor}_{\mathbb{Z}}(\operatorname{coker} \partial_{i+1, \Gamma}),$$

which, with (52), gives

$$\operatorname{tor}_{\mathbb{Z}}(H_i(\mathcal{C} \otimes \mathbb{Z}[A])) = \operatorname{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[A]). \quad (54)$$

Theorem 3, with identity (54), gives

$$\limsup_{\langle \Gamma \rangle \rightarrow \infty} \frac{\log |\operatorname{tor}_{\mathbb{Z}}(H_i(\mathcal{C} \otimes \mathbb{Z}[A]))|}{|\mathbb{Z}^n / \Gamma|} = \mathbb{M}(\Delta(\operatorname{tor}(M))),$$

from which together with (50) we have

$$\limsup_{\langle \Gamma \rangle \rightarrow \infty} \frac{\log |\operatorname{tor}_{\mathbb{Z}}(H_i(\mathcal{C} \otimes \mathbb{Z}[A]))|}{|\mathbb{Z}^n / \Gamma|} = \mathbb{M}(\Delta(\operatorname{tor}(H_i(\mathcal{C})))),$$

which completes the proof of Theorem 4. \square

4. Homology of abelian covering

4.1. Alexander polynomials of links. Suppose Z is an oriented integral homology 3-sphere, i.e. $H_i(Z, \mathbb{Z}) \cong H_i(S^3, \mathbb{Z})$, and $L \subset Z$ is an oriented link with n ordered components. Let $N(L)$ be a small open tubular neighborhood of L and $X = Z \setminus N(L)$. By Alexander duality $H_1(X, \mathbb{Z}) \cong \mathbb{Z}^n$, and there is a natural identification of $H_1(X, \mathbb{Z})$ with \mathbb{Z}^n such that t_i corresponds to the meridian of the i -th component of the link. We fix such an identification of $H_1(X, \mathbb{Z})$ with \mathbb{Z}^n .

Let \tilde{X} be the abelian covering corresponding to the abelianization $\pi_1(X) \rightarrow H_1(X, \mathbb{Z}) = \mathbb{Z}^n$. The homology groups $H_i(\tilde{X}, \mathbb{Z})$ has a structure of $\mathfrak{R} = \mathbb{Z}[\mathbb{Z}^n]$ module. The Alexander polynomials $\Delta_i(L)$ (or $\Delta_i(L \subset Z)$), by definition, are the polynomials $\Delta_i(H_1(\tilde{X}, \mathbb{Z}))$. Recall that if j is the smallest index such that $\Delta_j(H_1(\tilde{X}, \mathbb{Z})) \neq 0$, then one defines $\Delta(H_1(\tilde{X}, \mathbb{Z})) = \Delta_j(H_1(\tilde{X}, \mathbb{Z}))$. We also define $\Delta(L) = \Delta(H_1(\tilde{X}, \mathbb{Z}))$.

Note that X has Euler characteristic 0. It is known that X is homotopic to a finite 2-dimensional CW-complex Y , with one 0-cell, $m + 1$ 1-cells a_1, \dots, a_{m+1} , and m

2-cells b_1, \dots, b_m , for some number m . Certainly $m \geq n$. Let $\rho: \pi_1(Y) \rightarrow H_1(Y, \mathbb{Z})$ be the standard abelianization map. By choosing an appropriate CW-structure, we can assume further that $\rho(a_i) = t_i$ for $i = 1, \dots, n$.

Let \tilde{Y} be the abelian covering of Y corresponding to the abelianization $\rho: \pi_1(Y) \rightarrow H_1(Y, \mathbb{Z}) = \mathbb{Z}^n$. The CW-complex of \tilde{Y} can be considered as a chain complex over $\mathfrak{R} = \mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$, and has the form

$$0 \rightarrow \mathfrak{R}^m \xrightarrow{\partial_2} \mathfrak{R}^{m+1} \xrightarrow{\partial_1} \mathfrak{R} \rightarrow 0. \quad (55)$$

Here

$$\partial_1 = \begin{pmatrix} 1 - \rho(a_1) \\ 1 - \rho(a_2) \\ \dots \\ 1 - \rho(a_{m+1}) \end{pmatrix}$$

and ∂_2 is an $m \times (m+1)$ -matrix with entries in \mathfrak{R} which can be calculated using Fox derivative. There is only one 0-cell of \tilde{Y} , denoted by O . The lift of a_i beginning at O will be denoted by \tilde{a}_i , $i = 1, \dots, m+1$.

Remark 4.1. The module $M_2 = \text{coker}(\partial_2)$ is known as the Alexander module. In some texts, e.g. [Hi], the Alexander polynomials are defined as $\Delta_i(M_2)$, which differ from ours only by a shift of index: $\Delta_i(H_1(\tilde{X}, \mathbb{Z})) = \Delta_{i+1}(M_2)$ since both $H_1(\tilde{X}, \mathbb{Z})$ and M_2 have the same \mathfrak{R} -torsion, see (49). In particular, $\Delta(L) = \Delta(H_1(\tilde{X}, \mathbb{Z})) = \Delta(M_2)$.

4.2. Homology of the branched covering. Suppose $\Gamma \subset \mathbb{Z}^n$ is a subgroup of finite index, and $A = A_\Gamma = \mathbb{Z}^n / \Gamma$. Let X_Γ and Y_Γ be the covering of X and Y respectively corresponding to the epimorphism $\pi_1 \rightarrow H_1 \rightarrow A$. Then the CW complex of Y_Γ is $\mathcal{C}(\tilde{Y}) \otimes_{\mathfrak{R}} \mathbb{Z}[A]$:

$$0 \rightarrow \mathbb{Z}[A]^m \xrightarrow{\partial_{2,\Gamma}} \mathbb{Z}[A]^{m+1} \xrightarrow{\partial_{1,\Gamma}} \mathbb{Z}[A] \rightarrow 0. \quad (56)$$

The branched covering X_Γ^{br} , by definition, is obtained from X_Γ by Dehn fillings as follows. The boundary of X is the union of n tori, each surrounding a link component. The boundary of X_Γ is also the union of several tori, each is the covering of one of tori in the boundary of X . Suppose T is a torus in the boundary of X_Γ covering the i -th torus of the boundary of X . There is a simple closed curve C on T covering the meridian of the i -th torus. To every boundary component T one does the Dehn filling on T that kills the homology class of C . The resulting 3-manifold is X_Γ^{br} .

The homology group $H_1(X_\Gamma^{\text{br}}, \mathbb{Z})$ is the quotient of $H_1(X_\Gamma, \mathbb{Z})$ by the relation $C = 0$, for all the curves C described in the above Dehn filling operation. The difficulty with working $H_1(X_\Gamma^{\text{br}}, \mathbb{Z})$ is these relations $C = 0$ are local, they cannot be obtained from a global relation in terms of \mathfrak{R} -modules.

We now describe a universal \mathfrak{R} -module through which the \mathbb{Z} -torsion of $H_1(X_\Gamma^{\text{br}}, \mathbb{Z})$ can be calculated.

Recall that ∂_2 in (55) is an $m \times m + 1$ matrix. Let $I_{n,m+1}$ be the $n \times (m + 1)$ matrix obtained from the identity $n \times n$ matrix by adding $(m + 1 - n)$ columns of 0, and T be the $n \times n$ diagonal matrix

$$T = \text{diag}(1 - t_1, \dots, 1 - t_n).$$

Proposition 4.1. *Let M be the \mathfrak{R} -module with the following presentation matrix*

$$\begin{pmatrix} \partial_2 & 0 \\ I_{n,m+1} & T \end{pmatrix},$$

which has size $(m + n) \times (m + 1 + n)$. Then for any subgroup $\Gamma \subset \mathbb{Z}^n$ of finite index we have

$$\text{tor}_{\mathbb{Z}}(H_1(X_\Gamma^{\text{br}}, \mathbb{Z})) \cong \text{tor}_{\mathbb{Z}}(M \otimes_{\mathfrak{R}} \mathbb{Z}[A_\Gamma]).$$

Proof. For $i = 1, \dots, n$ let $d_i = d_i(\Gamma)$ be the degree of t_i in $A = \mathbb{Z}^n/\Gamma$, and $u_i = \sum_{l=0}^{d_i-1} (t_i)^l \in \mathbb{Z}[A]$.

The homology of the branched covering $H_1(X_\Gamma^{\text{br}}, \mathbb{Z})$ is

$$H_1(X_\Gamma^{\text{br}}, \mathbb{Z}) = H_1(Y_\Gamma, \mathbb{Z})/(\text{Rel}),$$

where (Rel) is the $\mathbb{Z}[A]$ -submodule of $H_1(Y_\Gamma, \mathbb{Z})$ generated by $u_i \tilde{a}_i, i = 1, \dots, n$.

By definition, $H_1(Y_\Gamma, \mathbb{Z})$ is H_1 of the complex (56). By adding relations $u_i \tilde{a}_i = 0, i = 1, \dots, n$, we see that $H_1(X_\Gamma^{\text{br}}, \mathbb{Z})$ is H_1 of the following complex:

$$0 \rightarrow \mathbb{Z}[A]^m \times \mathbb{Z}[A]^n \xrightarrow{D_1} \mathbb{Z}[A]^{m+1} \xrightarrow{\partial_{1,\Gamma}} \mathbb{Z}[A] \rightarrow 0, \quad (57)$$

where $D_1(x, y) = \partial_{2,\Gamma}(x) + U'(y)$, with $U': \mathbb{Z}[A]^n \rightarrow \mathbb{Z}[A]^{m+1}$ being the $\mathbb{Z}[A]$ -linear map defined by

$$U'(x_1, \dots, x_n) = (u_1 x_1, \dots, u_n x_n, 0, \dots, 0).$$

Let $U: \mathbb{Z}[A]^n \rightarrow \mathbb{Z}[A]^n$ be the $\mathbb{Z}[A]$ -linear map given by the diagonal matrix $U = \text{diag}(u_1, \dots, u_n)$. Certainly $\ker U = \ker U'$.

Applying (49) to the chain complex (57), we get

$$\text{tor}_{\mathbb{Z}}(H_1(X_\Gamma^{\text{br}}, \mathbb{Z})) \cong \text{tor}_{\mathbb{Z}} \text{coker } D_1. \quad (58)$$

The map $U': \mathbb{Z}[A]^n \rightarrow \mathbb{Z}[A]^{m+1}$ descends to $U'': (\mathbb{Z}[A]^n / \ker U') \rightarrow \mathbb{Z}[A]^{m+1}$, hence $\text{coker } D_1 = \text{coker } D'_1$, where

$$D'_1: \mathbb{Z}[A]^m \times (\mathbb{Z}[A]^n / \ker U) \rightarrow \mathbb{Z}[A]^{m+1},$$

defined by $D'_1(x, y) = \partial_2(x) + U''(y)$. From (58) we have

$$\mathrm{tor}_{\mathbb{Z}}(H_1(X_{\Gamma}^{\mathrm{br}}, \mathbb{Z})) \cong \mathrm{tor}_{\mathbb{Z}}(\mathrm{coker} D'_1). \quad (59)$$

By tensoring $T: \mathfrak{R}^n \rightarrow \mathfrak{R}^n$ with $\mathbb{Z}[A]$, we get $T_{\Gamma}: \mathbb{Z}[A]^n \rightarrow \mathbb{Z}[A]^n$, which is given by a diagonal matrix. Note that $T_{\Gamma}U = 0$, i.e. T_{Γ} is 0 on the image of U , hence T_{Γ} descend to a map $T'_{\Gamma}: (\mathbb{Z}[A]^n / \mathrm{Im} U) \rightarrow \mathbb{Z}[A]^n$.

We have the following commutative diagram with exact vertical lines:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z}[A]^{m+1} \times (\mathbb{Z}[A]^n / \ker U) & \xrightarrow{D'_1} & \mathbb{Z}[A]^{m+1} & \longrightarrow & 0 \\ \downarrow & & \downarrow i_1 & & \downarrow i_2 & & \downarrow \\ 0 & \longrightarrow & \mathbb{Z}[A]^{m+1} \times \mathbb{Z}[A]^n & \xrightarrow{D_{\Gamma}} & \mathbb{Z}[A]^{m+1} \times \mathbb{Z}[A]^n & \longrightarrow & 0 \\ \downarrow & & \downarrow j_1 & & \downarrow j_2 & & \downarrow \\ 0 & \longrightarrow & (\mathbb{Z}[A]^n / \mathrm{Im} U) & \xrightarrow{T'_{\Gamma}} & \mathbb{Z}[A]^n & \longrightarrow & 0, \end{array}$$

where $i_1(x, y) = (x, U(y))$, $i_2(x) = (x, 0)$, $j_1(x, y) = (0, y)$, $j_2(x, y) = y$, and D_{Γ} is the matrix of presentation of M , tensoring with $\mathbb{Z}[A]$.

Let the first complex be \mathcal{D}_1 , the second \mathcal{D}_2 , and the third \mathcal{D}_3 . From the exact sequence $0 \rightarrow \mathcal{D}_1 \rightarrow \mathcal{D}_2 \rightarrow \mathcal{D}_3 \rightarrow 0$ we have a long exact sequence

$$\cdots \rightarrow H_1(\mathcal{D}_3) \rightarrow H_0(\mathcal{D}_1) \rightarrow H_0(\mathcal{D}_2) \rightarrow H_0(\mathcal{D}_3) \rightarrow 0. \quad (60)$$

The first term is 0 and the last term is a free abelian group, by Lemma 4.2 below. Hence the second term and the third term in (60) have the same \mathbb{Z} -torsion. Since $H_0(\mathcal{D}_1) = \mathrm{coker} D'_1$ and $H_0(\mathcal{D}_2) = M \otimes \mathbb{Z}[A]$, we have

$$\mathrm{tor}_{\mathbb{Z}}(\mathrm{coker} D'_1) = \mathrm{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[A]),$$

which, together with (59), proves the proposition. \square

Lemma 4.2. *For chain complex \mathcal{D}_3 ,*

$$0 \rightarrow (\mathbb{Z}[A]^n / \mathrm{Im} U) \xrightarrow{T'_{\Gamma}} \mathbb{Z}[A]^n \rightarrow 0,$$

one has $H_1(\mathcal{D}) = 0$ and $H_0(\mathcal{D})$ is a free abelian group.

Proof. This is the same as to show that for the chain \mathcal{D}' ,

$$0 \rightarrow \mathbb{Z}[A]^n \xrightarrow{U} \mathbb{Z}[A]^n \xrightarrow{T_{\Gamma}} \mathbb{Z}[A]^n \rightarrow 0,$$

one has $H_1(\mathcal{D}') = 0$, and $H_0(\mathcal{D}')$ is free abelian.

Since both U and T_Γ are diagonal, $\mathcal{D}' = \bigoplus_{i=1}^n \mathcal{D}'_i$, where \mathcal{D}'_i is the complex

$$0 \rightarrow \mathbb{Z}[A] \xrightarrow{u_i} \mathbb{Z}[A]^n \xrightarrow{1-t_i} \mathbb{Z}[A] \rightarrow 0.$$

As seen in Section 1.7, the principal ideals (u_i) and $(1 - t_i)$ are annihilator of each other, hence $H_1(\mathcal{D}'_i) = 0$. Besides, the ideal $(1 - t_i)$ is primitive as a lattice in $\mathbb{Z}[A]$, hence $H_0(\mathcal{D}'_i)$ is a free abelian group. \square

4.3. Proof of Theorem 2

Proof. Part (a), the case of non-branched covering, follows immediately from Theorem 5.

Let us consider the case of branched covering. By Proposition 4.1, we have

$$\begin{aligned} \limsup_{\langle \Gamma \rangle \rightarrow \infty} \frac{\log |\operatorname{tor}_{\mathbb{Z}}(H_1(X_\Gamma^{\text{br}}, \mathbb{Z}))|}{|\mathbb{Z}^n / \Gamma|} &= \limsup_{\langle \Gamma \rangle \rightarrow \infty} \frac{\log \operatorname{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[A_\Gamma])}{|\mathbb{Z}^n / \Gamma|} \\ &= \mathbb{M}(\Delta(M)) \quad \text{by Theorem 3.} \end{aligned} \quad (61)$$

The module $M_1 = \mathfrak{R}^n / T(\mathfrak{R}^n)$ has a free resolution

$$0 \rightarrow \mathfrak{R}^n \xrightarrow{T} \mathfrak{R}^n \rightarrow M_1 \rightarrow 0,$$

hence its projective dimension is 1. Also $\Delta_0(M_1) = \prod_{i=1}^n (1 - t_i)$.

Let $M_2 = \operatorname{coker} \partial_2$. From the matrix of presentation of M we see that there is an exact sequence

$$0 \rightarrow M_2 \rightarrow M \rightarrow M_1 \rightarrow 0.$$

Since the projective dimension of M_1 is ≤ 1 , by Theorem 3.12 of [Hi], one has

$$\begin{aligned} \Delta_j(M) &= \Delta_j(M_2) \Delta_0(M_1) \\ &= \Delta_j(M_2) \prod_{i=1}^n (1 - t_i). \end{aligned}$$

It follows that $\Delta(M) = \Delta(M_2) \prod_{i=1}^n (1 - t_i)$. Since $\mathbb{M}(1 - t_i) = 1$ and $\Delta(M_2) = \Delta(L)$, we have

$$\mathbb{M}(\Delta(M)) = \mathbb{M}(\Delta(L)),$$

from which together with (61) one gets part (b) of Theorem 2. \square

5. Converging sequences

5.1. Statement. For a non-zero vector $x \in \mathbb{R}^n$ let $\mathbf{r}(x) = x/\|x\| \in S^{n-1}$ be the unit vector positively colinear with x . Here S^{n-1} is the $(n-1)$ -dimensional sphere of unit vectors in \mathbb{R}^n . For a subgroup $\Gamma \subset \mathbb{Z}^n$ of finite index let $d_i = d_i(\Gamma)$ be the degree of t_i in the quotient group $A_\Gamma = \mathbb{Z}^n/\Gamma$. Let $\mathbf{r}(\Gamma) = \mathbf{r}(d_1, \dots, d_n) \in S_+^{n-1}$, the part of S^{n-1} with non-negative coordinates.

Theorem 7. Suppose M is a finitely generated \mathfrak{R} -module. For any $\kappa \in S_+^{n-1}$, there exists a sequence of finite index subgroups $\Gamma_s \subset \mathbb{Z}^n$, $s = 1, 2, \dots$, such that

$$\lim_{s \rightarrow \infty} \mathbf{r}(\Gamma_s) = \kappa$$

and

$$\limsup_{s \rightarrow \infty} \frac{\log |\mathrm{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[\mathbb{Z}^n/\Gamma_s])|}{|\mathbb{Z}^n/\Gamma_s|} = \mathbb{M}(\Delta(M)).$$

Remark 5.1. One could prove a similar statement, replacing $M \otimes \mathbb{Z}[\mathbb{Z}^n/\Gamma_s]$ with $H_i(\mathcal{C} \otimes \mathbb{Z}[\mathbb{Z}^n/\Gamma_s])$ like in Theorem 4.

The proof and methods of this section are independent of Theorem 3. It gives an alternative proof of “half” of Theorem 3: The left-hand side in the identity of Theorem 3 is greater than or equal to the right-hand side.

5.2. A result of Bombieri and Zannier: reduction from \mathbb{Z}^n to \mathbb{Z} . For $\mathbf{k} \in \mathbb{Z}^n$ let $\mathbf{k}^\perp = \{\mathbf{m} \in \mathbb{Z}^n \mid \mathbf{k} \cdot \mathbf{m} = 0\}$, where $\mathbf{k} \cdot \mathbf{m}$ is the usual dot product. Define

$$\langle \mathbf{k} \rangle = \langle \mathbf{k}^\perp \rangle = \min\{|\mathbf{x}|, \mathbf{x} \in \mathbf{k}^\perp \setminus \{0\}\}.$$

The group homomorphism $\mathbb{Z}^n \rightarrow \mathbb{Z}$ given by $\mathbf{m} \rightarrow \mathbf{m} \cdot \mathbf{k}$ gives rise to the algebra homomorphism $\tau_{\mathbf{k}}: \mathbb{Q}[t_1^{\pm 1}, \dots, t_n^{\pm 1}] \rightarrow \mathbb{Q}[t^{\pm 1}]$ defined by

$$\tau_{\mathbf{k}}(t^{\mathbf{m}}) = t^{\mathbf{m} \cdot \mathbf{k}}.$$

The following is a deep result of Bombieri and Zannier [SZ], [BMZ], which was formulated as a conjecture by Schinzel.

Theorem 5.1. Suppose $p_1, p_2 \in \mathbb{Q}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ are co-prime. There is a constant $C = C(p_1, p_2)$ such that if $\langle \mathbf{k} \rangle > C$, then $\gcd(\tau_{\mathbf{k}}(p_1), \tau_{\mathbf{k}}(p_2))$ is the product of some (possibly none) cyclotomic polynomials.

From this one can easily deduce the following.

Proposition 5.2. *Suppose $p_1, \dots, p_\ell \in \mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$. There exists a constant $C = C(p_1, \dots, p_\ell)$ such that if $\langle k \rangle > C$, then*

$$\gcd(\tau_k(p_1), \dots, \tau_k(p_\ell)) = \phi \tau_k(\gcd(p_1, \dots, p_\ell)),$$

where ϕ is a product of cyclotomic polynomials.

Proof. By dividing each of the p_i 's by $\gcd(p_1, \dots, p_\ell)$ we can henceforth assume that $\gcd(p_1, \dots, p_\ell) = 1$.

We will use induction on ℓ . But first make the following well-known observation on the coefficients of $\tau_k(p)$ of a polynomial $p \in \mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$ having the form

$$p = \sum_{m \in \mathcal{N}} c_m t^m,$$

where $\mathcal{N} \subset \mathbb{Z}$ is a finite set. Then

$$\tau_k(p) = \sum_{m \in \mathcal{N}} c_m t^{m \cdot k}. \quad (62)$$

If $\langle k \rangle$ is greater than $|m - m'|$ for any two $m \neq m' \in \mathcal{N}$, then $k \cdot m \neq k \cdot m'$, and (62) shows that the coefficients of $\tau_k(p)$, in some order, are exactly the coefficients of p .

Now we proceed with induction. Suppose $\ell = 2$. By Bombieri and Zannier result, over $\mathbb{Q}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$, $\gcd(\tau_k(p_1), \tau_k(p_2)) = \phi$, a product of cyclotomic polynomials. Hence over $\mathbb{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$, $\gcd(p_1, p_2) = a\phi$, for some integer a . It follows that a is the gcd of all the coefficients of $\tau_k(p_1)$ and $\tau_k(p_2)$. By the above observation, with $\langle k \rangle$ big enough, this means a is the gcd of all the coefficients of p_1 and p_2 , which must be 1. This proves the statement when $\ell = 2$.

Now assume $\ell \geq 3$. One has

$$\begin{aligned} \gcd(\tau_k(p_1), \dots, \tau_k(p_\ell)) &= \gcd(\tau_k(p_1), \gcd(\tau_k(p_2), \dots, \tau_k(p_\ell))) \\ &= \gcd(\tau_k(p_1), \phi \tau_k(\gcd(p_2, \dots, p_\ell))) \quad \text{by induction} \\ &= \phi' \tau_k(\gcd(p_1, \gcd(p_2, \dots, p_\ell))) \quad \text{by case } \ell = 2 \\ &= \phi' \tau_k(\gcd(p_1, p_2, \dots, p_\ell)). \end{aligned}$$

Here ϕ, ϕ' are product of cyclotomic polynomials. The proof is completed. \square

5.3. A result of Lawton. Recall that the additive Mahler measure $\mathbb{M}(f)$ of $f \in \mathbb{C}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$, $f \neq 0$, is defined by

$$\mathbb{M}(f) = \int_{\mathbb{S}^n} \log |f(x)| d\sigma,$$

where \mathbb{S}^n is the n -dimensional torus, and $d\sigma$ is the invariant Haar measure on \mathbb{S}^n normalized so that $\int_{\mathbb{S}^n} d\sigma = 1$.

The Mahler measure is additive, $\mathbb{M}(fg) = \mathbb{M}(f) + \mathbb{M}(g)$. It is known that $\mathbb{M}(f) = 0$ if and only if f is a generalized cyclotomic polynomial, see e.g. [SZ], [Sch].

The following approximation result was a conjecture of D. Boyd, and was proved by Lawton, see e.g. [Law], [SZ], [Sch].

Theorem 5.3 (Lawton). *Suppose $f \in \mathbb{C}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$, $f \neq 0$. Then*

$$\lim_{\langle \mathbf{k} \rangle \rightarrow \infty} \mathbb{M}(\tau_{\mathbf{k}}(f)) = \mathbb{M}(f).$$

5.4. A converging sequence. The following follows from Bombieri–Zannier and Lawton results. Denote $\mathfrak{R}_1 = \mathbb{Z}[t^{\pm 1}]$.

Proposition 5.4. *Suppose that $\mathbf{k}^{(s)} \in \mathbb{Z}^n$, $s = 1, 2, \dots$, is any sequence such that $\lim_{s \rightarrow \infty} \langle \mathbf{k}^{(s)} \rangle = \infty$, and let M be finitely generated \mathfrak{R} -module. Let $M^{(s)} = M \otimes \mathfrak{R}_1$, where \mathfrak{R}_1 is considered as an \mathfrak{R} -module via $\tau_s := \tau_{\mathbf{k}^{(s)}} : \mathfrak{R} \rightarrow \mathfrak{R}_1$. Then*

$$\lim_{s \rightarrow \infty} \mathbb{M}(\Delta(M^{(s)})) = \mathbb{M}(\Delta(M)).$$

Remark 5.2. It is not true in general that $\lim_{s \rightarrow \infty} \Delta(M^{(s)}) = \Delta(M)$.

Proof. Suppose M has a presentation matrix ∂ of size $m_1 \times m_0$, with entries in \mathfrak{R} . Then $M^{(s)}$ has presentation matrix $\tau_s(\partial)$, with entries in \mathfrak{R}_1 .

Let j be the smallest integer such that $\Delta_j(M) \neq 0$. This means all the $(m_0 - j')$ -minors of ∂ are 0 if $j' < j$, and if the $(m_0 - j)$ -minors of ∂ are $p_1, \dots, p_\ell \in \mathfrak{R}$, then

$$\Delta(M) = \Delta_j(M) = \gcd(p_1, \dots, p_\ell).$$

Note that every minor of $\tau_s(\partial)$ is obtained from the corresponding minor by applying τ_s . It follows that all the $(m_0 - j')$ -minors of $\tau_s(\partial)$ are 0 if $j' < j$, and the $(m_0 - j)$ -minors are $\tau_s(p_1), \dots, \tau_s(p_\ell)$.

By Proposition 5.2, for s big enough,

$$\gcd(\tau_s(f_1), \dots, \tau_s(f_r)) = \phi \tau_s(\gcd(f_1, \dots, f_r)),$$

where ϕ is a product of cyclotomic polynomials. This means

$$\Delta(M^{(s)}) = \phi \tau_s(\Delta(M)).$$

Using additivity of the Mahler measure and the fact that the Mahler measure of a cyclotomic polynomial is 0, we have

$$\mathbb{M}(\Delta(M^{(s)})) = \mathbb{M}(\tau_s(\Delta(M))).$$

Since $\langle k^{(s)} \rangle \rightarrow \infty$ as $s \rightarrow \infty$, by Lawton's theorem, we have

$$\lim_{s \rightarrow \infty} \mathbb{M}(\Delta(M^{(s)})) = \lim_{s \rightarrow \infty} \mathbb{M}_{\tau_s}(\Delta(M)) = \mathbb{M}(\Delta(M)). \quad \square$$

5.5. Theorem 3, the case $n = 1$. In the previous section we approximate \mathbb{Z}^n by $\mathbb{Z}^n/(k^\perp)$, which has rank 1. Now we want to approximate abelian rank-1 group by finite cyclic group. Here we give a short proof, independent of previous sections, of the case $n = 1$ of Theorem 3.

Proposition 5.5. *Suppose M is a finitely generated \mathfrak{R}_1 -module. Then*

$$\lim_{\ell \rightarrow \infty} \frac{\text{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[\mathbb{Z}/\ell])}{\ell} = \mathbb{M}(\Delta(\text{tor} M)).$$

Proof. The reason the case $n = 1$ is easy is that $\mathbb{Z}[\mathbb{Z}/\ell] = \mathfrak{R}_1/(1-t^\ell)$, with $(1-t^\ell)$ a principal ideal.

For an \mathfrak{R}_1 -module N and an element $a \in \mathfrak{R}_1$, let ${}_a N$ be the a -torsion of N :

$${}_a N = \{x \in N \mid ax = 0\}.$$

A homological interpretation of ${}_a N$ is $\text{Tor}_1^{\mathfrak{R}_1}(N, \mathfrak{R}_1/(a)) = {}_a N$. If $a|b$ then ${}_a N \subset {}_b N$. If N is a finitely generated torsion module, then there is $b \in \mathfrak{R}_1$, called a *universal annihilator* of N , such that for every $a \in \mathfrak{R}_1$,

$${}_a N = {}_{\gcd(b,a)} N.$$

For example, such b can be the product of all the generators of all prime ideals associated to N . One can also define b as the least common multiple of the family of annihilators of a generating family for N .

Since $M' = M/\text{tor}(M)$ is torsion-free, by [Bou], Chapter VII, there is a free \mathfrak{R}_1 -module F such that $M' \subset F$ and F/M' is a torsion module. Let f be a universal annihilator of F/M' . Decompose $f = f_1 f_2$, where f_1 is the product of all cyclotomic factors (with multiplicity) in the prime decomposition of f . The identity map $F \rightarrow F$ descends to a surjective map

$$F/(f_1 F) \twoheadrightarrow {}_{f_1}(F/M').$$

Since f_1 is monic, $F/f_1 F$ is a finitely generated \mathbb{Z} -module. It follows that ${}_{f_1}(F/M')$ is a finitely generated \mathbb{Z} -module, hence its \mathbb{Z} -torsion part is a finite set.

Tensoring the exact sequence

$$0 \rightarrow M' \rightarrow F \rightarrow F/M' \rightarrow 0$$

with $\mathbb{Z}[\mathbb{Z}/\ell]$, we get

$$\begin{aligned} 0 \rightarrow \mathrm{Tor}_1^{\mathfrak{R}_1}(F/M', \mathbb{Z}[\mathbb{Z}/\ell]) &\rightarrow M' \otimes \mathbb{Z}[\mathbb{Z}/\ell] \\ &\rightarrow F \otimes \mathbb{Z}[\mathbb{Z}/\ell] \rightarrow (F/M') \otimes \mathbb{Z}[\mathbb{Z}/\ell] \rightarrow 0. \end{aligned}$$

Since F is a free \mathfrak{R}_1 -module, the third term is a free \mathbb{Z} -module. It follows that the \mathbb{Z} -torsions of the first and the second terms are the same

$$\mathrm{tor}_{\mathbb{Z}}(\mathrm{Tor}_1^{\mathfrak{R}_1}(F/M', \mathbb{Z}[\mathbb{Z}/\ell])) = \mathrm{tor}_{\mathbb{Z}}(M' \otimes \mathbb{Z}[\mathbb{Z}/\ell]).$$

Note that

$$\mathrm{Tor}_1^{\mathfrak{R}_1}(F/M', \mathbb{Z}[\mathbb{Z}/\ell]) = {}_{(1-t^\ell)}(F/M')$$

is a subset of ${}_f(F/M')$ since $1 - t^\ell$ is a product of cyclotomic polynomials. Since $|\mathrm{tor}_{\mathbb{Z}}({}_f(F/M'))|$ is finite and does not depend on ℓ , we conclude that $\mathrm{tor}_{\mathbb{Z}}(M' \otimes \mathbb{Z}[\mathbb{Z}/\ell])$ is finite and bounded from above.

Tensoring the exact sequence

$$0 \rightarrow \mathrm{tor}(M) \rightarrow M \rightarrow M' \rightarrow 0$$

with $\mathbb{Z}[\mathbb{Z}/\ell] = \mathfrak{R}_1/(1 - t^\ell)$, taking into account $\mathrm{Tor}_1^{\mathfrak{R}_1}(M', \mathfrak{R}_1/(1 - t^\ell)) = 0$, we have

$$0 \rightarrow \mathrm{tor}(M) \otimes \mathbb{Z}[\mathbb{Z}/\ell] \rightarrow M \otimes \mathbb{Z}[\mathbb{Z}/\ell] \rightarrow M' \otimes \mathbb{Z}[\mathbb{Z}/\ell] \rightarrow 0.$$

Since the \mathbb{Z} -torsion of the last term is bounded, we see that the \mathbb{Z} -torsion parts of the first two terms have the same growth, i.e. $M \sim \mathrm{tor}(M)$. The proposition now follows from the case of torsion modules, which was known [Sch] (see also [GS], [Ri], [Lü2]). \square

5.6. Converging sequences

Lemma 5.6. *Let M be a finitely generated \mathfrak{R} -module, and $\mathbf{k}^{(s)} \in \mathbb{Z}^n$, $s = 1, 2, \dots$, be any sequence such that $\lim_{s \rightarrow \infty} \langle \mathbf{k}^{(s)} \rangle = \infty$ and $\gcd(k_1^{(s)}, \dots, k_n^{(s)}) = 1$ for $s \geq 1$. For each positive integer j define the subgroup $\Gamma_{s,j} \subset \mathbb{Z}^n$ by*

$$\Gamma_{s,j} = (\mathbf{k}^{(s)})^\perp + j \mathbf{k}^{(s)}.$$

For every s there exists an integer $\eta_s > 0$ such that for every $j_s > \eta_s$, we have

$$\lim_{s \rightarrow \infty} \frac{\log |\mathrm{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[A_{\Gamma_{s,j_s}}])|}{|\mathbb{Z}^n / \Gamma_{s,j_s}|} = \mathbb{M}(\Delta(M)).$$

Proof. It is easy to see that for any $\mathbf{k} = (k_1, \dots, k_n) \in \mathbb{Z}^n$ with $\gcd(k_1, \dots, k_n) = 1$, and $0 < j \in \mathbb{Z}$, the map $\mathbf{m} \rightarrow \mathbf{m} \cdot \mathbf{k} \pmod{j}$ is an isomorphism between $\mathbb{Z}^n / (\mathbf{k}^\perp \oplus j\mathbf{k})$ and $\mathbb{Z} / (j|\mathbf{k}|^2)$.

It follows that, as \mathfrak{R} -modules,

$$M \otimes_{\mathfrak{R}} \mathbb{Z}[\mathbb{Z}^n / \Gamma_{s,j}] \cong M^{(s)} \otimes_{\mathfrak{R}_1} \mathbb{Z}[\mathbb{Z} / (j|\mathbf{k}^{(s)}|^2)].$$

Since $|\mathbb{Z}^n / \Gamma_{s,j}| = j|\mathbf{k}^{(s)}|^2$, one has

$$\begin{aligned} \lim_{j \rightarrow \infty} \frac{\log |\operatorname{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[A_{\Gamma_{s,j}}])|}{|\mathbb{Z}^n / \Gamma_{s,j}|} &= \lim_{j \rightarrow \infty} \frac{|\operatorname{tor}_{\mathbb{Z}}(M^{(s)} \otimes_{\mathfrak{R}_1} \mathbb{Z}[\mathbb{Z} / (j|\mathbf{k}^{(s)}|^2)])|}{j|\mathbf{k}^{(s)}|^2}, \\ \lim_{j \rightarrow \infty} \frac{\log |\operatorname{tor}_{\mathbb{Z}}(M \otimes \mathbb{Z}[A_{\Gamma_{s,j}}])|}{|\mathbb{Z}^n / \Gamma_{s,j}|} &= \mathbb{M}(\Delta(M^{(s)})) \quad \text{by Proposition 5.5.} \end{aligned} \quad (63)$$

Let $a_{s,j}$ be the left-hand side of (63). From (63), for fixed s , there is $\eta_s > 0$ such that if $j > \eta_s$, then

$$|a_{s,j} - \mathbb{M}(\Delta(M^{(s)}))| < 1/s.$$

It is clear that if $j_s > \eta_s$, then

$$\lim_{s \rightarrow \infty} a_{s,j_s} = \lim_{s \rightarrow \infty} \mathbb{M}(\Delta(M^{(s)})) = \mathbb{M}(\Delta(M)),$$

where the last identity follows from Proposition 5.4. \square

5.7. Proof of Theorem 7. Assume the sequence $\mathbf{k}^{(s)}$ of Lemma 5.6 satisfies $k_i^{(s)} \neq 0$ for $i = 1, \dots, n$. If we choose j_s divisible by the product $k_1^{(s)} \dots k_n^{(s)}$, then $d_i(\Gamma_{s,j_s}) = |j_s/k_i^{(s)}|$, and

$$\mathbf{r}(\Gamma_{s,j_s}) = \mathbf{r}(1/|k_1^{(s)}|, \dots, 1/|k_n^{(s)}|).$$

Thus Theorem 7 follows from Lemma 5.6 and the following result.

Lemma 5.7. Suppose $\kappa \in S_+^{n-1}$. There exists $\mathbf{k}^{(s)} = (k_1^{(s)}, \dots, k_n^{(s)}) \in \mathbb{Z}^n$ such that

- (i) $k_1^{(s)}, \dots, k_n^{(s)} > 0$ and have greatest common divisor 1,
- (ii) $\lim_{s \rightarrow \infty} \mathbf{r}(1/k_1^{(s)}, \dots, 1/k_n^{(s)}) = \kappa$,
- (iii) $\lim_{s \rightarrow \infty} \langle \mathbf{k}^{(s)} \rangle = \infty$.

Proof. Let S_{++}^{n-1} be the subset of S^{n-1} consisting of points with all positive coordinates. Let $\operatorname{Inv}: S_{++}^{n-1} \rightarrow S_{++}^{n-1}$ be the map defined by

$$\operatorname{Inv}(x_1, \dots, x_n) = \mathbf{r}(1/x_1, \dots, 1/x_n).$$

It is clear that Inv is an involution, and hence is a auto-diffeomorphism of S_{++}^{n-1} .

The set Q of all points of the form $\mathbf{r}(k_1, \dots, k_n)$, with k_1, \dots, k_n positive and co-prime, is dense in S_+^{n-1} . If \mathcal{L} is a finite collection of hyperplanes in \mathbb{R}^n , then $Q \setminus \mathcal{L}$ is still dense in S_+^{n-1} .

For $s \geq 1$ let P_s be the set of all points in \mathbb{Z}^n having norm $\leq s$, and

$$\mathcal{L}_s = \bigcup_{s \in P_s} \mathbf{k}^\perp.$$

By definition, if $\mathbf{k} \notin \mathcal{L}_s$, then $\langle \mathbf{k} \rangle > s$.

$Q \setminus \mathcal{L}_s$ is dense in S_+^{n-1} , hence so is $\text{Inv}(Q \setminus \mathcal{L}_s)$. This implies there is $\mathbf{x}^{(s)} \in Q \setminus \mathcal{L}_s$ such that

$$\|\text{Inv}(\mathbf{k}^{(s)}) - \kappa\| < 1/s. \quad (64)$$

By definition, $\mathbf{x}^{(s)} = \mathbf{r}(\mathbf{k}^{(s)})$ for some $\mathbf{k}^{(s)} = (k_1^{(s)}, \dots, k_n^{(s)})$ with positive and co-prime $k_i^{(s)}$. Since $\mathbf{k}^{(s)} \notin \mathcal{L}_s$, we have $\langle \mathbf{k}^{(s)} \rangle > s$, which establishes Property (ii). Inequality (64) shows that Property (iii) also holds. \square

References

- [AM] M. Atiyah and I. Macdonald, *Introduction to commutative algebra*. Addison-Wesley Publishing Co., Reading, Mass., 1969. [Zbl 0175.03601](#) [MR 0242802](#)
- [BM] H. Bass and M. P. Murthy, Grothendieck groups and Picard groups of abelian group rings. *Ann. of Math.* (2) **86** (1967), 16–73. [Zbl 0157.08202](#) [MR 0219592](#)
- [BV] N. Bergeron and A. Venkatesh, The asymptotic growth of torsion homology for arithmetic groups. *J. Inst. Math. Jussieu* **12** (2013), no. 2, 391–447. [Zbl 1266.22013](#) [MR 3028790](#)
- [Ber] D. Bertrand, Duality on tori and multiplicative dependence relations. *J. Austral. Math. Soc. Ser. A* **62** (1997), 198–216. [Zbl 0886.11035](#) [MR 1433209](#)
- [BMZ] E. Bombieri, D. Masser, and U. Zannier, *Anomalous subvarieties structure theorems and applications*, *Internat. Math. Res. Notices IMRN* **2007** (2007), no. 19, Art. ID rnm057, 33 pp. [Zbl 1145.11049](#) [MR 2359537](#)
- [Bou] N. Bourbaki, *Commutative algebra*. Chapters 1–7, Elements of Mathematics (Berlin), Springer-Verlag, Berlin 1989. [Zbl 0902.13001](#) [MR 1727221](#)
- [EW] M. Einsiedler and T. Ward, Fitting ideals for finitely presented algebraic dynamical systems. *Aequationes Math.* **60** (2000), 57–71. [Zbl 0972.22005](#) [MR 1777892](#)
- [GS] F. Gonzalez-Acuna and H. Short, Cyclic branched coverings of knots and homology spheres. *Rev. Mat. Univ. Complut. Madrid* **4** (1991), no. 1, 97–120. [Zbl 0756.57001](#) [MR 1142552](#)
- [Go] C. McA. Gordon, Knots whose branched coverings have periodic homology. *Trans. Amer. Math. Soc.* **168** (1972), 357–370. [Zbl 0238.55001](#) [MR 0295327](#)
- [Hi] J. Hillman, *Algebraic invariants of links*. Series on Knots and Everything 32, World Scientific Publishing, River Edge, NJ, 2002. [Zbl 1007.57001](#) [MR 1932169](#)

- [HS] J. Hillman and M. Sakuma On the homology of finite abelian coverings of links. *Canad. Math. Bull.* **40** (1997), 309–315. [Zbl 0888.57005](#) [MR 1464839](#)
- [Ka] I. Kaplansky, *Commutative rings*. The University of Chicago Press, Chicago, Ill., 1974. [Zbl 0296.13001](#) [MR 0345945](#)
- [Lau] M. Laurent, Equations diophantiennes exponentielles. *Invent. Math.* **78** (1984), 299–327. [Zbl 0554.10009](#) [MR 0767195](#)
- [Law] W. Lawton, A problem of Boyd concerning geometric means of polynomials. *J. Number Theory* **16** (1983), 356–362. [Zbl 0516.12018](#) [MR 0707608](#)
- [LSW] D. Lind, K. Schmidt, and T. Ward, Mahler measure and entropy for commuting automorphisms of compact groups. *Invent. Math.* **101** (1990), 593–629. [Zbl 0774.22002](#) [MR 1062797](#)
- [Le1] T. Le, Hyperbolic volume, Mahler measure, and homology growth. Talk at Columbia University 2009, slides available at <http://www.math.columbia.edu/textasciitilde volconf09/notes/leconf.pdf>.
- [Le2] T. Le, Hyperbolic volume and homology growth. In preparation.
- [Lü1] W. Lück, Approximating L^2 -invariants by their finite-dimensional analogues. *Geom. Funct. Anal.* **4** (1994), 455–481. [Zbl 0853.57021](#) [MR 1280122](#)
- [Lü2] W. Lück, L^2 -invariants: theory and applications to geometry and KK-theory. *Ergeb. Math. Grenzgeb.* (3) 44, Springer-Verlag, Berlin 2002. [Zbl 1009.55001](#) [MR 1926649](#)
- [Mü] W. Müller, Analytic torsion and cohomology of hyperbolic 3-manifolds. Preprint of the Max-Planck Institute, Bonn 2009.
- [MM] J. Mayberry and K. Murasugi, Torsion-groups of abelian coverings of links. *Trans. Amer. Math. Soc.* **271** (1982), 143–173. [Zbl 0487.57001](#) [MR 0648083](#)
- [Ra] J. Raimbault, Exponential growth of torsion in abelian coverings. *Algebr. Geom. Topol.* **12** (2012), no. 3, 1331–1372. [Zbl 1250.57004](#) [MR 2966689](#)
- [Ri] R. Riley, Growth of order of homology of cyclic branched covers of knots. *Bull. London Math. Soc.* **22** (1990), 287–297. [Zbl 0727.57002](#) [MR 1041145](#)
- [SZ] A. Schinzel, *Polynomials with special regard to reducibility*. With an appendix by Umberto Zannier, *Encyclopedia Math. Applications* 77, Cambridge University Press, Cambridge 2000. [Zbl 0956.12001](#) [MR 1770638](#)
- [Sch] K. Schmidt, *Dynamical systems of algebraic origin*. *Progr. Math.* 128, Birkhäuser Verlag, Basel 1995. [Zbl 0833.28001](#) [MR 1345152](#)
- [Sch-W] W. Schmidt, Heights of points on subvarieties of G_m^n . In *Number theory* (Tiruchirappalli, 1996), *Contemp. Math.* 210, Amer. Math. Soc., Providence, RI, 1998, 97–99. [Zbl 0897.11021](#) [MR 1478487](#)
- [SW] D. Silver and S. Williams, Mahler measure, links and homology growth. *Topology* **41** (2002), 979–991. [Zbl 1024.57007](#) [MR 1923995](#)
- [Tu] V. Turaev, *Introduction to combinatorial torsions*. *Lectures Math.* ETH Zurich, Birkhäuser, Basel 2001. [Zbl 0970.57001](#) [MR 1809561](#)

Received November 14, 2012

Thang Le, Department of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332–0160, U.S.A.

E-mail: letu@math.gatech.edu