Zeitschrift:	Commentarii Mathematici Helvetici
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	89 (2014)
Artikel:	Ricci flow on three-dimensional manifolds with symmetry
Autor:	Lott, John / Sesum, Natasa
DOI:	https://doi.org/10.5169/seals-515664

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 04.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Ricci flow on three-dimensional manifolds with symmetry

John Lott and Natasa Sesum*

Abstract. We describe the Ricci flow on two classes of compact three-dimensional manifolds: 1. Warped products with a circle fiber over a two-dimensional base. 2. Manifolds with a free local isometric T^2 -action.

Mathematics Subject Classification (2010). 53C44,57M50.

Keywords. Ricci flow, warped product, geometrization.

1. Introduction

In understanding three-dimensional Ricci flow solutions, a special role is played by three-manifolds with symmetry. It is plausible that one can get more precise results about such Ricci flows than in the general case. Consequently, such Ricci flows with symmetry have been studied for quite a while. We begin by describing some of the earlier results.

Locally homogeneous three-dimensional Ricci flow solutions were examined by Isenberg–Jackson [IJ92] and Knopf–McLeod [KM01]. The flow equations reduced to a system of three coupled ODEs. The solutions are now fairly well understood; see, for example, [Lot07, Section 3].

Certain three-dimensional Ricci flow solutions with a two-dimensional isometry group were analyzed by Carfora–Isenberg–Jackson [CIJ90] and Hamilton [Ham95, Section 11]. They considered Riemannian three-manifolds that admit a free isometric T^2 -action. The base is a circle and the total space is (necessarily) diffeomorphic to a 3-torus. Under some additional assumptions (\mathcal{F} -metric [CIJ90] or square torus metric [Ham95, Section 11]) it was shown that the Ricci flow exists for all time and converges to a flat metric.

Hamilton and Isenberg considered a twisted version of such torus bundles [HI93]. That is, there was a local (with respect to the base) free isometric T^2 -action. The T^2 -bundle over the circle was globally twisted by a hyperbolic element of SL(2, \mathbb{Z});

^{*}The research of the first author was partially supported by NSF grant DMS-0900968. The research of the second author was partially supported by NSF grant DMS-1110145.

J. Lott and N. Sesum

see Subsection 3.1 for a more precise description. Under the additional assumption of a "solv-Gowdy" metric, it was shown that the Ricci flow approaches that of a locally homogeneous Sol-metric; see also [Kno00].

Passing to one-dimensional isometry groups, a natural class of geometries comes from warped product metrics with circle fibers and a closed surface base M. One starts with a product metric on $N = M \times S^1$ and then allows the circle length over $m \in M$ to become *m*-dependent. That is, we consider Riemannian metrics h on Nof the form

$$h = g + e^{2u} d\theta^2, \tag{1.1}$$

where g is a Riemannian metric on $M, u \in C^{\infty}(M)$ and θ is the standard coordinate on S^1 . It is not hard to see that the Ricci flow preserves the warped product structure. The Ricci flow equation on N becomes two coupled evolution equations on M for g(t) and u(t). The evolution equation for g(t) is like the Ricci flow equation on the surface M, but there is an extra term involving u(t). Because of this extra term, the techniques used to analyze Ricci flow on surfaces [CK04, Chapter 5] break down.

When M is a two-sphere, Xiaodong Cao showed that the product of \mathbb{R} and a cigar soliton cannot arise as a finite-time dilation limit of a warped product Ricci flow on $S^2 \times S^1$ ([Cao05]). His argument used an isoperimetric inequality. (Cao's work preceded Perelman's proof that the product of \mathbb{R} and a cigar soliton can never arise as a finite-time dilation limit for the Ricci flow on a compact three-manifold [Per02].)

In the present paper we use new techniques to give general results about the Ricci flow on three-manifolds with symmetries. One of our tools is the result of [Lot10] giving the long-time behavior of a three-dimensional Ricci flow solution $(N, h(\cdot))$ satisfying $\max_{p \in N} |\operatorname{Rm}^N|(p,t) = O(t^{-1})$ and $\operatorname{diam}(N, h(t)) = O(\sqrt{t})$. Thus one of our main goals is to show that these bounds are satisfied in the relevant cases. In particular, we show that $\max_{p \in N} |\operatorname{Rm}^N|(p,t) = O(t^{-1})$ for all of the immortal Ricci flows under consideration. (A Ricci flow solution is immortal if it exists for $t \in [0, \infty)$.) It is an open question whether an immortal Ricci flow solution on a compact three-manifold always satisfies this curvature bound.

To describe the results of the paper, we start with warped products. In the special case of a product metric, the Ricci flow solution is the isometric product of S^1 with a Ricci flow solution on the base M. Such two-dimensional Ricci flows are well understood [CK04, Chapter 5]. If h(0) is a warped product metric on $N = M \times S^1$ then a natural conjecture is that the Ricci flow asymptotically approaches a product flow.

Theorem 1.1. Let $h(\cdot)$ be a Ricci flow solution on a closed connected orientable three-dimensional manifold N. Suppose that h(0) is a warped product metric as in (1.1) with a two-dimensional orientable base M.

(i) If $\chi(M) > 0$ then there is a finite singularity time $T < \infty$. As $t \to T^-$, the lengths of the circle fibers remain uniformly bounded above and below. For any

 $p \in N$, the pointed smooth limit $\lim_{t\to T^-} (N, p, \frac{1}{T-t}h(t))$ exists and is the isometric product of \mathbb{R} with a sphere S^2 of constant curvature $\frac{1}{2}$.

- (ii) If $\chi(M) \leq 0$ then the Ricci flow exists for all $t \in [0, \infty)$. Also, there is a constant $C < \infty$ such that for all $p \in N$ and $t \in [0, \infty)$, one has $|\operatorname{Rm}^N| (p, t) \leq \frac{C}{t}$.
- (iii) If $\chi(M) = 0$ then $\lim_{t \to \infty} h(t)$ exists and is a flat metric on T^3 . The convergence is exponentially fast.
- (iv) If $\chi(M) < 0$, put $\hat{g}(t) = \frac{g(t)}{t}$. For any $i_0 > 0$, define the i_0 -thick part of $(M, \hat{g}(t))$ by

$$X_{i_0}(t) = \{ m \in M : \operatorname{inj}_{\hat{g}(t)}(m) \ge i_0 \}.$$
 (1.2)

Then

$$\lim_{t \to \infty} \max_{x \in X_{i_0}(t)} |R_{\hat{g}(t)}(x) + 1| = 0$$
(1.3)

and

$$\lim_{t \to \infty} \max_{x \in X_{i_0}(t)} |\widehat{\nabla} u|_{\hat{g}(t)}(x) = 0.$$
 (1.4)

For all sufficiently small i_0 , if t is sufficiently large then $X_{i_0}(t)$ is nonempty.

Remark 1.2. The proof of Theorem 1.1 (i) is essentially contained in List's paper [Lis08] on a modified Ricci flow. The only thing missing from [Lis08] is the observation that his flow differs from the warped product flow by a Lie derivative.

Remark 1.3. The proof of the curvature bound in Theorem 1.1 (ii) is by contradiction, using a blowup argument, a sharp volume estimate and the Gauss–Bonnet theorem.

Remark 1.4. The proof of Theorem 1.1 (iii) is somewhat indirect. We first show that $Vol(M, g(t)) = O(\sqrt{t})$ and that the length of the shortest noncontractible curve on M is nondecreasing in t. A geometric argument then shows that $diam(M, g(t)) = O(\sqrt{t})$. From [Lot10], we deduce that

$$\left(\max_{p \in N} |\operatorname{Rm}^{N}|(p,t)\right) \cdot \operatorname{diam}^{2}(N,h(t)) = o(t).$$
(1.5)

Rescaling at a given time t to diameter one, if t is sufficiently large then we can assume that $\max_{p \in N} |\operatorname{Rm}^N|(p, t)$ is arbitrarily small. After passing to a finite cover \hat{N} , we can assume that there is a universal lower bound on the injectivity radius of the pullback metric $\hat{h}(t)$. By the linear stability of flat metrics [GIK02], $\lim_{t\to\infty} \hat{h}(t)$ exists and is a flat metric. Hence $\lim_{t\to\infty} h(t)$ exists and is a flat metric.

Remark 1.5. Theorem 1.1 (iv) says that in the case $\chi(M) < 0$ and as $t \to \infty$, over a large part of M the flow (N, h(t)) approaches a product flow of S^1 times a finite-volume surface of constant sectional curvature $-\frac{1}{2t}$. Our result here is possibly nonoptimal; see Remark 2.20.

Our other main result is about Ricci flow solutions with a local $U(1) \times U(1)$ symmetry.

Theorem 1.6. Let N be an orientable three-manifold that fibers over S^1 with T^2 -fibers. Choosing an orientation for S^1 , let $H \in SL(2, \mathbb{Z}) = \pi_0(\text{Diff}^+(T^2))$ be the holonomy of the torus bundle. We can consider N to be the total space of a twisted principal $U(1) \times U(1)$ bundle, where the twisting is determined by H.

Let $h(\cdot)$ be a Ricci flow solution on N. Suppose that h(0) is invariant under the local $U(1) \times U(1)$ actions. Then the Ricci flow exists for all $t \in [0, \infty)$. There is a constant $C < \infty$ such that for all $p \in N$ and $t \in [0, \infty)$, one has $|\operatorname{Rm}^N| (p, t) \leq \frac{C}{t}$.

- (i) If H is elliptic, i.e. has finite order, then $\lim_{t\to\infty} h(t)$ exists and is a flat metric on N. The convergence is exponentially fast.
- (ii) Suppose that H is hyperbolic, i.e. has two distinct real eigenvalues. We write h(t) in the form

$$h(t) = g_{yy}(y,t) dy^{2} + (dx)^{T} G(y,t) dx, \qquad (1.6)$$

where $\{x^1, x^2\}$ are local coordinates on T^2 and $y \in [0, 1)$ is a local coordinate on S^1 . Then up to an overall change of parametrizations for S^1 and T^2 , we have

$$\lim_{t \to \infty} \frac{g_{yy}(y,t)}{t} = \frac{1}{2} \operatorname{Tr}(X^2), \qquad (1.7)$$
$$\lim_{t \to \infty} G(y,t) = e^{yX},$$

where X is the real symmetric matrix such that $e^X = H^T H$. The convergence in (1.7) is power-decay fast in t.

Remark 1.7. Theorem 1.6 (ii) says that $(N, h(\cdot))$ approaches a locally homogeneous Ricci flow solution of Sol-type. The proof of Theorem 1.6 (ii) uses the monotonicity of the modified W_+ -functional from [Lot10, Section 4.2.3], along with the local stability result of [Kno09, Theorem 3].

Remark 1.8. Theorem 1.6 (i) includes the cases considered in [CIJ90] and [Ham95, Section 11]. Theorem 1.6 (ii) includes the case considered in [HI93].

The structure of the paper is as follows. In Section 2 we prove Theorem 1.1. In Section 3 we prove Theorem 1.6. More detailed descriptions are at the beginnings of the sections.

We are grateful to Jim Isenberg for introducing us to these problems and for sharing his knowledge with us. We learned the estimates (2.19), (2.24) and (2.26) from Jim.

In what follows we will use the Einstein summation convention freely.

2. Warped products

In this section we prove Theorem 1.1. In Subsection 2.1 we write the warped product Ricci flow equations and the corresponding evolution equations for geometric quantities. In Subsection 2.2 we give *a priori* bounds using the maximum principle and integral estimates. In Subsection 2.3 we deal with the case $\chi(M) > 0$. In Subsection 2.4 we prove that the Ricci flow exists for $t \in [0, \infty)$ when $\chi(M) \leq 0$. In Subsection 2.5 we deal with the case $\chi(M) = 0$. In Subsection 2.6 we deal with the case $\chi(M) < 0$.

2.1. Warped product Ricci flow. Let $N = M \times S^1$ be endowed with a warped product metric

$$h = g + e^{2u} d\theta^2. \tag{2.1}$$

Here g is a Riemannian metric on M, u is a smooth function on M and $\theta \in [0, 2\pi)$. We use i, j, k, l for the indices on M. Nonzero components of the curvature tensor of (N, h) are

$$R_{ijkl}^{N} = R_{ijkl}^{M},$$

$$R_{\theta i\theta j}^{N} = -e^{2u} \left[\nabla_{i} \nabla_{j} u + (\nabla_{i} u) (\nabla_{j} u) \right].$$
(2.2)

Its square norm is

$$\left|\operatorname{Rm}_{h}^{N}\right|^{2} = \left|\operatorname{Rm}_{g}^{M}\right|^{2} + 2g^{ii'}g^{jj'}\left[\nabla_{i}\nabla_{j}u + (\nabla_{i}u)(\nabla_{j}u)\right]\left[\nabla_{i'}\nabla_{j'}u + (\nabla_{i'}u)(\nabla_{j'}u)\right].$$
(2.3)

We will often think of $|\operatorname{Rm}_{h}^{N}|(\cdot, t)$ as a function on M rather than N, since it pulls back from M, and write $|\operatorname{Rm}_{h}^{N}|(m, t)$. Note that a sectional curvature bound on (N, h) implies the same sectional curvature bound on (M, g).

The nonzero components of the Ricci tensor are

$$\operatorname{Ric}_{ij}^{N} = \operatorname{Ric}_{ij}^{M} - \nabla_{i}\nabla_{j}u - (\nabla_{i}u)(\nabla_{j}u),$$

$$\operatorname{Ric}_{\theta\theta}^{N} = -e^{2u} \left(\Delta u + |\nabla u|^{2} \right).$$
(2.4)

The scalar curvature is

$$R^N = R^M - 2\triangle u - 2|\nabla u|^2.$$
(2.5)

The Ricci flow equation

$$\frac{dh}{dt} = -2\operatorname{Ric}_{h(t)} \tag{2.6}$$

on N preserves the warped product structure. In terms of g and u, it becomes

$$\frac{\partial g_{ij}}{\partial t} = -2R_{ij} + 2\nabla_i \nabla_j u + 2(\nabla_i u)(\nabla_j u),$$

$$\frac{\partial u}{\partial t} = \Delta u + |\nabla u|^2,$$
(2.7)

where $\Delta = \Delta_{g(t)}$. Adding the Lie derivative with respect to $-\nabla u$ to the right-hand side of (2.7) gives the modified equations

$$\frac{\partial g_{ij}}{\partial t} = -2R_{ij} + 2(\nabla_i u)(\nabla_j u),
\frac{\partial u}{\partial t} = \Delta u.$$
(2.8)

Hereafter we will mainly work with the system (2.8), since geometric statements about (2.8) will imply the corresponding statements about the Ricci flow (2.7).

Given s > 0 and a solution $(g(\cdot), u(\cdot))$ of (2.8), we obtain another solution $(g_s(\cdot), u_s(\cdot))$ of (2.8) by putting

$$g_s(t) = \frac{1}{s}g(st),$$

$$u_s(t) = u(st).$$
(2.9)

Note that the rescaling in (2.9) differs from the three-dimensional rescaling of the Ricci flow $(N, h(\cdot))$, which would give

$$G_{s}(t) = \frac{1}{s}g(st),$$

$$U_{s}(t) = u(st) - \frac{1}{2}\ln(s).$$
(2.10)

The rescaling in (2.9) can be interpreted in the following way. There is a \mathbb{Z} -cover $\hat{N} = M \times \mathbb{R}$ of N with a pullback Ricci flow $\hat{h}(\cdot)$. On \hat{N} , the rescaling in (2.9) amounts to looking at the Ricci flow solution $\frac{1}{s}\phi_s^*\hat{h}(st)$, where $\phi_s(m, r) = (m, \sqrt{sr})$. We refer to [Lot07, Section 4] for further discussion of this point.

Put

$$h_s = g_s + e^{2u_s} d\theta^2. aga{2.11}$$

One sees from (2.3) and (2.9) that

$$\left|\operatorname{Rm}_{h_s}\right|^2 = s^2 \left|\operatorname{Rm}_h\right|^2.$$
(2.12)

This can also be seen from the discussion of the previous paragraph on \hat{N} along with the fact that under the three-dimensional rescaling of (2.10), the norm square of the curvature gets multiplied by s^2 .

Vol. 89 (2014)

Ricci flow on three-dimensional manifolds with symmetry

Following [Lis08], we introduce the tensor

$$S_{ij} = R_{ij} - (\nabla_i u)(\nabla_j u) \tag{2.13}$$

and its trace

$$S = R - |\nabla u|^2. \tag{2.14}$$

Putting $\alpha_n = 1$ in [Lis08, Lemma 3.2] gives

$$\frac{\partial |\nabla u|^2}{\partial t} = \Delta |\nabla u|^2 - 2|\operatorname{Hess}(u)|^2 - 2|\nabla u|^4$$
(2.15)

and

$$\frac{\partial S}{\partial t} = \Delta S + 2|S_{ij}|^2 + 2|\Delta u|^2.$$
(2.16)

Hereafter we specialize to the case when M is closed, connected and orientable, with dim(M) = 2. Then (2.8) simplifies to

$$\frac{\partial g_{ij}}{\partial t} = -Rg_{ij} + 2(\nabla_i u)(\nabla_j u), \qquad (2.17)$$
$$\frac{\partial u}{\partial t} = \Delta u.$$

Note that if u is nonconstant then the conformal class of g(t) is t-dependent.

2.2. A priori bounds

2.2.1. Bounds from the maximum principle. We take the Ricci flow to start at time zero. Applying the maximum principle to the second equation in (2.8) shows that there exist constants $C_1, C_2 > 0$ such that for all $m \in M$ and all t for which the flow exists,

$$C_1 \le u(m,t) \le C_2.$$
 (2.18)

Next, applying the maximum principle to (2.15) implies (cf. [Lis08, Lemma 5.6]) that

$$|\nabla u|^2(\cdot, t) \le \frac{c}{2ct+1},\tag{2.19}$$

where $c = \max_{p \in M} |\nabla u|^2(p, 0)$. Similarly, applying the maximum principle to (2.16) implies that

$$S(\cdot, t) \ge -\frac{1}{t}.$$
(2.20)

In particular,

$$R(\cdot, t) \ge -\frac{1}{t}.$$
(2.21)

J. Lott and N. Sesum

2.2.2. Volume estimates. Let V(t) denote the volume of $(M^2, g(t))$. As

$$\frac{d}{dt}dV_{g(t)} = \frac{1}{2}g^{ij}\frac{\partial g_{ij}}{\partial t}dV_{g(t)} = \left(-R + |\nabla u|^2\right)dV_{g(t)}, \qquad (2.22)$$

we have

$$\frac{dV}{dt} = -4\pi \cdot \chi(M) + \int_M |\nabla u|^2 \, dV_{g(t)},\tag{2.23}$$

where $\chi(M)$ is the Euler characteristic of M. Thus $\frac{dV}{dt} \ge -4\pi \cdot \chi(M)$ and so

$$V(t) \ge -4\pi \cdot \chi(M) \cdot t + V(0). \tag{2.24}$$

Using (2.19) in addition, we can control V(t) from above. Namely, from (2.19),

$$\frac{dV}{dt} \le -4\pi \cdot \chi(M) + \frac{c}{2ct+1} \cdot V \tag{2.25}$$

where c is as in (2.19). Then by ODE comparison,

$$V(t) \le -\frac{4\pi}{c} \cdot \chi(M) \cdot (2ct+1) + \sqrt{2ct+1} \cdot \left(\frac{4\pi \cdot \chi(M)}{c} + V(0)\right). \quad (2.26)$$

If $\chi(M) = 0$ then estimates (2.24) and (2.26) imply that

$$V(0) \le V(t) \le C(1 + \sqrt{t})$$
 (2.27)

for an appropriate constant $C < \infty$.

When $\chi(M) < 0$, the linear term on the right-hand side of (2.26) is $-8\pi\chi(M)t$. We would like to improve this to $-4\pi\chi(M)t$, to bring it in line with (2.24). Put $E(t) = \int_M |\nabla u|^2 dV_{g(t)}$.

Lemma 2.1. We have

$$\frac{dE}{dt} \le -\frac{E^2(t)}{V(t)}.$$
(2.28)

Proof. First, as in (2.22), the volume density changes by

$$\frac{d}{dt}(dV_{g(t)}) = -SdV_{g(t)}.$$
(2.29)

Note that

$$\int_{M} S \, dV_{g(t)} = \int_{M} (R - |\nabla u|^2) \, dV_{g(t)} = 4\pi \chi(M) - E(t).$$
(2.30)

CMH

Then

$$-\frac{dE}{dt} = \frac{d}{dt} \int_{M} S \, dV_{g(t)} = \int_{M} (\Delta S + 2|S_{ij}|^{2} + 2(\Delta u)^{2} - S^{2}) \, dV_{g(t)}$$

$$= 2 \int_{M} \left|S_{ij} - \frac{1}{2} S g_{ij}\right|^{2} dV_{g(t)} + 2 \int_{M} (\Delta u)^{2} \, dV_{g(t)}$$

$$= \int_{M} \left(2 \left| -\nabla_{i} u \nabla_{j} u + \frac{1}{2} |\nabla u|^{2} g_{ij}\right|^{2} + 2(\Delta u)^{2}\right) dV_{g(t)}$$

$$= \int_{M} (|\nabla u|^{4} + 2(\Delta u)^{2}) \, dV_{g(t)}$$

$$\geq \int_{M} |\nabla u|^{4} \, dV_{g(t)}$$

$$\geq \frac{1}{V(t)} \left(\int_{M} |\nabla u|^{2} \, dV_{g(t)}\right)^{2},$$
(2.31)

which proves the lemma.

We now show that E(t) decays logarithmically in time when $\chi(M) < 0$.

Corollary 2.2. If $\chi(M) < 0$ then there exist constants A, B > 0 such that

$$\int_{M} |\nabla u|^2 \, dV_{g(t)} \le \frac{A}{1 + B \ln(t+1)} \tag{2.32}$$

at all t for which the flow exists.

Proof. By Lemma 2.1 and the volume estimate (2.26) we have

$$\frac{dE}{dt} \le -\frac{E^2(t)}{V(t)} \le -\frac{E^2(t)}{c_1 t + c_2} \tag{2.33}$$

for appropriate constants $c_1, c_2 > 0$. The corollary follows from ODE comparison.

Corollary 2.3. If $\chi(M) < 0$ then there is a function $\alpha \colon [0, \infty) \to [0, \infty)$, with $\lim_{t\to\infty} \alpha(t) = 0$, such that

$$\left|\frac{V(t)}{-4\pi\chi(M)\cdot(t+1)} - 1\right| \le \alpha(t) \tag{2.34}$$

at all t for which the flow exists.

Proof. This follows from (2.23) and (2.32).

Lemma 2.4. The quantity $\frac{V(t)}{t}$ is nonincreasing along the flow (2.8).

Proof. From (2.20) and (2.29),

$$\frac{d}{dt}\left(\frac{V(t)}{t}\right) = -\frac{1}{t}\int_{M}\left(S + \frac{1}{t}\right)dV_{g(t)} \le 0.$$
(2.35)

This proves the lemma.

In the sequel we will distinguish between the cases $\chi(M) > 0$, $\chi(M) = 0$ and $\chi(M) < 0$.

2.3. Positive Euler characteristic

Proposition 2.5. If $\chi(M) > 0$ then there is a finite singularity time $T < \infty$. For any $p \in N$, the pointed smooth limit $\lim_{t\to T^-} (N, p, \frac{1}{T-t}h(t))$ exists and is the isometric product of \mathbb{R} with a sphere S^2 of constant curvature $\frac{1}{2}$.

Proof. If a smooth flow existed for all $t \in [0, \infty)$ then equation (2.26) would imply that V(t) < 0 for large t, which is impossible. Thus there is a singularity at some time $T < \infty$.

From [Lis08, Theorem 5.15], $\lim_{t\to T^-} \max_{m\in M} |R|(m,t) = \infty$, where R is the scalar curvature of M. Let $\{t_k\}_{k=1}^{\infty}$ be a sequence of times such that for sufficiently large k, t_k is the first time t for which $\max_{m\in M} |R|(m,t) = k$. Let $m_k \in M$ be such that $|R|(m_k, t_k) = k$. From [Lis08, Theorem 7.9], a subsequence of the rescaled pointed solutions

$$(M, m_k, g_k(t), u_k(t)) = (M, m_k, kg(t_k + t/k), u(t_k + t/k))$$
(2.36)

converges smoothly to a solution $(M_{\infty}, m_{\infty}, g_{\infty}(\cdot), u_{\infty}(\cdot))$ defined for $t \in (-\infty, 0]$, where $g_{\infty}(\cdot)$ is a κ -solution on M_{∞} in the sense of [Per02, Section 11.1] and $u_{\infty}(\cdot)$ is constant both spatially and temporally. The proof of this statement uses a modified W-functional which was introduced in [Lis08] and [Lot10], and which becomes

$$W(g, u, f, \tau) = \int_{M} \left[\tau \left(|\nabla f|^{2} + R - |\nabla u|^{2} \right) + f - 2 \right] (4\pi\tau)^{-1} e^{-f} dV_{g(t)}.$$
(2.37)

in our three-dimensional warped product case.

The only κ -solution on an orientable surface is the round shrinking 2-sphere [KL08, Corollary 40.1]. In particular, $\lim_{k\to\infty} (M, kg(t_k), u(t_k)) = (S^2, g_{S^2}, u_{\infty})$, where g_{S^2} has constant scalar curvature one. (Because S^2 is compact, we no longer have to refer to basepoints. At this point our convergence is still modulo diffeomorphisms.) Standard arguments show that the system (2.7) is stable around the solution given by the round shrinking S^2 and constant u, with exponential convergence for

CMH

the normalized flow. It follows that $\lim_{t\to T^-} (M, \frac{1}{T-t}g(t), u(t)) = (S^2, g_{S^2}, u_\infty)$, where the convergence is now taken without diffeomorphisms.

In terms of the three-dimensional geometry, from (2.18) the fiber lengths are uniformly bounded above and below by positive constants, up to time T. The three-dimensional pointed limit $\lim_{t\to T^-} (N, \frac{1}{T-t}h(t), p)$ is the isometric product of \mathbb{R} with (S^2, g_{S^2}) . This proves the proposition.

Remark 2.6. We could also prove Proposition 2.5 by looking at the three-dimensional singularity models with a nowhere-vanishing Killing vector field. Such a proof would be less elementary, since it would use the results of [Per02, Section 11].

Remark 2.7. The method of proof of Proposition 2.5 works for finite-time singularities of warped products $M \times S^1$ if M is a compact manifold of arbitrary dimension n-1. Blowing up at points of maximal curvature, one obtains an (n-1)-dimensional ancient solution which is κ -noncollapsed at all scales.

2.4. Nonsingularity when $\chi(M) \leq 0$

Proposition 2.8. If $\chi(M^2) \leq 0$ then the Ricci flow exists for $t \in [0, \infty)$.

Proof. If not then there is a singularity at some time $T < \infty$. The same argument as in Subsection 2.3 gives $\lim_{k\to\infty} (M, kg(t_k), u(t_k)) = (S^2, g_{S^2}, u_\infty)$. In particular, M is diffeomorphic to S^2 , which contradicts our assumption.

2.5. Vanishing Euler characteristic

Proposition 2.9. If $\chi(M) = 0$ then $\lim_{t\to\infty} h(t)$ exists and is a flat metric on T^3 . The convergence is exponentially fast.

Proof. We first show in the following two propositions that if $\chi(M) = 0$ then $\max_{p \in N} |\operatorname{Rm}^{N}|(p,t) = O(t^{-1})$ and $\operatorname{diam}(N,h(t)) = O(\sqrt{t})$. Here Rm^{N} denotes the three-dimensional sectional curvatures.

Proposition 2.10. If $\chi(M) = 0$ then there is a $C < \infty$ such that for all $t \in [0, \infty)$, we have

$$t \cdot \max_{p \in N} \left| \operatorname{Rm}^{N} \right| (p, t) \le C.$$
(2.38)

Proof. If not, $\limsup_{t\to\infty} t \cdot \max_{m\in M} |\operatorname{Rm}^N|(m,t) = \infty$. (Since the function $|\operatorname{Rm}^N|(\cdot,t)$ pulls back from M, we can think of it as a function on M.) We perform an analog of Hamilton's pointpicking algorithm for a type IIb Ricci flow solution; see

J. Lott and N. Sesum

[CLN06, Chapter 8.2.1.3]. Namely, take any sequence $\{T_i\}_{i=1}^{\infty}$ with $\lim_{i\to\infty} T_i = \infty$ and let $(m_i, t_i) \in M \times [0, T_i]$ be such that

$$t_i(T_i - t_i) \left| \mathrm{Rm}^N \right| (m_i, t_i) = \sup_{(m,t) \in M \times [0, T_i]} t(T_i - t) \left| \mathrm{Rm}^N \right| (m, t).$$
(2.39)

Dilate the flow in space and time by $Q_i = |\operatorname{Rm}^N|(m_i, t_i)$ in the following way:

$$g_i(t) = Q_i \cdot g(t_i + tQ_i^{-1}), \quad u_i(t) = u(t_i + tQ_i^{-1}).$$
 (2.40)

Then $(g_i(\cdot), u_i(\cdot))$ satisfies (2.8) on a time interval (A_i, Ω_i) , with $\lim_i A_i = -\infty$ and $\lim_i \Omega_i = \infty$. Put

$$h_i(t) = g_i(t) + e^{2u_i} d\theta^2.$$
 (2.41)

By construction, $\left| \operatorname{Rm}_{h_i}^N \right| (m_i, 0) = 1$ and after redefining A_i and Ω_i , there is a sequence $\{\gamma_i\}_{i=1}^{\infty}$ with $\lim_{i \to \infty} \gamma_i = 1$ such that

$$\max_{(m,t)\in M\times[A_i,\Omega_i]} \left| \operatorname{Rm}_{h_i}^N \right|(m,t) \le \gamma_i.$$
(2.42)

The curvature bound on N implies a curvature bound on M which, along with the *a priori* bounds on u_i and $|\nabla u_i|$, implies higher derivative bounds on $\operatorname{Rm}^M(g_i)$ and u_i [Lis08, Theorem 5.12]. We would now like to take a convergent subsequence of the pointed flows $\{(M \times [A_i, \Omega_i], (m_i, 0), g_i(\cdot), u_i(\cdot))\}_{i=1}^{\infty}$ to obtain an eternal solution $(M_{\infty} \times \mathbb{R}, (m_{\infty}, 0), g_{\infty}(\cdot), u_{\infty}(\cdot))$ of (2.8), where \mathbb{R} denotes a time interval. To do so, we need a uniform positive lower bound on the injectivity radius at $(m_i, 0)$. If the two-dimensional manifolds $(M, g_i(0))$ were positively curved then such a bound would be automatic. Since we don't know that $(M, g_i(0))$ is positively curved, we argue differently.

Lemma 2.11. There is some $\epsilon > 0$ such that for all *i*, the injectivity radius of $(M, g_i(0))$ at m_i is bounded below by ϵ .

Proof. Suppose that the lemma is false. Then after passing to a subsequence, we can assume that $\lim_{i\to\infty} \inf_{g_i(0)}(m_i) = 0$. Passing to a further subsequence, we can assume that $\{(M \times [A_i, \Omega_i], (m_i, 0), g_i(\cdot), u_i(\cdot))\}_{i=1}^{\infty}$ converges to a solution of (2.8) on an étale groupoid. For information about the use of étale groupoids in Ricci flow, we refer to [Lot07] and [Lot10]. The upshot is that after passing to a subsequence, we have smooth pointed convergence to an eternal solution $(M_{\infty} \times \mathbb{R}, (m_{\infty}, 0), g_{\infty}(\cdot), u_{\infty}(\cdot))$ of (2.8), where M_{∞} is a two-dimensional étale groupoid. Furthermore, this solution has uniformly bounded curvature by (2.42), and for each $t \in \mathbb{R}, (M_{\infty}, g_{\infty}(t))$ is a complete closed effective Riemannian groupoid.

By assumption, $\lim_{i\to\infty} t_i Q_i = \infty$. As

$$\max_{m \in M} |\nabla u_i|_{g_i}^2(m, t) = Q_i^{-1} \max_{m \in M} |\nabla u|_g^2(m, t_i + tQ_i^{-1}), \qquad (2.43)$$

when combined with (2.19) we conclude that $\nabla u_{\infty} = 0$ on any time slice. Then u_{∞} is constant spatially and temporally, and g_{∞} is just a Ricci flow on M_{∞} . As it is an eternal solution, it has nonnegative curvature, with positive curvature at $(m_{\infty}, 0)$.

We already know that $(M_{\infty}, m_{\infty}, g_{\infty}(t))$ is a bounded curvature limit of $\{(M, m_i, g_i(t))\}_{i=1}^{\infty}$. The Riemannian groupoid $(M_{\infty}, g_{\infty}(t))$ has a locally constant sheaf of finite dimensional Lie algebras g which act as germs of Killing vector fields on the unit space $M_{\infty}^{(0)}$. Because of the bounded curvature assumption, these local Killing vector fields do not have a point of common vanishing; see, for example, [Ron07, Theorem 5.1].

As $M_{\infty}^{(0)}$ is two-dimensional, and we are in the collapsing situation, the only possibilities for g are \mathbb{R}^2 and \mathbb{R} . If $g = \mathbb{R}^2$ then $(M_{\infty}, g_{\infty}(t))$ is flat, which is a contradiction.

Suppose that $\mathfrak{g} = \mathbb{R}$. Locally, $g_{\infty}(t)$ can be written as $dx^2 + f^2(x,t)dy^2$. Since its curvature is $K = -\frac{f''}{f}$, the function f is concave. Fixing the value of f at a single point in $M_{\infty}^{(0)}$, the function f pulls back from a function on the orbit space \mathcal{O} of M_{∞} , which in our case is a one-dimensional orbifold [Lot10, Proposition 5.2]. If \mathcal{O} is a circle then we immediately get a contradiction, since the concave function fmust be constant, but this contradicts the fact that (M_{∞}, g_{∞}) has nonzero curvature at $(m_{\infty}, 0)$. If \mathcal{O} is an interval orbifold then we can pass to a double cover and argue as before. If \mathcal{O} is \mathbb{R} then the positive concave function f must be constant, which again contradicts the fact that (M_{∞}, g_{∞}) has nonzero curvature at $(m_{\infty}, 0)$. If \mathcal{O} is $[0, \infty)$ then we can pass to a double cover and argue as before.

This proves the lemma.

We can now take a convergent subsequence of the pointed flows

$$\{(M \times [A_i, \Omega_i], (m_i, 0), g_i(\cdot), u_i(\cdot))\}_{i=1}^{\infty}$$

to obtain an eternal solution $(M_{\infty} \times \mathbb{R}, (m_{\infty}, 0), g_{\infty}(\cdot), u_{\infty}(\cdot))$ of (2.8). From (2.19) and (2.43), u_{∞} is constant. Thus we have a nonflat eternal Ricci flow solution on the two-dimensional manifold M_{∞} , with uniformly bounded curvature (from (2.42)) and complete time slices. In particular, $\int_{M_{\infty}} R_{g_{\infty}(0)} dV_{g_{\infty}(0)} > 0$. Although we won't really need it, $(M_{\infty}, g_{\infty}(\cdot))$ must be the cigar soliton, as follows from the fact that the spacetime supremum of R is achieved at a point, along with a differential Harnack inequality [CLN06, Theorem 9.4]. Hence

$$\int_{M_{\infty}} R_{g_{\infty}(0)} \, d\, V_{g_{\infty}(0)} = 4\pi.$$
(2.44)

For large *i*, there is a bounded domain $S_i \subset (M, g(t_i))$ which, after rescaling, is almost isometric to a large piece of the time-zero slice of the cigar soliton. Then for large *i*,

$$\int_{S_i} R_{g(t_i)} \, dV_{g(t_i)} \ge 3\pi. \tag{2.45}$$

On the other hand,

$$\int_{M-S_i} R_{g(t_i)} \, dV_{g_i(t)} \ge -\frac{1}{t_i} \operatorname{Vol}_{g(t_i)}(M-S_i) \ge -\frac{V(t_i)}{t_i}.$$
 (2.46)

Adding (2.45) and (2.46), and using (2.27), we see that for large i,

$$\int_{M} R_{g(t_i)} \, dV_{g(t_i)} \ge 2\pi. \tag{2.47}$$

This contradicts the assumption that $\chi(M) = 0$ and proves Proposition 2.10

Proposition 2.12. diam $(N, h(t)) = O(\sqrt{t})$.

Proof. We will use the following general result about Riemannian submersions.

Lemma 2.13. If $\pi: (N,h) \to (M,g)$ is a Riemannian submersion, with N compact and connected, then

$$\operatorname{diam}(M,g) \le \operatorname{diam}(N,h) \le \operatorname{diam}(M,g) + 2 \max_{m \in M} \operatorname{diam}(\pi^{-1}(m)), \qquad (2.48)$$

where diam $(\pi^{-1}(m))$ is the intrinsic diameter of $\pi^{-1}(m)$.

Proof. Given $m_1, m_2 \in M$, we have

$$d_M(m_1, m_2) = d_N(\pi^{-1}(m_1), \pi^{-1}(m_2)).$$
(2.49)

It immediately follows that $diam(M, g) \leq diam(N, h)$.

Given $p_1, p_2 \in N$, put $m_1 = \pi(p_1)$ and $m_2 = \pi(p_2)$. Then

$$d_N(p_1, p_2) \le \operatorname{diam}(\pi^{-1}(m_1)) + d_N(\pi^{-1}(m_1), \pi^{-1}(m_2)) + \operatorname{diam}(\pi^{-1}(m_2)).$$
(2.50)
he lemma follows.

The lemma follows.

From (2.18) and Lemma 2.13, diam $(N, h(t)) \leq \text{diam}(M, g(t)) + 2e^{C_2}$. Thus it suffices to show that diam $(M, g(t)) = O(\sqrt{t})$.

Let L(t) be the length (with respect to g(t)) of a shortest noncontractible closed geodesic γ_t on M. We parametrize γ_t by an arclength parameter s. We will first show the following claim, which is an analog of [Ham95, Theorem 12.1].

Claim 2.14. L(t) is nondecreasing in t.

Proof. Given the curve γ_t at time t > 0, we obtain an upper bound on $L(t - \Delta t)$ by considering the length of the same curve γ_t at the earlier time $t - \Delta t$. Then

$$\lim_{\Delta t \to 0} \inf_{\Delta t} \frac{L(t) - L(t - \Delta t)}{\Delta t} \ge \frac{1}{2} \int_0^{L(t)} \frac{\partial g}{\partial t} (\gamma'_t, \gamma'_t) \, ds$$
$$= \int_0^{L(t)} \left[-\frac{1}{2} R(\gamma_t(s)) + \langle \gamma'_t, \nabla u \rangle^2 \right] ds.$$
(2.51)

As γ_t is stable, applying the second variation formula with respect to a parallel normal field along γ_t gives

$$-\int_{0}^{L(t)} R(\gamma_t(s)) \, ds \ge 0. \tag{2.52}$$

The claim follows.

Since $\chi(M) = 0$, equations (2.27) and (2.38) imply that

$$\lim_{t \to \infty} \left[\left(\max_{m \in M} |R|(m,t) \right) \cdot V(t) \right] = 0.$$
(2.53)

The next claim is purely geometric and has nothing to do with flows.

Claim 2.15. There exist $\epsilon > 0$ and $C < \infty$ such that for every metric g on $M = T^2$ with $(\max_{m \in M} |R_g|(m)) \cdot \operatorname{Vol}(M, g) \leq \epsilon$, we have

$$L(g) \cdot \operatorname{diam}(M, g) \le C \cdot \operatorname{Vol}(M, g), \qquad (2.54)$$

where L(g) is the length of the shortest noncontractible closed geodesic.

Proof. Suppose first that the metric g is flat. Let γ be a shortest closed geodesic and let \hat{M} be the corresponding cyclic cover of M. Then $\hat{M} = S^1 \times \mathbb{R}$, where the circle has length $L(\gamma) = L(g)$. A generator of the covering group \mathbb{Z} acts on \hat{M} by translation in the \mathbb{R} -direction, by some distance L', along with rotation around S^1 by some angle $\theta \in [-\pi, \pi]$. For any $\hat{m} \in \hat{M}$, we have

$$d_{\hat{M}}(\hat{m}, g\hat{m}) = \sqrt{(L')^2 + \left(\frac{\theta L}{2\pi}\right)^2}.$$
 (2.55)

This is the length of a closed geodesic on M, so we must have

$$\sqrt{(L')^2 + \left(\frac{\theta L}{2\pi}\right)^2} \ge L.$$
(2.56)

This implies that $L' \ge \frac{\sqrt{3}}{2}L$. Now diam $(M, g) \le \frac{1}{2}(L + L')$ and Vol(M, g) = LL', so the claim holds in this case with C = 2. Hence we can assume that g is nonflat.

Suppose that the claim is false. Then there is a sequence $\epsilon_i \to 0$ such that for each *i*, there is a metric g_i on *M* with

$$\left(\max_{m \in M} |R_{g_i}|(m)\right) \cdot \operatorname{Vol}(M, g_i) \le \epsilon_i$$
(2.57)

but

$$L(g_i) \cdot \operatorname{diam}(M, g_i) > i \cdot \operatorname{Vol}(M, g_i).$$
(2.58)

Rescale g_i so that $\max_{m \in M} |R_{g_i}|(m) = 1$. Then $\operatorname{Vol}(M, g_i) \leq \epsilon_i$.

Let $\delta \ll 1$ be a new parameter. Suppose first that for an infinite number of *i*, the pointed Gromov-Hausdorff distance from (M, m, g_i) to a point is greater than δ for all $m \in M$, i.e. diam $(M, g_i) > \delta$. After relabelling the sequence $\{(M, g_i)\}_{i=1}^{\infty}$, we can assume that for all *i* and all $m \in M$, the pointed Gromov-Hausdorff distance from (M, m, g_i) to a point is greater than δ . Since $\lim_{i\to\infty} \operatorname{Vol}(M, g_i) = 0$, there is a sequence $\{c_i\}_{i=1}^{\infty}$ with $\lim_{i\to\infty} c_i = 0$ such that for any $m \in M$, the pointed closed metric ball $(\overline{B}(m, \frac{\delta}{10}), m, g_i)$ is c_i -close to $([-\frac{\delta}{10}, \frac{\delta}{10}], 0)$ in the pointed Gromov-Hausdorff topology.

Fix $\delta' \ll 1$. From [CFG92], for large *i* there is a diffeomorphism ϕ_i from (M, g_i) to $(S^1 \times S^1, g'_i)$, where

- g'_i is a warped product metric $dx^2 + f^2_i(x)dy^2$,
- $\max_{x \in S^1} f_i(x) \to 0$ as $i \to \infty$, and
- ϕ_i is a $e^{\delta'}$ -biLipschitz map.

More precisely, we are using the fact that the results of [CFG92] hold in a localized sense, i.e. without an upper diameter bound; see [CT06, Section 2] for discussion. The paper [CFG92] gives a biLipschitz approximation of (M, g_i) by a Riemannian nilbundle with affine holonomy; see also [Fuk89]. In the present case such a Riemannian nilbundle is a Riemannian submersion $S^1 \times S^1 \rightarrow S^1$.

For the metric g'_i , let A_i be the length of the circle base. We know that $A_i \ge \frac{\delta}{10}$. To get an upper bound on $L(g_i)$, we locate a shortest circle fiber of the fiber bundle $S^1 \times S^1 \to S^1$ and take its preimage under ϕ_i . This gives a noncontractible closed curve in M, so

$$L(g_i) \le e^{\delta'} \min_{x \in S^1} f_i(x).$$
 (2.59)

Next, using Lemma 2.13, the diameter of $(S^1 \times S^1, g'_i)$ is bounded above by $A_i + 2 \max_{x \in S^1} f_i(x)$, so

diam
$$(M, g_i) \le e^{\delta'} (A_i + 2 \max_{x \in S^1} f_i(x)).$$
 (2.60)

The volume of $(S^1 \times S^1, g'_i)$ is bounded below by A_i times the length of the smallest circle fiber, so

$$Vol(M, g_i) \ge e^{-2\delta'} A_i \min_{x \in S^1} f_i(x).$$
 (2.61)

For large *i*, equations (2.59), (2.60) and (2.61) contradict (2.58).

Thus we can assume that for all but a finite number of *i*, there is some point m_i such that the pointed Gromov-Hausdorff distance from (M, m_i, g_i) to a point is at most δ . We now rescale (M, g_i) to a metric (M, \hat{g}_i) with diameter one. After this rescaling, $\max_{m \in M} |R_{\hat{g}_i}| (m) \leq \delta^2$. Suppose that $\liminf_{i \to \infty} \operatorname{Vol}(M, \hat{g}_i) = 0$. After passing to a subsequence, we can assume that $\lim_{i \to \infty} \operatorname{Vol}(M, \hat{g}_i) = 0$. As in the argument of the preceding paragraphs, for large *i* there is a biLipschitz approximation of (M, \hat{g}_i) by a warped product metric on a fiber bundle $S^1 \times S^1 \to S^1$, whose base has diameter close to one, and we obtain a contradiction to (2.58).

Thus there is some $v_0 > 0$ such that for all i, we have $Vol(M, \hat{g}_i) \ge v_0$. Cheeger compactness now gives a contradiction to (2.58). This proves the claim.

To conclude the proof of Proposition 2.12 we argue as follows. By (2.27), $V(t) = O(\sqrt{t})$. By (2.2) and Proposition 2.10, $\max_{m \in M} |R^M|(m, t) = O(t^{-1})$. Thus for large t, we can apply Claim 2.15 to conclude that

diam
$$(M, g(t)) \le C \frac{V(t)}{L(t)} \le C \frac{V(t)}{L(0)},$$
 (2.62)

where Claim 2.14 is used in the last inequality. Proposition 2.12 follows.

By [Lot10, Theorem 1.2.1], $\max_{m \in M} |\operatorname{Rm}_n^N|(m, t) = o(t^{-1})$ by Propositions 2.10 and 2.12. Thus

$$\lim_{t \to \infty} \left[\left(\max_{m \in M} |\operatorname{Rm}^N|(m, t) \right) \cdot \operatorname{diam}^2(N, h(t)) \right] = 0.$$
 (2.63)

For some given t, we rescale (N, h(t)) so that it has diameter one. We now wish to apply the local stability of flat metrics on T^3 under the Ricci flow. For this, we need the following lemma, which is purely geometric.

Claim 2.16. There are a compact subset K of the moduli space of flat metrics on T^3 and a function $\epsilon': [0, \infty) \to [0, \infty)$, with $\lim_{\epsilon \to 0} \epsilon'(\epsilon) = 0$, such that the following holds. Suppose that h is a Riemannian metric on $N = T^3$ with $\operatorname{diam}(N, h) = 1$ and $\max_{p \in N} |\operatorname{Rm}_h|(p) \le \epsilon$. Then there is a finite cover \hat{N} of N such that (\hat{N}, \hat{h}) has distance at most $\epsilon'(\epsilon)$ in the C^1 -topology from an element of K.

Proof. Given i_0 , $D_1 > 0$ and $D_2 < \infty$, let K_{i_0,D_1,D_2} denote the isometry classes of flat Riemannian metrics on T^3 with diameter in $[D_1, D_2]$ and injectivity radius bounded below by i_0 . For any $\alpha \in \mathbb{Z}^+$, the set K_{i_0,D_1,D_2} is compact in the C^{α} -topology. We will take $K = K_{i_0,D_1,D_2}$, where the parameters i_0 , D_1 and D_2 will be determined in the proof.

Given K, suppose that the claim is false. Then there is some $\epsilon' > 0$ along with a sequence $\{(N_i, h_i)\}_{i=1}^{\infty}$ of Riemannian 3-tori with diam $(N_i, h_i) = 1$ and $\max_{p \in N_i} |\operatorname{Rm}_{g_i}|(p) \leq \frac{1}{i}$, but with the property that for no *i* is there a finite cover (\hat{N}_i, \hat{h}_i) of (N_i, h_i) with distance less than ϵ' in the C^1 -topology from an element of K. Let \tilde{N}_i be the universal cover of N_i . Pick $p_i \in N_i$ and let \tilde{p}_i be a lift to \tilde{N}_i . As $i \to \infty$, $(\tilde{N}_i, \tilde{p}_i)$ converges to $(\mathbb{R}^3, 0)$ in the pointed C^1 -topology; this follows from a uniform lower bound on the injectivity radius at \tilde{p}_i [Ron07, (3.2)] and C^1 -convergence results. Let $r_{1,i} \in \pi_1(N_i, p_i)$ be a nontrivial element which

minimizes $d(\tilde{p}_i, r_{1,i} \tilde{p}_i)$. Let $r_{2,i} \in \pi_1(N_i, p_i) - \langle r_{1,i} \rangle$ minimize $d(\tilde{p}_i, r_{2,i} \tilde{p}_i)$. Let $r_{3,i} \in \pi_1(N_i, p_i) - \langle r_{1,i}, r_{2,i} \rangle$ minimize $d(\tilde{p}_i, r_{3,i} \tilde{p}_i)$. By the diameter condition, for large *i* we have $d(\tilde{p}_i, r_{3,i} \tilde{p}_i) \leq 10$.

Let $\gamma_{1,i}$, $\gamma_{2,i}$, $\gamma_{3,i}$ be minimizing geodesics from \tilde{p} to $r_{1,i}$, \tilde{p} , $r_{2,i}$, \tilde{p} , $r_{3,i}$, \tilde{p} . There is a universal constant c > 0 such that for each i, pairs from $\{\gamma'_{1,i}(0), \gamma'_{2,i}(0), \gamma'_{3,i}(0)\}$ form an angle at \tilde{p} which is at least c [Gro78, Section 2.3]. Let $k_{1,i}$ and $k_{2,i}$ be positive integers such that $k_{1,i}d(\tilde{p}_i, r_{1,i}, \tilde{p}_i)$ and $k_{2,i}d(\tilde{p}_2, r_{2,i}, \tilde{p}_i)$ lie in $[\frac{1}{10}, 10]$. Let \hat{N}_i be the cover of N_i such that $\pi_1(\hat{N}_i, \hat{p}_i)$ has generators $r_{1,i}^{k_{1,i}}$, $r_{2,i}^{k_{2,i}}$, $r_{3,i}$. After passing to a subsequence, we can assume that as $i \to \infty$, with respect to the approximations of \tilde{N}_i by \mathbb{R}^3 , the triples $r_{1,i}^{k_{1,i}}$, $r_{2,i}^{k_{2,i}}$, \tilde{p}_i , $r_{3,i}, \tilde{p}_i$ converge to vectors v_1, v_2, v_3 in \mathbb{R}^3 with lengths in $[\frac{1}{10}, 10]$ and mutual angles at least c. Then $\lim_{i\to\infty} (\hat{N}_i, \hat{h}_i) = \mathbb{R}^3/(\mathbb{Z}v_1 + \mathbb{Z}v_2 + \mathbb{Z}v_3)$ in the C^1 -topology. We can find i_0 , $D_1 > 0$ and $D_2 < \infty$, computed in terms of c, such that $\mathbb{R}^3/(\mathbb{Z}v_1 + \mathbb{Z}v_2 + \mathbb{Z}v_3)$ has diameter in $[D_1, D_2]$ and injectivity radius bounded below by i_0 . This contradicts the properties of the sequence $\{(N_i, h_i)\}_{i=1}^{\infty}$.

By (2.63), for any $\epsilon' > 0$, if t is large enough then we can apply Claim 2.16 to (N, h(t)). Using the higher derivative curvature estimates coming from the Ricci flow and applying a similar argument as in the proof of Claim 2.16, for any $\alpha \in \mathbb{Z}^+$ we can also say that there is a compact subset K_{α} of the moduli space of flat metrics on T^3 so that for any $\epsilon' > 0$, if t is large enough then a finite cover $(\hat{N}, \hat{h}(t))$ is ϵ' -close in the C^{α} -topology to an element of K_{α} . Taking α large, we can now apply [GIK02, Theorem 3.7] to conclude that as $t \to \infty$, $(\hat{N}, \hat{h}(t))$ (or more precisely the solution of (2.7)) converges to a flat metric on T^3 . Furthermore, the convergence is exponentially fast. (Strictly speaking, [GIK02] considers the evolution of a Riemannian metric h which is sufficiently close to a fixed flat metric, but the arguments clearly extend to the setting of closeness to a compact set K_{α} of flat metrics, since there will be uniform control on the constants.) Because of the equivariance of Ricci flow under isometries, the same is true for (N, h(t)).

This proves Proposition 2.9.

2.6. Negative Euler characteristic. By Proposition 2.8, the flow (2.8) or, equivalently, (2.7) exists forever. We start with the following proposition.

Proposition 2.17. We have $\max_{p \in N} |\text{Rm}^{N}|(p, t) = O(t^{-1})$.

Proof. The argument is similar to that in the proof of Proposition 2.10. If the statement were not true, i.e. if $\limsup_{t\to\infty} t \cdot \max_{m\in M} |\operatorname{Rm}^N|(m,t) = \infty$, then we begin the argument exactly the same as in the proof of Proposition 2.10 to take a limit of rescalings at times $\{t_i\}_{i=1}^{\infty}$. This limit is an eternal solution of the form $h_{\infty}(t) = g_{\infty}(t) + Cd\theta^2$, where C is a constant and $g_{\infty}(t)$ is a nonflat eternal solution to

the Ricci flow on a two-dimensional manifold M_{∞} , with complete time slices and nonnegative bounded curvature. As in the proof of Proposition 2.10, $(M_{\infty}, g_{\infty}(\cdot))$ must be a cigar soliton and

$$\int_{M_{\infty}} R_{g_{\infty}(0)} \, d\, V_{g_{\infty}(0)} = 4\pi. \tag{2.64}$$

For large *i*, there is a bounded domain $S_i \subset (M, g(t_i))$ which, after rescaling, is almost isometric to a large region in the cigar soliton. Then

$$\int_{S_i} R_{g(t_i)} \, dV_{g(t_i)} \ge 3\pi. \tag{2.65}$$

Furthermore, by (2.21) and Corollary 2.3, we have

$$\int_{M-S_i} R_{g(t_i)} \, dV_{g(t_i)} \ge -\frac{1}{t_i} \operatorname{Vol}_{g(t_i)}(M-S_i) \ge -\frac{V(t_i)}{t_i} \ge 4\pi \chi(M) \cdot (1+o(i^0)).$$
(2.66)

Adding (2.65) and (2.66), and taking $i \to \infty$, contradicts the Gauss–Bonnet theorem for M.

Put $\hat{g}(t) = \frac{g(t)}{t}$. Let $\hat{\nabla}$ denote the corresponding Levi-Civita connection.

Claim 2.18. There exist $i_0 > 0$ and $t_0 > 0$ such that for every $t \ge t_0$, there is a point $m_t \in M$ where the injectivity radius satisfies $\operatorname{inj}_{\hat{g}(t)}(m_t) \ge i_0$.

Proof. By (2.2) and Proposition 2.17, for $t \ge 1$ the metrics $(M, \hat{g}(t))$ have uniformly bounded curvature. If the claim were not true then for every $\epsilon > 0$, there would be some $t_{\epsilon} \ge 1$ such that $\operatorname{inj}_{\hat{g}(t_{\epsilon})}(m) < \epsilon$ for all $m \in M$. Then M would have an F-structure and hence a vanishing Euler characteristic [CG90]. This is a contradiction.

Proposition 2.19. For any $i_0 > 0$, define the i_0 -thick part of $(M, \hat{g}(t))$ by

$$X_{i_0}(t) = \{ m \in M : \operatorname{inj}_{\hat{g}(t)}(m) \ge i_0 \}.$$
(2.67)

Then

$$\lim_{t \to \infty} \sup_{x \in X_{i_0}(t)} |R_{\hat{g}(t)}(x) + 1| = 0$$
(2.68)

and

$$\lim_{t \to \infty} \sup_{x \in X_{i_0}(t)} |\widehat{\nabla} u|_{\hat{g}(t)}(x) = 0.$$
(2.69)

Proof. Suppose that the proposition is not true. Then there are some $i_0, \epsilon > 0$ along with sequences $t_i \to \infty$ and $\{m_i\}_{i=1}^{\infty}$ such that for each i, $\operatorname{inj}_{\hat{g}(t_i)}(m_i) \ge i_0$ and either $|R_{\hat{g}(t_i)}(m_i) + 1| \ge \epsilon$ or $|\widehat{\nabla}u|_{\hat{g}(t_i)}(m_i) \ge \epsilon$.

By Hamilton's compactness theorem and the derivative estimates on u from [Lis08, Section 5], after passing to a subsequence we can assume that there is a smooth pointed limit of flows

$$\lim_{i \to \infty} \left(M, m_i, \frac{1}{t_i} g(t_i t), u(t_i t) \right) = \left(M_{\infty}, m_{\infty}, g_{\infty}(t), u_{\infty}(t) \right).$$
(2.70)

For any bounded domain $S \subset M_{\infty}$, using Corollary 2.2 we have that at any time a,

$$\int_{S} |\nabla u_{\infty}|^{2}_{g_{\infty}(a)} dV_{g_{\infty}(a)} \leq \limsup_{i \to \infty} \int_{M} |\nabla u(t_{i}a)|^{2}_{g(t_{i}a)} dV_{g(t_{i}a)} = 0.$$
(2.71)

Thus u_{∞} is spatially constant at time *a*. Because *a* is arbitrary, and u_{∞} satisfies the time-dependent heat equation, it follows that u_{∞} is also temporally constant. In particular,

$$0 = |\nabla u_{\infty}|_{g_{\infty}(1)}(m_{\infty}) = \lim_{i \to \infty} |\widehat{\nabla} u|_{\widehat{g}(t_i)}(m_i).$$

$$(2.72)$$

Now $R_{g_{\infty}(t)} + \frac{1}{t} \ge 0$. Given $0 < a < b < \infty$, equations (2.35) and (2.72), along with Corollary 2.3, give

$$\int_{a}^{b} \int_{M_{\infty}} \left(R_{g_{\infty}(t)} + \frac{1}{t} \right) dV_{g_{\infty}(t)} \frac{dt}{t}$$

$$= \int_{a}^{b} \int_{M_{\infty}} \left(R_{g_{\infty}(t)} - |\nabla u_{\infty}|^{2}(t) + \frac{1}{t} \right) dV_{g_{\infty}(t)} \frac{dt}{t}$$

$$\leq \lim_{i \to \infty} \int_{t_{i}a}^{t_{i}b} \int_{M} \left(R_{g(t)} - |\nabla u|^{2}(t) + \frac{1}{t} \right) dV_{g(t)} \frac{dt}{t}$$

$$= \lim_{i \to \infty} \left(\frac{V(t_{i}a)}{t_{i}a} - \frac{V(t_{i}b)}{t_{i}b} \right) = 0.$$
(2.73)

Since a and b were arbitrary, we obtain $R_{g_{\infty}(t)}(m) = -\frac{1}{t}$ for all $m \in M_{\infty}$ and $t \in (0, \infty)$. In particular,

$$-1 = R_{g_{\infty}(1)}(m_{\infty}) = \lim_{i \to \infty} R_{\hat{g}(t_i)}(m_i).$$
(2.74)

Equations (2.72) and (2.74) together contradict our assumptions about $\{t_i\}_{i=1}^{\infty}$ and $\{m_i\}_{i=1}^{\infty}$, thereby proving the proposition.

Remark 2.20. In the case $\chi(M) < 0$ one could hope for a bound diam $(M, g(t)) = O(\sqrt{t})$. With such a bound one could conclude that in Proposition 2.19, there is some $i_0 > 0$ such that for large t, $X_{i_0}(t)$ is all of M. Without such a bound, one could imagine that as $t \to \infty$, the manifolds $(M, \hat{g}(t))$ approach a family of surfaces of constant curvature $-\frac{1}{2}$ that slowly pinch off a closed geodesic.

3. Torus bundles

In this section we prove Theorem 1.6. In Subsection 3.1 we write down the Ricci flow equations with a $U(1) \times U(1)$ symmetry and give some direct consequences. In Subsection 3.2 we show that the Ricci flow exists for all $t \in [0, \infty)$. In Subsection 3.3 we prove that the curvature decays like $O(t^{-1})$. In Subsection 3.4 we show that the length of the circle base is $O(\sqrt{t})$. In Subsection 3.5 we finish the proof of Theorem 1.6.

3.1. Twisted principal $U(1) \times U(1)$ **bundles.** Let *N* be a an orientable 3-manifold which is the total space of a fiber bundle $\pi : N \to S^1$, with T^2 -fibers. Choosing an orientation of S^1 , the fiber bundle has a holonomy $H \in SL(2, \mathbb{Z}) = \pi_0(\text{Diff}^+(T^2))$. Taken up to inverses, *H* determines the topological type of the fiber bundle. We refer to [Sco83, Theorem 5.5] for the Thurston types of such fiber bundles. If *H* is elliptic, i.e. has finite order, then *N* has a flat structure. If *H* is parabolic, i.e. $|\text{Tr}(H)| \leq 2$ but *H* is not elliptic, then *N* has a Nil structure. If *H* is hyperbolic, i.e. has no eigenvalues on the unit circle, then *N* has a Sol structure.

If *H* is the identity then $N = S^1 \times T^2$ is the total space of a principal $U(1) \times U(1)$ bundle. That is, *N* admits a free $U(1) \times U(1)$ action. In general, *N* is the total space of a twisted principal $U(1) \times U(1)$ bundle, where "twisted" refers to the fact that *H* may be nontrivial. The setup is a special case of that in [Lot10, Section 4.1]. Let \mathcal{E} be a local system over S^1 of groups isomorphic to $U(1) \times U(1)$. We assume that the holonomy of the local system is $H \in Aut(U(1) \times U(1))$. Then there is a notion of a free \mathcal{E} -action on the T^2 -bundle *N*, which generalizes the global $U(1) \times U(1)$ action that exists when *H* is the identity.

Let *h* be a Riemannian metric on *N* which is \mathcal{E} -invariant. There is a corresponding horizontal distribution \mathcal{H} on *N*. Since \mathcal{H} is one-dimensional, it is integrable. Note that even if *H* is the identity, \mathcal{H} can have a nontrivial holonomy in $U(1) \times U(1)$, when going around the circle base. Thus the flat structure on \mathcal{E} is logically distinct from the flat structure on *N* coming from \mathcal{H} .

Let V be a coordinate chart of S^1 , with local coordinate y. The integrability of \mathcal{H} gives a local trivialization $V \times T^2$ of $\pi^{-1}(V)$. The restriction of h to $\pi^{-1}(V)$ is invariant under the $U(1) \times U(1)$ action coming from $\mathcal{E}|_V$. Using this action and the local trivialization, let x_1 , x_2 denote local angular coordinates on the T^2 -fibers. In terms of these coordinates we can write

$$h = \sum_{i,j=1}^{2} G_{ij}(y) dx^{i} dx^{j} + g_{yy}(y) dy^{2}.$$
 (3.1)

We use i, j, k, l for vertical indices. From [Lot10, Section 4.2], nonzero compo-

nents of the curvature tensor of (N, h) are

$$R_{ijkl}^{N} = -\frac{1}{4}g^{yy}G_{ik,y}G_{jl,y} + \frac{1}{4}g^{yy}G_{il,y}G_{jk,y},$$

$$R_{iyjy}^{N} = -\frac{1}{2}G_{ij;yy} + \frac{1}{4}G^{kl}G_{ik,y}G_{jl,y},$$
(3.2)

where

$$G_{ij;yy} = G_{ij,yy} - \Gamma_{yy}^{y}G_{ij,y} = G_{ij,yy} - \frac{1}{2}\frac{g_{yy,y}}{g_{yy}}G_{ij,y}.$$
 (3.3)

The nonzero components of the Ricci tensor are

$$R_{ij} = -\frac{1}{2}g^{yy}G_{ij;yy} - \frac{1}{4}g^{yy}G^{kl}G_{kl,y}G_{ij,y} + g^{yy}\frac{1}{2}G^{kl}G_{ik,y}G_{lj,y},$$

$$R_{yy} = -\frac{1}{2}G^{ij}G_{ij;yy} + \frac{1}{4}G^{ij}G_{jk,y}G^{kl}G_{li,y}.$$

The scalar curvature is

$$R = -g^{yy}G^{ij}G_{ij;yy} + \frac{3}{4}g^{yy}G^{ij}G_{jk,y}G^{kl}G_{li,y} - \frac{1}{4}g^{yy}G^{ij}G_{ij,y}G^{kl}G_{kl,y}.$$
 (3.4)

We will use matrix notation $G = (G_{ij})$. Note that G is symmetric and positivedefinite. The Ricci flow equation

$$\frac{dh}{dt} = -2\operatorname{Ric}_{h(t)} \tag{3.5}$$

preserves the local $U(1) \times U(1)$ invariance of the metric. In terms of g and G, it becomes

$$\frac{\partial g_{yy}}{\partial t} = \operatorname{Tr} \left(G^{-1} G_{;yy} \right) - \frac{1}{2} \operatorname{Tr} \left(\left(G^{-1} G_{,y} \right)^2 \right),$$

$$\frac{\partial G}{\partial t} = g^{yy} G_{;yy} + \frac{1}{2} g^{yy} \operatorname{Tr} \left(G^{-1} G_{,y} \right) G_{,y} - g^{yy} G_{,y} G^{-1} G_{,y}.$$
(3.6)

Adding a Lie derivative with respect to $-\nabla \ln \sqrt{\det(G)}$ to the right-hand side gives the modified equations

$$\frac{\partial g_{yy}}{\partial t} = \frac{1}{2} \operatorname{Tr} \left(\left(G^{-1} G_{,y} \right)^2 \right),
\frac{\partial G}{\partial t} = g^{yy} \left(G_{;yy} - G_{,y} G^{-1} G_{,y} \right).$$
(3.7)

Taking y to run over \mathbb{R} , the periodicity condition on G is

$$G(y+1,t) = H^T G(y,t) H.$$
 (3.8)

CMH

The manifold N can be recovered by taking the quotient of $\mathbb{R} \times T^2$ by the equivalence relation $(y + 1, x) \sim (y, Hx + b)$, where b is some fixed element of $\mathbb{R}^2/\mathbb{Z}^2$.

Hereafter we will mainly work with (3.7). An example of a solution to (3.7) is

$$g_{yy}(y,t) = 4c^{2}(t+a),$$

$$G(y,t) = \begin{pmatrix} e^{2cy} & 0\\ 0 & e^{-2cy} \end{pmatrix}.$$
(3.9)

If we take y to be defined in \mathbb{R}/\mathbb{Z} then we get a metric on a bundle with hyperbolic holonomy $H \in SL(2,\mathbb{Z})$, provided that H has eigenvalues e^c and e^{-c} . Here ∂_{x^1} and ∂_{x^2} are corresponding eigenvectors. The length of the circle base is $2c\sqrt{t+a}$.

Given s > 0 and a solution $(g(\cdot), G(\cdot))$ of (3.7), we obtain another solution of (3.7) by putting

$$g_s(t) = \frac{1}{s}g(st),$$

$$G_s(t) = G(st).$$
(3.10)

Put

$$h_s = (dx)^T G_s dx + g_s.$$
 (3.11)

Then

$$|\mathrm{Rm}_{h_s}|^2 = s^2 |\mathrm{Rm}_h|^2.$$
 (3.12)

Lemma 3.1.

$$\frac{\partial \ln \det(G)}{\partial t} = \Delta \ln \det(G). \tag{3.13}$$

Proof. We have

$$\frac{\partial \ln \det(G)}{\partial t} = \operatorname{Tr}\left(G^{-1}\frac{\partial G}{\partial t}\right)$$

$$= g^{yy}\operatorname{Tr}\left(G^{-1}(G_{;yy} - G_{,y}G^{-1}G_{,y})\right).$$
(3.14)

Passing to an arc-length parameter s on S^1 at a given time t, we obtain

$$\frac{\partial \ln \det(G)}{\partial t} = \operatorname{Tr}\left(G^{-1}(G_{,ss} - G_{,s}G^{-1}G_{,s})\right).$$
(3.15)

On the other hand

$$\Delta \ln \det(G) = \frac{d^2}{ds^2} \ln \det(G) = \frac{d}{ds} \operatorname{Tr} \left(G^{-1} G_{,s} \right)$$

$$= \operatorname{Tr} \left(G^{-1} G_{,ss} - G^{-1} G_{,s} G^{-1} G_{,s} \right).$$
(3.16)

This proves the lemma.

Note that det(G) is globally defined on S^1 since the holonomy lies in SL(2, \mathbb{Z}). It represents the squares of the volumes of the fibers.

Corollary 3.2. There are constants $C_1, C_2 > 0$ such that for all $y \in S^1$ and all times t for which the flow exists.

$$C_1 \le \sqrt{\det(G)}(y,t) \le C_2. \tag{3.17}$$

Proof. This follows from applying the maximum principle to Lemma 3.1. \Box

Lemma 3.3. Put

$$\mathcal{E} = g^{yy} \operatorname{Tr} \left((G^{-1} G_{,y})^2 \right).$$
 (3.18)

Then

$$\frac{\partial \mathcal{E}}{\partial t} = \Delta \mathcal{E} - \frac{\mathcal{E}^2}{2} - 2 g^{yy} g^{yy} \operatorname{Tr} \left(\left(G^{-1} G_{;yy} - (G^{-1} G_{,y})^2 \right)^2 \right).$$
(3.19)

Proof. Differentiating (3.18) with respect to t gives

$$\frac{\partial \mathcal{E}}{\partial t} = -\frac{1}{2} g^{yy} g^{yy} \left(\operatorname{Tr} \left((G^{-1}G_{,y})^2 \right) \right)^2
- 2g^{yy} \operatorname{Tr} \left(G^{-1}G_{,y}G^{-1}g^{yy}(G_{;yy} - G_{,y}G^{-1}G_{,y})G^{-1}G_{,y} \right)
+ 2g^{yy} \operatorname{Tr} \left(G^{-1}G_{,y}G^{-1} \left(g^{yy}(G_{;yy} - G_{,y}G^{-1}G_{,y}) \right)_{,y} \right).$$
(3.20)

Switching to an arc-length parameter s at a given time t, we obtain

$$\frac{\partial \mathcal{E}}{\partial t} = -\frac{\mathcal{E}^2}{2} - 2 \operatorname{Tr} \left(G^{-1} G_{,s} G^{-1} (G_{,ss} - G_{,s} G^{-1} G_{,s}) G^{-1} G_{,s} \right) + 2 \operatorname{Tr} \left(G^{-1} G_{,s} G^{-1} (G_{,ss} - G_{,s} G^{-1} G_{,s})_{,s} \right).$$
(3.21)

One now computes that

$$\frac{\partial \mathscr{E}}{\partial t} = -\frac{\mathscr{E}^2}{2} + \left(\operatorname{Tr} \left((G^{-1}G_{,s})^2 \right) \right)_{,ss} - 2\operatorname{Tr} \left(\left(G^{-1}G_{,ss} - (G^{-1}G_{,s})^2 \right)^2 \right).$$
(3.22)

This proves the lemma.

Note that

$$\mathcal{E} = g^{yy} \operatorname{Tr}\left(\left(G^{-\frac{1}{2}} G_{,y} G^{-\frac{1}{2}} \right)^2 \right)$$
(3.23)

is nonnegative, since $G^{-\frac{1}{2}}G_{,y}G^{-\frac{1}{2}}$ is symmetric.

CMH

Corollary 3.4. For all $y \in S^1$ and all $t \ge 0$ for which the flow exists, we have $\mathcal{E}(y,t) \le \frac{2}{t}$.

Proof. We have

$$\operatorname{Tr}\left(\left(G^{-1}G_{;yy} - (G^{-1}G_{,y})^{2}\right)^{2}\right) = \operatorname{Tr}\left(\left(G^{-\frac{1}{2}}G_{;yy}G^{-\frac{1}{2}} - \left(G^{-\frac{1}{2}}G_{,y}G^{-\frac{1}{2}}\right)^{2}\right)^{2}\right),$$
(3.24)

which is nonnegative since $G^{-\frac{1}{2}}G_{;yy}G^{-\frac{1}{2}} - (G^{-\frac{1}{2}}G_{,y}G^{-\frac{1}{2}})^2$ is symmetric. The corollary now follows from applying the maximum principle to Lemma 3.3.

3.2. Nonsingularity of the flow

Proposition 3.5. Given h(0) as described in Subsection 3.1, the flow (3.7) exists for $t \in [0, \infty)$.

Proof. If not then there is a singularity at some time $T < \infty$. Specializing the modified W-functional of [Lot10, Definition 4.48] to our case, it becomes

$$W(G, g, f, \tau)$$
(3.25)
= $\int_{S^1} \left[\tau |\nabla f|^2 - \frac{1}{4} g^{yy} \operatorname{Tr} \left(\left(G^{-1} G_{,y} \right)^2 \right) \right) + f - 1 \right] (4\pi\tau)^{-\frac{1}{2}} e^{-f} \sqrt{g_{yy}} dy.$

Using this modified W-functional, we can go through the same steps as in the proof of [Lis08, Theorem 7.9] to conclude that there is a blowup limit

$$(M_{\infty}, m_{\infty}, g_{\infty}(\cdot), G_{\infty}(\cdot))$$

where

• $G_{\infty} = \text{const.}$ and

• g_{∞} is a nonflat Ricci flow solution on the 1-manifold M_{∞} .

However, there is no such nonflat Ricci flow solution. This proves the proposition. \Box

3.3. Curvature bound

Proposition 3.6. We have $\max_{p \in N} |\text{Rm}^{N}|(p, t) = O(t^{-1})$.

Proof. Suppose that the proposition is false. We take a rescaling limit as in the proof of Proposition 2.10 to obtain a nonflat eternal solution $(M_{\infty}, p_{\infty}, g_{\infty}(\cdot), G_{\infty}(\cdot))$ on a one-dimensional étale groupoid M_{∞} . (It will follow from Lemma 3.12 that M_{∞} is a one-dimensional manifold.) From Corollary 3.4, G_{∞} is constant. Then g_{∞} is a nonflat Ricci flow solution on a one-dimensional space, which is a contradiction.

3.4. Diameter bound. We compute how the length of the base circle varies with time.

Lemma 3.7. Put $L(t) = \int_{S^1} \sqrt{g_{yy}}(y) dy$. Then

$$\frac{dL}{dt} = \frac{1}{4} \int_{S^1} \mathcal{E}(y, t) \sqrt{g_{yy}}(y) dy.$$
 (3.26)

Proof. We have

$$\frac{dL}{dt} = \frac{1}{2} \int_{S^1} g^{yy} \frac{\partial g_{yy}}{\partial t} \sqrt{g_{yy}}(y) dy$$

$$= \frac{1}{4} \int_{S^1} g^{yy} \operatorname{Tr}\left(\left(G^{-1} G_{,y} \right)^2 \right) \sqrt{g_{yy}}(y) dy.$$
(3.27)

This proves the lemma.

In particular, L(t) is monotonically nondecreasing in t.

Lemma 3.8. $t^{-\frac{1}{2}}L(t)$ is monotonically nonincreasing in t.

Proof. From Corollary 3.4 and Lemma 3.7, we derive

$$\frac{dL}{dt} \le \frac{L(t)}{2t}.$$
(3.28)

The lemma follows.

3.5. Long-time behavior

Lemma 3.9. Given $x \in T^2$ and linearly independent vectors $v_1, v_2 \in T_x T^2$, there is a constant $C(v_1, v_2) < \infty$ with the following property. Suppose that g_{T^2} is a flat metric on T^2 . Then

diam
$$(T^2, g_{T^2}) \le C(v_1, v_2) (|v_1|_{g_{T^2}} + |v_2|_{g_{T^2}}).$$
 (3.29)

Proof. Let \mathcal{V}_1 and \mathcal{V}_2 be the affine-parallel vector fields on T^2 that extend v_1 and v_2 , respectively. Let g_0 be a fixed flat metric on T^2 . There is some $c < \infty$ such that any pair of points in T^2 can be joined by flowing first in the \mathcal{V}_1 -direction for a length at most c and then flowing in the \mathcal{V}_2 -direction for a length at most c. With respect to g_{T^2} , the length of this path is bounded above by $c \frac{|v_1|_{\mathcal{S}_T^2}}{|v_1|_{\mathcal{S}_0}} + c \frac{|v_2|_{\mathcal{S}_T^2}}{|v_2|_{\mathcal{S}_0}}$. The lemma follows.

We use the notation in the statement of Theorem 1.6.

1	-	-	

3.5.1. Elliptic holonomy. If *H* is elliptic then after pulling back the T^2 -bundle from a finite covering $S^1 \to S^1$, we can assume that H = I. Then G(y, t) is a globally defined matrix-valued function of $y \in \mathbb{R}/\mathbb{Z}$ and $t \in [0, \infty)$.

Lemma 3.10. The intrinsic diameter of the T^2 -fibers is uniformly bounded above in t.

Proof. Given $v \in \mathbb{R}^2$, equation (3.7) implies that

$$\frac{\partial}{\partial t}\langle v, Gv \rangle = g^{yy} \langle v, Gv \rangle_{;yy} - g^{yy} \langle v, G_{,y} G^{-1} G_{,y} v \rangle.$$
(3.30)

The maximum principle now implies that $\langle v, G(y, t)v \rangle$ is uniformly bounded above for $y \in S^1$ and $t \in [0, \infty)$. The lemma follows from Lemma 3.9.

Proposition 3.11. diam $(N, h(t)) = O(\sqrt{t})$.

Proof. This follows from Lemmas 2.13, 3.8 and 3.10.

From [Lot10, Theorem 1.2.1],

$$\left(\max_{p \in N} \left| \operatorname{Rm}_{h}^{N} \right|(p,t) \right) \cdot \operatorname{diam}^{2}(N,h(t)) = o(t).$$
(3.31)

The argument of Subsection 2.5 now shows that $\lim_{t\to\infty} h(t)$ exists and is a flat metric on N, with the convergence being exponentially fast.

3.5.2. Hyperbolic holonomy. Suppose that *H* is hyperbolic.

Let $P(2, \mathbb{R})$ denote the positive-definite symmetric 2×2 matrices. Given $G \in P(2, \mathbb{R})$ and symmetric 2×2 matrices $\delta_1 G$, $\delta_2 G \in T_G P(2, \mathbb{R})$, we define their inner product by

$$\langle \delta_1 G, \delta_2 G \rangle = \frac{1}{2} \operatorname{Tr} \left(G^{-1}(\delta_1 G) G^{-1}(\delta_2 G) \right).$$
(3.32)

Consider the map $\Phi \colon \mathbb{R} \times SL(2, \mathbb{R}) \to P(2, \mathbb{R})$ given by $(u, M) \to e^u M^T M$. Identifying SO(2)\SL(2, \mathbb{R}) with the hyperbolic space H^2 , the map Φ passes to an isometry $\mathbb{R} \times H^2 \to P(2, \mathbb{R})$.

With respect to this isometry, the action of $SL(2, \mathbb{Z})$ on $P(2, \mathbb{R})$ (by $(A, G) \rightarrow A^T G A$) becomes the product of the trivial action of $SL(2, \mathbb{Z})$ on \mathbb{R} with the isometric action of $SL(2, \mathbb{Z})$ on H^2 . Letting $\langle H \rangle$ denote the cyclic subgroup of $SL(2, \mathbb{Z})$ generated by the holonomy H, equation (3.8) can be interpreted as saying that for each t, the function $G(\cdot, y)$ describes a smooth map $S^1 \rightarrow (\mathbb{R} \times H^2/\langle H \rangle)$ whose homotopy class is specified by H.

Lemma 3.12. There is a constant c > 0 such that for all $t \in [0, \infty)$, $L(t) \ge c\sqrt{t}$.

Proof. Lemma 3.4 says that the Lipschitz constant of the map $G(\cdot, t)$ from [0, 1] to $P(2, \mathbb{R})$ is bounded above by $\frac{1}{\sqrt{t}}$. As H is hyperbolic, there is a minimal length c > 0 among all noncontractible closed curves in $H^2/\langle H \rangle$. Consequently, the distance between G(0, t) and $G(1, t) = H^T G(0, t) H$ in $P(2, \mathbb{R})$ is bounded below by c. Thus $L(t) \ge c\sqrt{t}$.

Recall the definitions of g_s and G_s from (3.10). By an appropriate *s*-dependent choice of basis for \mathbb{R}^2 , we can assume that $G_s(0, 1) = I$. In making such a choice of basis, we are ignoring the lattice structure that comes from writing T^2 as a quotient of \mathbb{R}^2 . We are simply treating *G* and *g* as functions which satisfy (3.7) and (3.8).

Let X be the real symmetric matrix so that $e^X = H^T H$.

Proposition 3.13. For any sequence $\{s_j\}_{j=1}^{\infty}$ going to infinity, after passing to a subsequence and possibly reparametrizing S^1 , we have

$$\lim_{j \to \infty} g_{s_j}(y, t) = \frac{t}{2} \operatorname{Tr}(X^2) dy^2$$
(3.33)

and

$$\lim_{j \to \infty} G_{s_j}(y, t) = e^{yX}, \tag{3.34}$$

with smooth convergence on compact subsets of $S^1 \times [0, \infty)$.

Proof. The proof is similar to that in [Lot10, Propositions 4.39 and 4.79]; see also [FIN05, Theorem 1.3].

We first construct a positive solution \tilde{u}_{∞} of the conjugate heat equation

$$\frac{\partial u}{\partial t} = -\Delta u - \frac{1}{4} g^{yy} \operatorname{Tr} \left((G^{-1} G_{,y})^2 \right) u$$
(3.35)

that is defined for $t \in [0, \infty)$. To do so, note that if u is a solution to (3.35) then $\int_{S^1} u \sqrt{g_{yy}} dy$ is constant in t. Let $\{t_j\}_{j=1}^{\infty}$ be a sequence of times going to infinity. Let $\tilde{u}_j(\cdot)$ be a solution to (3.35) on the interval $[0, t_j]$ with initial condition $\tilde{u}_j(t_j) = \frac{1}{L(t_j)}$. One shows that one can extract a subsequence of the \tilde{u}_j 's that converges smoothly on compact subsets of $S^1 \times [0, \infty)$ to a positive solution $\tilde{u}_{\infty}(\cdot)$ of (3.35).

Define $\tilde{f}_{\infty}(\cdot)$ by $\tilde{u}_{\infty}(t) = (4\pi t)^{-\frac{n}{2}} e^{-\tilde{f}_{\infty}(t)}$. Put

$$W_{+}(G, g, \tilde{f}, t)$$

$$= \int_{S^{1}} \left[t \left(|\nabla \tilde{f}|^{2} - \frac{1}{4} g^{yy} \operatorname{Tr} \left((G^{-1} G_{,y})^{2} \right) \right) - \tilde{f} + 1 \right] (4\pi t)^{-\frac{1}{2}} e^{-\tilde{f}} \sqrt{g_{yy}} dy.$$
(3.36)

Vol. 89 (2014) Ricci flow on three-dimensional manifolds with symmetry

From [Lot10, Proposition 4.64],

$$\frac{d}{dt}W_{+}(G(t),g(t),\tilde{f}_{\infty}(t),t)$$

$$= \frac{t}{2} \int_{S^{1}} \operatorname{Tr}\left(\left(G^{-1}(\triangle G - g^{yy}G_{,y}G^{-1}G_{,y} - g^{yy}G_{,y}\tilde{f}_{\infty,y})\right)^{2}\right)\tilde{u}_{\infty}\sqrt{g_{yy}}\,dy$$

$$+ 2t \int_{S^{1}} g^{yy}g^{yy}\left(-\frac{1}{4}\operatorname{Tr}\left((G^{-1}G_{,y})^{2}\right) + \tilde{f}_{;yy} + \frac{1}{2t}g_{yy}\right)^{2}\tilde{u}_{\infty}\sqrt{g_{yy}}\,dy.$$
(3.37)

In particular, $W_+(G(t), g(t), \tilde{f}_{\infty}(t), t)$ is monotonically nondecreasing in t. Put

$$W_{\infty} = \lim_{t \to \infty} W_{+}(G(t), g(t), \tilde{f}_{\infty}(t), t),$$
 (3.38)

which at the moment could be infinity.

Using the curvature bound from Proposition 3.6 and the diameter bounds from Lemmas 3.8 and 3.12, one shows that after passing to a subsequence,

$$\lim_{j\to\infty}(g_{s_j}(\cdot),G_{s_j}(\cdot))$$

exists in the topology of smooth convergence on compact subsets of $S^1 \times [0, \infty)$, and equals a solution $(g_{\infty}(\cdot), G_{\infty}(\cdot))$ of (3.7) on a circle of time-1 length

$$L_{\infty} = \lim_{t \to \infty} \frac{L(t)}{\sqrt{t}}.$$
(3.39)

(The notion of convergence allows for j-dependent diffeomorphisms of S^1 .)

Put $u_j(t) = \tilde{u}_{\infty}(t + s_j)$. After passing to a subsequence, we can assume that $\lim_{j\to\infty} u_j(t) = u_{\infty}(t)$ for some solution $u_{\infty}(\cdot)$ to (3.35) (relative to $g_{\infty}(\cdot)$ and $G_{\infty}(\cdot)$), with smooth convergence on compact subsets of $S^1 \times [0, \infty)$. Define $f_{\infty}(\cdot)$ by $u_{\infty}(t) = (4\pi t)^{-\frac{n}{2}} e^{-f_{\infty}(t)}$. Then we have $W_+(G_{\infty}(t), g_{\infty}(t), f_{\infty}(t), t) = W_{\infty}$ for all t. In particular, $W_{\infty} < \infty$.

From (3.37), we obtain

$$\Delta G_{\infty} - g_{\infty}^{yy} G_{\infty,y} G_{\infty}^{-1} G_{\infty,y} - g_{\infty}^{yy} G_{\infty,y} f_{\infty,y} = 0, -\frac{1}{4} \operatorname{Tr} \left(\left(G_{\infty}^{-1} G_{\infty,y} \right)^2 \right) + f_{\infty;yy} + \frac{1}{2t} g_{\infty,yy} = 0.$$
(3.40)

From [Lot10, Proposition 4.80], there is a traceless symmetric matrix X such that $g_{\infty,yy}(y,t) = \frac{t}{2} \operatorname{Tr}(X^2)$ and $G_{\infty}(y,t) = e^{yX}$. Here y is a parametrization of S^1 whose time-1 velocity is L_{∞} . For each j, we had $G_{s_j}(y+1,t) = H^T G_{s_j}(y,t)H$. Hence $G_{\infty}(y+1,t) = H^T G_{\infty}(y,t)H$ and so $e^X = H^T H$.

We now prove part (ii) of Theorem 1.6. From Proposition 3.13, for any $K \in \mathbb{Z}^+$ and any $\epsilon > 0$, there is some $t_0 < \infty$ such that $\left(\frac{g_{yy}(y,t_0)}{t_0}, G(y,t_0)\right)$ is ϵ -close in the J. Lott and N. Sesum

 C^{K} -norm to $(\frac{1}{2}\text{Tr}(X^{2}), e^{yX})$. From the local stability result of [Kno09, Theorem 3], after an overall reparametrization of S^{1} , we have

$$\lim_{t \to \infty} \frac{g_{yy}(y,t)}{t} = \frac{1}{2} \operatorname{Tr}(X^2),$$

$$\lim_{t \to \infty} G(y,t) = e^{yX}.$$
(3.41)

The convergence is exponentially fast in the variable $\ln(t)$, i.e. power-law fast in t. (Strictly speaking, the result in [Kno09, Theorem 3] is for the modified Ricci flow (3.7) but there is a similar result for the unmodified Ricci flow (3.6).)

This proves Theorem 1.6.

Remark 3.14. Suppose that *H* is parabolic. By Corollary 3.2, the fiber volumes are uniformly bounded above and below by positive constants. After pulling back to a double cover of the base S^1 , if necessary, we can assume that Tr(H) = 2. If $v \in \mathbb{R}^2$ is a nonzero *H*-invariant vector then (3.30) gives a uniform upper bound on the squared length $\langle v, G(y, t)v \rangle$.

Lemma 3.7 implies that the length L(t) of the base circle is monotonically nondecreasing in t. We claim that $\lim_{t\to\infty} \frac{L(t)}{\sqrt{t}} = 0$. If not then we would conclude from the proof of Proposition 3.13 that N has a Sol-structure, which contradicts the topological fact that it has a Nil-structure.

If we knew that the diameters of the T^2 -fibers were $O(\sqrt{t})$ then we could conclude from Lemma 2.13 and [Lot10, Theorem 1.2.2] that the pullback Ricci flow solution on the universal cover \tilde{N} approaches the Nil expanding soliton. Based on the calculation in the locally homogeneous Nil case, as in [Lot07, Subsubsection 3.3.3], we expect that both L(t) and the fiber diameters are $O(t^{\frac{1}{6}})$.

References

- [Cao05] X. Cao, Isoperimetric estimate for the Ricci flow on $S^2 \times S^1$. Comm. Anal. Geom. 13 (2005), no. 4, 727–739. Zbl 1104.53065 MR 2191905
- [CFG92] J. Cheeger, K. Fukaya, and M. Gromov, Nilpotent structures and invariant metrics on collapsed manifolds. J. Amer. Math. Soc. 5 (1992), no. 2, 327–372. Zbl 0758.53022 MR 1126118
- [CG90] J. Cheeger and M. Gromov, Collapsing Riemannian manifolds while keeping their curvature bounded. II. J. Differential Geom. 32 (1990), no. 1, 269–298. Zbl 0727.53043 MR 1064875
- [CIJ90] M. Carfora, J. Isenberg, and M. Jackson, Convergence of the Ricci flow for metrics with indefinite Ricci curvature. J. Differential Geom. 31 (1990), no. 1, 249–263. Zbl 0647.53033 MR 1030672
- [CK04] B. Chow and D. Knopf, *The Ricci flow: an introduction*. Math. Surveys Monogr. 110, Amer. Math. Soc., Providence, RI, 2004. Zbl 1086.53085 MR 2061425

- [CLN06] B. Chow, P. Lu, and L. Ni, *Hamilton's Ricci flow*. Grad. Stud. Math. 77, Amer. Math. Soc., Providence, RI, 2006. Zbl 1118.53001 MR 2274812
- [CT06] J. Cheeger and G. Tian, Curvature and injectivity radius estimates for Einstein 4-manifolds. J. Amer. Math. Soc. 19 (2006), no. 2, 487–525. Zbl 1092.53034 MR 2188134
- [FIN05] M. Feldman, T. Ilmanen, and L. Ni, Entropy and reduced distance for Ricci expanders. J. Geom. Anal. 15 (2005), no. 1, 49–62. Zbl 1071.53040 MR 2132265
- [Fuk89] K. Fukaya, Collapsing Riemannian manifolds to ones with lower dimension. II. J. Math. Soc. Japan 41 (1989), no. 2, 333–356. Zbl 0703.53042 MR 0984756
- [GIK02] C. Guenther, J. Isenberg, and D. Knopf, Stability of the Ricci flow at Ricci-flat metrics. *Comm. Anal. Geom.* **10** (2002), no. 4, 741–777. Zbl 1028.53043 MR 1925501
- [Gro78] M. Gromov, Almost flat manifolds. J. Differential Geom. 13 (1978), no. 2, 231–241. Zbl 0432.53020 MR Almost flat manifolds
- [Ham95] R. S. Hamilton, The formation of singularities in the Ricci flow. In Surveys in differential geometry, Vol. II (Cambridge, Mass., 1993), International Press, Cambridge, Mass., 1995, 7–136. Zbl 0867.53030 MR 1375255
- [HI93] R. Hamilton and J. Isenberg, Quasi-convergence of Ricci flow for a class of metrics. Comm. Anal. Geom. 1 (1993), no. 3–4, 543–559. Zbl 0841.53058 MR 1266479
- [IJ92] J. Isenberg and M. Jackson, Ricci flow of locally homogeneous geometries on closed manifolds. J. Differential Geom. 35 (1992), no. 3, 723–741. Zbl 0808.53044 MR 1163457
- [KL08] B. Kleiner and J. Lott, Notes on Perelman's papers. Geom. Topol. 12 (2008), no. 5, 2587–2855. Zbl 1204.53033 MR 2460872
- [KM01] D. Knopf and K. McLeod, Quasi-convergence of model geometries under the Ricci flow. Comm. Anal. Geom. 9 (2001), no. 4, 879–919. Zbl 1020.53044 MR 1868923
- [Kno00] D. Knopf, Quasi-convergence of the Ricci flow. Comm. Anal. Geom. 8 (2000), no. 2, 375–391. Zbl 1020.53044 MR 1753322
- [Kno09] D. Knopf, Convergence and stability of locally ℝ^N-invariant solutions of Ricci flow. J. Geom. Anal. 19 (2009), no. 4, 817–846. Zbl 1176.53068 MR 2538938
- [Lis08] B. List, Evolution of an extended Ricci flow system. Comm. Anal. Geom. 16 (2008), no. 5, 1007–1048. Zbl 1166.53044 MR 2471366
- [Lot07] J. Lott, On the long-time behavior of type-III Ricci flow solutions. Math. Ann. 339 (2007), no. 3, 627–666. Zbl 1135.53046 MR 2336062
- [Lot10] J. Lott, Dimensional reduction and the long-time behavior of Ricci flow. Comment. Math. Helv. 85 (2010), no. 3, 485–534. Zbl 1195.53094 MR 2653690
- [Per02] G. Perelman, The entropy formula for the Ricci flow and its geometric applications. arXiv:math/0211159 [math.DG]
- [Ron07] X. Rong, Collapsed manifolds with bounded sectional curvature and applications. In Surveys in differential geometry, Vol. XI, International Press, Somerville, Mass., 2007, 1–23. Zbl 1155.53023 MR 2408262
- [Sco83] P. Scott, The geometries of 3-manifolds. Bull. London Math. Soc. 15 (1983), no. 5, 401–487. Zbl 0561.57001 MR 0705527

Received April 26, 2011

John Lott, Department of Mathematics, University of California - Berkeley, Berkeley, CA 94720-3840, U.S.A.

E-mail: lott@math.berkeley.edu

Natasa Sesum, Department of Mathematics, Rutgers University, Piscataway, NJ 08854-8019, U.S.A.

E-mail: natasas@math.rutgers.edu