

Simple length rigidity for Kleinian surface groups and applications

Autor(en): **Bridgeman, Martin / Canary, Richard D.**

Objekttyp: **Article**

Zeitschrift: **Commentarii Mathematici Helvetici**

Band (Jahr): **92 (2017)**

Heft 4

PDF erstellt am: **27.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-730826>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Simple length rigidity for Kleinian surface groups and applications

Martin Bridgeman* and Richard D. Canary*

Abstract. We prove that a Kleinian surface group is determined, up to conjugacy in the isometry group of \mathbb{H}^3 , by its simple marked length spectrum. As a first application, we show that a discrete faithful representation of the fundamental group of a compact, acylindrical, hyperbolizable 3-manifold M is similarly determined by the translation lengths of images of elements of $\pi_1(M)$ represented by simple curves on the boundary of M . As a second application, we show the group of diffeomorphisms of quasifuchsian space which preserve the renormalized pressure intersection is generated by the (extended) mapping class group and complex conjugation.

Mathematics Subject Classification (2010). 57M50.

Keywords. Kleinian surface groups, length spectrum and rigidity.

1. Introduction

We show that if ρ_1 and ρ_2 are two discrete, faithful representations of a surface group $\pi_1(S)$ into $\mathrm{PSL}(2, \mathbb{C})$ with the same simple marked length spectrum, then ρ_1 is either conjugate to ρ_2 or its complex conjugate. (Two such representations have the same simple marked length spectrum if whenever $\alpha \in \pi_1(S)$ is represented by a simple closed curve, then the images of α have the same translation length. The complex conjugate of a representation is obtained by conjugating the representation by $z \rightarrow \bar{z}$.) Marché and Wolff [21, Sec. 3] have exhibited non-elementary representations of a closed surface group of genus two into $\mathrm{PSL}(2, \mathbb{R})$ with the same simple marked length spectrum which do not have the same marked length spectrum, so the corresponding statement does not hold for non-elementary representations.

We give two applications of our main result. First, if M is a compact, acylindrical, hyperbolizable 3-manifold, we show that if ρ_1 and ρ_2 are discrete faithful representations of $\pi_1(M)$ into $\mathrm{PSL}(2, \mathbb{C})$ such that translation lengths of

*M.B. was partially supported by grant DMS-1500545 and R.C. was partially supported by grant DMS-1306992, from the National Science Foundation. The authors also acknowledge support from U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 “RNMS: GEometric structures And Representation varieties” (the GEAR Network).

the images of elements of $\pi_1(M)$ corresponding to simple curves in the boundary of M agree, then ρ_1 is either conjugate to ρ_2 or its complex conjugate. For our second application we consider the renormalized pressure intersection, first defined by Burger [9] and further studied by Bridgeman–Taylor [7]. Bridgeman [5] (see also [6]) showed that the Hessian of the renormalized pressure intersection gives rise to a path metric on quasifuchsian space $QF(S)$. We show that the group of diffeomorphisms of $QF(S)$ which preserve the renormalized pressure intersection is generated by the (extended) mapping class group and the involution of $QF(S)$ determined by complex conjugation.

1.1. Simple length rigidity for Kleinian surface groups. A *Kleinian surface group* is a discrete, faithful representation $\rho : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ where S is a closed, connected, orientable surface of genus at least two. If $\alpha \in \pi_1(S)$, then let $\ell_\rho(\alpha)$ denote the translation of length of $\rho(\alpha)$, or equivalently the length of the closed geodesic in the homotopy class of α in the quotient hyperbolic 3-manifold $\mathbb{H}^3/\rho(\pi_1(S))$. We say that two Kleinian surface groups $\rho_1 : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ have the *same marked length spectrum* if $\ell_{\rho_1}(\alpha) = \ell_{\rho_2}(\alpha)$ for all $\alpha \in \pi_1(S)$. Similarly, we say that ρ_1 and ρ_2 have the *same simple marked length spectrum* if $\ell_{\rho_1}(\alpha) = \ell_{\rho_2}(\alpha)$ whenever α has a representative on S which is a simple closed curve. If $\rho : G \rightarrow \mathrm{PSL}(2, \mathbb{C})$ is a representation we define its *complex conjugate* $\bar{\rho}_2$ to be the representation obtained by conjugating by $z \rightarrow \bar{z}$.

Theorem 1.1 (Simple length rigidity for Kleinian surface groups). *If S is a closed, connected, orientable surface of genus at least two, and $\rho_1 : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ are Kleinian surface groups with the same simple marked length spectrum, then ρ_1 is conjugate to either ρ_2 or $\bar{\rho}_2$.*

Since the full isometry group of \mathbb{H}^3 may be identified with the group generated by $\mathrm{PSL}(2, \mathbb{C})$, regarded as the group of fractional linear transformations, and $z \rightarrow \bar{z}$, one may reformulate our main result as saying that two Kleinian surface groups with the same simple marked length spectrum are conjugate in the isometry group of \mathbb{H}^3 .

Historical remarks. It is a classical consequence of the Fenchel–Nielsen coordinates for Teichmüller space that there are finitely many simple curves on S whose lengths determine a Fuchsian (i.e. discrete and faithful) representation of $\pi_1(S)$ into $\mathrm{PSL}(2, \mathbb{R})$ up to conjugacy in $\mathrm{PGL}(2, \mathbb{R})$, which we may identify with the isometry group of \mathbb{H}^2 . However, Marché and Wolff [21, Sec. 3] showed that there exist non-Fuchsian representations of the fundamental group of a surface of genus two into $\mathrm{PSL}(2, \mathbb{R})$ with the same simple marked length spectrum which do not have the same marked length spectrum. The representations constructed by Marché and Wolff do not lift to $\mathrm{SL}(2, \mathbb{R})$, so do not lie in the same component of the $\mathrm{PSL}(2, \mathbb{C})$ -representation variety as the discrete faithful representations.

Kourounitis [19] showed that there are finitely many simple curves on S whose complex lengths (see Section 2 for a discussion of complex length) determine a quasifuchsian surface group up to conjugacy in $\mathrm{PSL}(2, \mathbb{C})$. Culler and Shalen [13, Prop. 1.4.1] showed that there are finitely many curves whose traces determine a non-elementary representation into $\mathrm{SL}(2, \mathbb{C})$, up to conjugacy in $\mathrm{SL}(2, \mathbb{C})$, while Charles–Marché [11, Thm. 1.1] showed that one may choose the finite collection to consist of simple closed curves.

Kim [18] previously showed that two Zariski dense representations into the isometry group $\mathrm{Isom}(X)$ of a rank one symmetric space X with the same full marked length spectrum are conjugate in $\mathrm{Isom}(X)$. More generally, Dal’Bo and Kim [14] showed that any surjective homomorphism between Zariski dense subgroups of a semi-simple Lie group G , with trivial center and no compact factors, which preserves translation length on the associated symmetric space extends to an automorphism of G .

1.2. Simple length rigidity for acylindrical hyperbolic 3-manifolds. A compact, orientable 3-manifold M with non-empty boundary is said to be *hyperbolizable* if its interior admits a complete hyperbolic metric, which implies that there exists a discrete, faithful representation of $\pi_1(M)$ into $\mathrm{PSL}(2, \mathbb{C})$. A compact, hyperbolizable 3-manifold is said to be *acylindrical* if every π_1 -injective proper map of an annulus into M is properly homotopic into the boundary of M . (Recall that a map of a surface into a 3-manifold is said to be proper if it maps the boundary of the surface into the boundary of 3-manifold and that a proper homotopy is a homotopy through proper maps.)

In this setting, we use Theorem 1.1 show that a discrete, faithful representation of $\pi_1(M)$ into $\mathrm{PSL}(2, \mathbb{C})$ is determined, up to conjugacy in the isometry group of \mathbb{H}^3 , by the translation lengths of images of simple curves in the boundary ∂M of M .

Theorem 1.2. *If M is a compact, acylindrical, hyperbolizable 3-manifold, and $\rho_1 : \pi_1(M) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(M) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ are two discrete faithful representations, such that $\ell_{\rho_1}(\alpha) = \ell_{\rho_2}(\alpha)$ if $\alpha \in \pi_1(M)$ is represented by a simple closed curve on ∂M , then ρ_1 is conjugate to either ρ_2 or $\bar{\rho}_2$.*

1.3. Isometries of the renormalized pressure intersection. Burger [9] introduced a renormalized pressure intersection between convex cocompact representations into rank one Lie groups. Bridgeman and Taylor [7] extensively studied this renormalized pressure intersection in the setting of quasifuchsian representation. We say that $\rho : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ is *quasifuchsian* if it is topologically conjugate, in terms of its action on $\widehat{\mathbb{C}}$, to a Fuchsian representation into $\mathrm{PSL}(2, \mathbb{R})$. If $T > 0$ we let

$$R_T(\rho) = \{[\alpha] \in [\pi_1(S)] \mid \ell_\rho(\alpha) \leq T\}$$

where $[\pi_1(S)]$ is the set of conjugacy classes in $\pi_1(S)$. We define the *entropy*

$$h(\rho) = \limsup \frac{\log(\#(R_T(\rho)))}{T}$$

of a quasifuchsian representation ρ . Sullivan [32] showed that $h(\rho)$ is the Hausdorff dimension of the limit set of $\rho(\pi_1(S))$.

Let $QF(S)$ denote the space of $\mathrm{PSL}(2, \mathbb{C})$ -conjugacy classes of quasifuchsian representations. Bers [1] showed that $QF(S)$ is an analytic manifold which may be naturally identified with $\mathcal{T}(S) \times \mathcal{T}(S)$. If $\rho_1, \rho_2 \in QF(S)$, the *renormalized pressure intersection* of ρ_1 and ρ_2 is given by

$$\mathbf{J}(\rho_1, \rho_2) = \frac{h(\rho_2)}{h(\rho_1)} \lim_{T \rightarrow \infty} \left(\frac{1}{\#(R_T(\rho_1))} \sum_{[\alpha] \in R_T(\rho_1)} \frac{\ell_{\rho_2}(\alpha)}{\ell_{\rho_1}(\alpha)} \right).$$

Bridgeman and Taylor [7] showed that the Hessian of J gives rise to a non-negative bilinear form on the tangent space $TQF(S)$ of quasifuchsian space, called the pressure form. Motivated by work of McMullen [24] in the setting of Teichmüller space, Bridgeman [5] used the thermodynamic formalism to show that the only degenerate vectors for the pressure form correspond to pure bending at points on the Fuchsian locus. Moreover, the pressure form gives rise to a path metric on $QF(S)$, called the *pressure metric* (see also [6, Cor. 1.7]).

We say a smooth immersion $f : QF(S) \rightarrow QF(S)$ is a *smooth isometry of the renormalized pressure intersection* if

$$\mathbf{J}(f(\rho_1), f(\rho_2)) = \mathbf{J}(\rho_1, \rho_2)$$

for all $\rho_1, \rho_2 \in QF(S)$. We recall that the (extended) mapping class group $\mathrm{Mod}^*(S)$ is the group of isotopy classes of homeomorphisms of S . Since J is invariant under the action of $\mathrm{Mod}^*(S)$, every element of $\mathrm{Mod}^*(S)$ is a smooth isometry of the renormalized pressure intersection. There exists an involution $\tau : QF(S) \rightarrow QF(S)$ given by taking $[\rho]$ to $[\bar{\rho}]$. Since τ preserves the marked length spectrum, it is an isometry of the renormalized pressure intersection. We use our main result and work of Bonahon [4] to show that these give rise to all smooth isometries of the renormalized pressure intersection.

Theorem 1.3. *If S is a closed, orientable surface of genus at least two, then the group of smooth isometries of the renormalized pressure intersection on $QF(S)$ is generated by the (extended) mapping class group $\mathrm{Mod}^*(S)$ and complex conjugation τ .*

Royden [29] showed that $\mathrm{Mod}^*(S)$ is the isometry group of the Teichmüller metric on $T(S)$. Masur and Wolf [23] proved that $\mathrm{Mod}^*(S)$ is the isometry group of the Weil–Petersson metric on $T(S)$. Bridgeman [5] used work of Wolpert [33] to show that the restriction of the pressure form to the Fuchsian locus is a multiple of the Weil–Petersson metric.

One may thus view Theorem 1.3 as evidence in favor of the following natural conjecture.

Conjecture. *The isometry group of the pressure metric on quasifuchsian space $QF(S)$ is generated by the (extended) mapping class group and complex conjugation.*

In the proof of Theorem 1.3, we establish the following strengthening of our main result which may be of independent interest.

Theorem 1.4. *If S is a closed, connected, orientable surface of genus at least two, $\rho_1 : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ are Kleinian surface groups, and there exists k so that $\ell_{\rho_1}(\alpha) = k\ell_{\rho_2}(\alpha)$ for all $\alpha \in \pi_1(S)$ which are represented by simple curves on S , then ρ_1 is conjugate to either ρ_2 or $\bar{\rho}_2$.*

Kim [18, Thm. 3] showed that if ρ_1 and ρ_2 are irreducible, non-elementary, nonparabolic representations of a finitely presented group Γ into the isometry group of a rank one symmetric space and there exists $k > 0$ such that $\ell_{\rho_1}(\gamma) = k\ell_{\rho_2}(\gamma)$ for all $\gamma \in \Gamma$ (where $\ell_{\rho_i}(\gamma)$ the translation length of $\rho_i(\gamma)$), then $k = 1$ and ρ_1 and ρ_2 are conjugate representations.

Outline of paper. In Section 2 we analyze the complex length spectrum of Kleinian surface groups with the same simple marked length spectrum, then in Section 3, we give the proof of our main result. In Section 4 we prove Theorem 1.2, while in Section 5 we establish Theorems 1.3 and 1.4.

Acknowledgements. The authors would like to thank Maxime Wolff for several enlightening conversations on the length spectra of surface group representations, Jeff Brock and Mike Wolf for conversations about the Weil–Petersson metric and Alan Reid for conversations about the character variety. This material is partially based upon work supported by the National Science Foundation under grant No. 0932078 000 while the authors were in residence at the Mathematical Sciences Research Institute in Berkeley, CA, during the Spring 2015 semester.

2. The complex length spectrum

In this section, we investigate the complex length spectra of Kleinian surface groups with the same simple marked length spectrum.

Given $\alpha \in \pi_1(S)$ and $\rho \in AH(S)$, let $\lambda_\rho^2(\alpha)$ be the square of the largest eigenvalue of $\rho(\alpha)$. Notice that $\lambda_\rho^2(\alpha)$ is well-defined even though the largest eigenvalue of a matrix in $\mathrm{PSL}(2, \mathbb{C})$ is only well-defined up to sign. If we choose $\log \lambda_\rho^2(\alpha)$ to have imaginary part in $[0, 2\pi)$, then $\log \lambda_\rho^2(\alpha)$ is the *complex length* of $\rho(\alpha)$.

If α is a simple, non-separating closed curve on S , we let $W(\alpha)$ denote the set of all simple, non-separating curves on S which intersect α at most once. We say that ρ_1

and ρ_2 have the same marked complex length spectrum on $W(\alpha)$ if $\lambda_{\rho_1}^2(\beta) = \lambda_{\rho_2}^2(\beta)$ for all $\beta \in W(\alpha)$. Similarly, we say that ρ_1 and ρ_2 have conjugate marked complex length spectrum on $W(\alpha)$ if $\lambda_{\rho_1}^2(\beta) = \overline{\lambda_{\rho_2}^2(\beta)}$ for all $\beta \in W(\alpha)$.

We will show that if two Kleinian surface groups ρ_1 and ρ_2 have the same simple marked length spectrum, then, there exists a simple non-separating curve α on S such that ρ_1 and ρ_2 either have the same or conjugate complex length spectrum on $W(\alpha)$.

Proposition 2.1. *If S is a closed, connected, orientable surface of genus at least two, $\rho : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ are Kleinian surface groups with the same simple marked length spectrum, then there exists a simple non-separating curve α on S such that $\rho_1(\alpha)$ is hyperbolic and either*

- (1) ρ_1 and ρ_2 have the same marked complex length spectrum on $W(\alpha)$, or
- (2) ρ_1 and ρ_2 have conjugate marked complex length spectrum on $W(\alpha)$.

Proposition 2.1 will be a nearly immediate consequence of three lemmas. The first lemma shows that for two Kleinian surface groups with the same length spectrum, then the complex lengths of a simple non-separating curve either agree, differ by complex conjugation, or differ by sign (and are both real). The second lemma deals with the case where the complex length of every simple, non-separating curve is real, while the final lemma handles the case where some complex length is not real. All the proofs revolve around an analysis of the asymptotic behavior of complex lengths of curves of the form $\alpha^n \beta$ where α and β intersect exactly once. We begin by recording computations which will be used repeatedly in the remainder of the paper.

2.1. A convenient normalization. We recall that two elements $\alpha, \beta \in \pi_1(S)$ are *coprime* if they share no common powers. We say that a representation $\rho : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ is (α, β) -normalized if $\alpha, \beta \in \pi_1(S)$ are coprime and $\rho(\alpha)$ is hyperbolic and has attracting fixed point ∞ and repelling fixed point 0. In this case,

$$\rho(\alpha) = \pm \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}$$

where $|\lambda| > 1$, and

$$\rho(\beta) = \pm \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

where $ad - bc = 1$. Notice that the matrix representations of elements of $\mathrm{PSL}(2, \mathbb{C})$ are only well-defined up to multiplication by $\pm I$, but many related quantities like the square of the trace, the product of any two co-efficients, and the modulus of the eigenvalue of maximal modulus are well-defined.

Lemma 2.2. *Suppose that S is a closed, connected, orientable surface of genus at least two and $\rho : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ is an (α, β) -normalized Kleinian surface*

group. In the above notation,

$$\rho(\alpha^n \beta) = \pm \begin{pmatrix} \lambda^n a & \lambda^n b \\ \lambda^{-n} c & \lambda^{-n} d \end{pmatrix}$$

and all the matrix coefficients of $\rho(\beta)$ are non-zero. Moreover, if $\mu(n)$ is the modulus of the eigenvalue of $\rho(\alpha^n \beta)$ with largest modulus, then

$$\log \mu(n) = n \log |\lambda| + \log |a| + \Re \left(\lambda^{-2n} \frac{bc}{a^2} \right) + O(|\lambda|^{-4n}).$$

Proof. The first claim follows from a simple computation. If any of the coefficients of $\rho(\beta)$ are 0, then $\rho(\beta)$ takes some fixed point of $\rho(\alpha)$ to a fixed point of $\rho(\alpha)$, e.g. if $a = 0$, then $\rho(\beta)(\infty) = 0$. This would imply that $\rho(\beta\alpha\beta^{-1})$ shares a fixed point with $\rho(\alpha)$. Since $\rho(\pi_1(S))$ is discrete, this would imply that there is an element which is a power of both $\rho(\alpha)$ and $\rho(\beta\alpha\beta^{-1})$ which would contradict the facts that ρ is faithful and the subgroup of $\pi_1(S)$ generated by α and β is free of rank two.

The eigenvalues of $\rho(\alpha^n \beta)$, which are only well-defined up to sign, are then given by

$$\pm \left(\frac{(\lambda^n a + \lambda^{-n} d) \pm \sqrt{(\lambda^n a + \lambda^{-n} d)^2 - 4}}{2} \right)$$

So, since $|\lambda| > 1$, for all large enough n , one may use the Taylor expansion for $\sqrt{1+x}$ to conclude that they have the form

$$\pm \left(\lambda^n a \left(1 + \lambda^{-2n} \left(\frac{ad-1}{a^2} \right) + O(\lambda^{-4n}) \right) \right)$$

Therefore, since $ad - bc = 1$,

$$\log \mu(n) = n \log |\lambda| + \log |a| + \log \left| 1 + \lambda^{-2n} \frac{bc}{a^2} + O(\lambda^{-4n}) \right|.$$

We then use the expansion of $\log |1+z|$ about $z=0$ given by

$$\log |1+z| = \frac{1}{2} \log (|1+z|^2) = \frac{1}{2} \log (1 + 2\Re(z) + |z|^2) = \Re(z) + O(|z|^2)$$

to show that

$$\log \mu(n) = n \log |\lambda| + \log |a| + \Re \left(\lambda^{-2n} \frac{bc}{a^2} \right) + O(|\lambda|^{-4n}). \quad \square$$

2.2. Basic relationships between complex lengths. Our first lemma shows that if two Kleinian surface groups have the same simple marked length spectrum, then the complex lengths of any simple, non-separating curve either agree or differ by either complex conjugation or sign.

Lemma 2.3. *If ρ_1 and ρ_2 in $AH(S)$ have the same simple marked length spectrum and α is a simple non-separating curve on S , then either*

- (1) $\lambda_{\rho_1}^2(\alpha) = \lambda_{\rho_2}^2(\alpha)$,
- (2) $\lambda_{\rho_1}^2(\alpha) = \overline{\lambda_{\rho_2}^2(\alpha)}$, or
- (3) $\lambda_{\rho_1}^2(\alpha) = -\lambda_{\rho_2}^2(\alpha)$ and $\lambda_{\rho_1}^2(\alpha)$ is real.

Proof. If $\rho_1(\alpha)$ is parabolic, then $\rho_2(\alpha)$ is parabolic (since $\ell_{\rho_2}(\alpha) = \ell_{\rho_1}(\alpha) = 0$). In this case, $\lambda_{\rho_1}^2(\alpha) = \lambda_{\rho_2}^2(\alpha) = 1$. So we may assume that $\rho(\alpha)$ is hyperbolic.

Let β be a curve intersecting α exactly once. We may assume that both ρ_1 and ρ_2 are (α, β) -normalized, so

$$\rho_i(\alpha) = \pm \begin{pmatrix} \lambda_i & 0 \\ 0 & \lambda_i^{-1} \end{pmatrix}$$

where $|\lambda_i| > 1$, and

$$\rho_i(\beta) = \pm \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix},$$

where $a_i d_i - b_i c_i = 1$.

Since, ρ_1 and ρ_2 have the same simple marked length spectrum and $|\lambda_1| = |\lambda_2|$, Lemma 2.2 implies that

$$\log |a_1| + \Re \left(\lambda_1^{-2n} \frac{b_1 c_1}{a_1^2} \right) + O(|\lambda_1|^{-4n}) = \log |a_2| + \Re \left(\lambda_2^{-2n} \frac{b_2 c_2}{a_2^2} \right) + O(|\lambda_2|^{-4n})$$

for all n . Taking limits as $n \rightarrow \infty$, we see that $\log |a_1| = \log |a_2|$, so

$$\Re \left(\lambda_1^{-2n} \frac{b_1 c_1}{a_1^2} \right) + O(|\lambda_1|^{-4n}) = \Re \left(\lambda_2^{-2n} \frac{b_2 c_2}{a_2^2} \right) + O(|\lambda_2|^{-4n})$$

for all n . Therefore, after multiplying both sides by $|\lambda_1|^{2n} = |\lambda_2|^{2n}$, we see that

$$\lim_{n \rightarrow \infty} \Re \left(u_1^n v_1 - u_2^n v_2 \right) = 0$$

where

$$u_i = \left(\frac{\lambda_i}{|\lambda_i|} \right)^{-2} \quad \text{and} \quad v_i = \frac{b_i c_i}{a_i^2} \neq 0.$$

Lemma 2.3 is then an immediate consequence of the following elementary lemma. □

Lemma 2.4. *If $u_1, u_2 \in \mathbb{S}^1$, $v_1, v_2 \in \mathbb{C} - \{0\}$ and*

$$\lim_{n \rightarrow \infty} \Re(u_1^n v_1 - u_2^n v_2) = 0,$$

then either

- (1) $u_1 = u_2$,
- (2) $u_1 = \bar{u}_2$, or
- (3) $u_1 = -u_2 = \pm 1$.

Proof. We choose θ_i so that

$$u_i = e^{i\theta_i}.$$

If $s = \{n_k\}_{k=1}^\infty$ is a strictly increasing sequence of integers, let $S_i(s)$ be the set of accumulation points of $\{\Re(u_i^{n_k} v_i)\}$. Then, by assumption, $S_1(s) = S_2(s)$ for any sequence s .

If θ_i is an irrational multiple of 2π , then $S_i(\mathbb{N})$ is the interval $[-|v_i|, |v_i|]$. If θ_i is a rational multiple of 2π then $S_i(\mathbb{N})$ is finite. Therefore either (a) both θ_1 and θ_2 are irrational with $|v_1| = |v_2|$ or (b) both θ_1 and θ_2 are rational multiples of 2π . We handle these two cases separately.

Case I. Both θ_1 and θ_2 are irrational multiples of 2π and $|v_1| = |v_2|$. Since θ_1 is an irrational multiple of 2π , there is a sequence $\{n_k\}$ such that $\lim_{k \rightarrow \infty} e^{in_k \theta_1} = \frac{\bar{v}_1}{|v_1|}$. Therefore,

$$|v_1| = \lim_{k \rightarrow \infty} \Re(e^{in_k \theta_1} v_1) = \lim_{k \rightarrow \infty} \Re(e^{in_k \theta_2} v_2) = |v_2|,$$

so $\lim_{k \rightarrow \infty} e^{in_k \theta_2} = \frac{\bar{v}_2}{|v_2|}$. If $\{m_k\} = \{n_k + 1\}$, then

$$|v_1| \cos \theta_1 = \lim_{k \rightarrow \infty} \Re(e^{im_k \theta_1} v_1) = \lim_{k \rightarrow \infty} \Re(e^{im_k \theta_2} v_2) = |v_2| \cos \theta_2.$$

Since $|v_1| = |v_2| \neq 0$, it follows that $\theta_1 = \pm \theta_2$, so either $u_1 = u_2$ or $u_1 = \bar{u}_2$ and we are either in case (1) or in case (2).

Case II. Both θ_1 and θ_2 are rational multiples of 2π . Let $\theta_i = 2\pi p_i/q_i$ where $0 \leq p_i < q_i$ and p_i and q_i are relatively prime (and $q_i = 1$ if $p_i = 0$).

If $r \in \mathbb{Z}$ and $s_r = \{r + kq_1q_2\}$, then

$$S_1(s_r) = \{\Re(u_1^r v_1)\} = S_2(s_r) = \{\Re(u_2^r v_2)\},$$

so

$$\Re(u_1^r v_1) = \Re(u_2^r v_2) \quad \text{for all } r \in \mathbb{Z}.$$

If $\Re(u_1^r v_1) = \Re(u_2^r v_2) = 0$ for all $r \in \mathbb{Z}$, then $u_1 = \pm u_2 = \pm 1$, and we are in either case (1) or case (3).

If $\Re(u_1^r v_1) = \Re(u_2^r v_2) \neq 0$ for some r , then

$$\begin{aligned} 2 \cos(\theta_1) \Re(u_1^r v_1) &= \Re(u_1^{r+1} v_1) + \Re(u_1^{r-1} v_1) \\ &= \Re(u_2^{r+1} v_2) + \Re(u_2^{r-1} v_2) = 2 \cos(\theta_2) \Re(u_2^r v_2). \end{aligned}$$

Since $\Re(u_1^r v_1) = \Re(u_2^r v_2) \neq 0$, this implies that $\cos(\theta_1) = \cos(\theta_2)$, so $\theta_1 = \pm\theta_2$. Therefore, either $u_1 = u_2$ or $u_1 = \bar{u}_2$ and we are in either case (1) or case (2).

This completes the proof, since in all situations we have seen that either case (1), (2) or (3) occurs. \square

2.3. When the simple non-separating complex length spectrum is totally real.

We use a similar analysis to show that if the complex lengths of every simple, non-separating curve is real for two Kleinian surface groups with the same simple marked length spectrum, then the complex lengths agree for every simple, non-separating curve.

Lemma 2.5. *If S is a closed, connected, orientable surface of genus at least two and $\rho_1 : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ are Kleinian surface groups with same simple marked length spectrum, then either*

- (1) *there exists a simple non-separating curve γ on S such that $\lambda_{\rho_1}^2(\gamma) \notin \mathbb{R}$, or*
- (2) *$\lambda_{\rho_1}^2(\gamma) = \lambda_{\rho_2}^2(\gamma) \in \mathbb{R}$ whenever γ is a simple non-separating curve on S .*

Proof. Suppose that (1) does not hold, so $\lambda_{\rho_1}^2(\gamma) \in \mathbb{R}$ whenever γ is a simple non-separating curve on S . Lemma 2.3 then implies that $\lambda_{\rho_1}^2(\gamma) = \pm\lambda_{\rho_2}^2(\gamma)$ whenever γ is a simple non-separating curve on S .

Suppose that there is a simple non-separating curve α such that $\lambda_{\rho_1}^2(\alpha) = -\lambda_{\rho_2}^2(\alpha)$. Notice that if $\rho_1(\alpha)$ is parabolic, then $\ell_{\rho_1}(\alpha) = 0 = \ell_{\rho_2}(\alpha)$, so $\lambda_{\rho_1}^2(\alpha) = 1 = \lambda_{\rho_2}^2(\alpha)$. Therefore, $\rho_1(\alpha)$, and hence $\rho_2(\alpha)$, must be hyperbolic.

We choose a simple non-separating curve β intersecting α exactly once. We adapt the normalization and notation of Lemma 2.3. Lemma 2.2 implies that

$$\begin{aligned} t_i(n) &= \mathrm{Tr}^2(\rho_i(\alpha^n \beta)) = \lambda_i^{2n} a_i^2 + 2a_i d_i + \lambda_i^{-2n} d_i^2 \\ &= \lambda_{\rho_i}^2(\alpha^n \beta) + 2 + \lambda_{\rho_i}^{-2}(\alpha^n \beta). \end{aligned}$$

(Notice that the trace $\mathrm{Tr}(\rho_i(\alpha^n \beta))$ of $\rho_i(\alpha^n \beta)$ is well-defined up to sign, so $\mathrm{Tr}^2(\rho_i(\alpha^n \beta))$ is well-defined.) Since $\lambda_{\rho_1}^2(\alpha^n \beta) = \pm\lambda_{\rho_2}^2(\alpha^n \beta)$ for all n , by assumption, either (i) $t_1(n) = t_2(n)$ or (ii) $t_1(n) = 4 - t_2(n)$ for all n and $\lambda_1^2 = -\lambda_2^2$.

The proof divides into two cases.

Case I. There is an infinite sequence $\{n_k\}$ of even integers so that $t_1(n_k) = t_2(n_k)$.

Dividing by $\lambda_1^{2n_k} = \lambda_2^{2n_k}$ and taking limits we see that

$$a_1^2 = \lim_{k \rightarrow \infty} a_1^2 + \frac{2a_1 d_1}{\lambda_1^{2n_k}} + \frac{d_1^2}{\lambda_1^{4n_k}} = \lim_{k \rightarrow \infty} a_2^2 + \frac{2a_2 d_2}{\lambda_2^{2n_k}} + \frac{d_2^2}{\lambda_2^{4n_k}} = a_2^2.$$

It follows that

$$2a_1 d_1 + \frac{d_1^2}{\lambda_1^{2n_k}} = 2a_2 d_2 + \frac{d_2^2}{\lambda_2^{2n_k}}$$

for all n_k , so, after again taking limits, we see that

$$a_1 d_1 = a_2 d_2.$$

Then, by considering the final term, we see that

$$d_1^2 = d_2^2.$$

If there exists an infinite sequence $\{m_j\}$ of odd integers so that $t_1(m_j) = t_2(m_j)$ for all m_j , then

$$\lambda_1^{2m_j} a_1^2 + 2a_1 d_1 + \lambda_1^{-2m_j} d_1^2 = -\lambda_1^{2m_j} a_1^2 + 2a_1 d_1 - \lambda_1^{-2m_j} d_1^2.$$

Then we may divide each side by $\lambda_1^{2m_j}$ and pass to a limit to conclude that $a_1^2 = -a_1^2$. This would imply that $a_1 = 0$, which would contradict Lemma 2.2.

On the other hand, if there exists an infinite sequence $\{m_j\}$ of odd integers so that $t_1(m_j) = 4 - t_2(m_j)$ for all m_j , then

$$\lambda_1^{2m_j} a_1^2 + 2a_1 d_1 + \lambda_1^{-2m_j} d_1^2 = 4 - (-\lambda_1^{2m_j} a_1^2 + 2a_1 d_1 - \lambda_1^{-2m_j} d_1^2),$$

so

$$2a_1 d_1 = 4 - 2a_1 d_1.$$

Therefore, $a_1 d_1 = 1$, which implies that $b_1 c_1 = 0$, so either $b_1 = 0$ or $c_1 = 0$, which again contradicts Lemma 2.2.

Case II. There is an infinite sequence $\{n_k\}$ of even integers so that $t_1(n_k) = 4 - t_2(n_k)$.

We then argue, as in Case I, to show that

$$a_1^2 = -a_2^2, \quad 2a_1 d_1 = 4 - 2a_2 d_2 \quad \text{and} \quad d_1^2 = -d_2^2.$$

If there exists an infinite sequence $\{m_j\}$ of odd integers so that $t_1(m_j) = t_2(m_j)$ for all m_j , then

$$\lambda_1^{2m_j} a_1^2 + 2a_1 d_1 + \lambda_1^{-2m_j} d_1^2 = \lambda_1^{2m_j} a_1^2 + (4 - 2a_1 d_1) + \lambda_1^{-2m_j} d_1^2.$$

So, $a_1 d_1 = 1$, again giving a contradiction.

On the other hand, if there exists an infinite sequence $\{m_j\}$ of odd integers so that $t_1(m_j) = 4 - t_2(m_j)$ for all m_j , then

$$\lambda_1^{2m_j} a_1^2 + 2a_1 d_1 + \lambda_1^{-2m_j} d_1^2 = 4 - (\lambda_1^{2m_j} a_1^2 + (4 - 2a_1 d_1) + \lambda_1^{-2m_j} d_1^2).$$

Dividing both sides by $\lambda_1^{2m_j}$ and passing to a limit, we conclude that $a_1^2 = -a_1^2$, which is again a contradiction.

Therefore, neither Case I or Case II can occur, so case (2) must hold. \square

2.4. When the complex length is not always totally real. We now show that if $\lambda_{\rho_1}^2(\alpha)$ is not real, for some simple non-separating curve α , then $\lambda_{\rho_1}^2(\beta)$ and $\lambda_{\rho_2}^2(\beta)$ either agree for all $\beta \in W(\alpha)$, or differ by complex conjugation for all $\beta \in W(\alpha)$.

Lemma 2.6. *Suppose that S is a closed, connected, orientable surface of genus at least two and $\rho_1 : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ are Kleinian surface groups with same simple marked length spectrum. If α is simple non-separating curve on S such that $\lambda_{\rho_1}^2(\alpha) \notin \mathbb{R}$, then either*

- (1) ρ_1 and ρ_2 have the same marked complex length spectrum on $W(\alpha)$, or
- (2) ρ_1 and ρ_2 have conjugate marked complex length spectrum on $W(\alpha)$.

Proof. Lemma 2.3 implies that either $\lambda_{\rho_1}^2(\alpha) = \lambda_{\rho_2}^2(\alpha)$ or $\lambda_{\rho_1}^2(\alpha) = \overline{\lambda_{\rho_2}^2(\alpha)}$. If $\lambda_{\rho_1}^2(\alpha) = \overline{\lambda_{\rho_2}^2(\alpha)}$, then we consider the representation $\bar{\rho}_2$. In this case, $\lambda_{\rho_2}^2(\gamma) = \overline{\lambda_{\bar{\rho}_2}^2(\gamma)}$ for all $\gamma \in \pi_1(S)$. In particular, $\lambda_{\rho_1}^2(\alpha) = \lambda_{\bar{\rho}_2}^2(\alpha)$. Therefore, it suffices to prove that ρ_1 and ρ_2 have the same marked complex length spectrum on $W(\alpha)$ whenever $\lambda^2 = \lambda_{\rho_1}^2(\alpha) = \lambda_{\rho_2}^2(\alpha)$ and ρ_1 and ρ_2 have the same simple marked length spectrum.

First, suppose that β is a simple non-separating curve on S which intersects α once. We adopt the normalization and notation of Lemmas 2.3 and 2.5, so

$$t_i(n) = \mathrm{Tr}^2(\rho_i(\alpha^n \beta)) = \lambda^{2n} a_i^2 + 2a_i d_i + \lambda^{-2n} d_i^2.$$

Lemma 2.3 implies that for any n , either (i) $t_1(n) = t_2(n)$; (ii) $t_1(n) = \overline{t_2(n)}$; or (iii) $t_1(n) = 4 - t_2(n)$ and $t_1(n)$ is real.

If there is an infinite set of values of n such that $t_1(n) = t_2(n)$, then, by taking limits, we see that

$$a_1^2 = a_2^2, \quad a_1 d_1 = a_2 d_2 \quad \text{and} \quad d_1^2 = d_2^2.$$

It follows then that $t_1(n) = t_2(n)$ for all n . Moreover, since a_i and d_i are non-zero, either $a_1 = a_2$ and $d_1 = d_2$ or $a_1 = -a_2$ and $d_1 = -d_2$, so $\mathrm{Tr}^2(\rho_1(\beta)) = \mathrm{Tr}^2(\rho_2(\beta))$ which implies that $\lambda_{\rho_1}^2(\beta) = \lambda_{\rho_2}^2(\beta)$.

If there is an infinite set of values n such that $t_1(n) = 4 - t_2(n)$ with $t_i(n)$ real, then taking limits we have

$$a_1^2 = -a_2^2, \quad 2a_1 d_1 = 4 - 2a_2 d_2 \quad \text{and} \quad d_1^2 = -d_2^2.$$

It follows that $t_1(n) = 4 - t_2(n)$ for all n , so by Lemma 2.3, $t_1(n)$ is real for all n . Since $\Im(t_i(n)) = 0$ for all n ,

$$\lim_{n \rightarrow \infty} \frac{\Im(t_i(n))}{|\lambda|^{2n}} = \lim_{n \rightarrow \infty} \Im\left(\left(\frac{\lambda^2}{|\lambda|^2}\right)^n a_i^2\right) = 0.$$

Since $a_i^2 \neq 0$, by Lemma 2.2, this can only happen if $\frac{\lambda^2}{|\lambda|^2} = \pm 1$. Thus, $\lambda^2 = \lambda_{\rho_1}(\alpha^2)$ is real, contradicting our assumption.

Finally, if $t_1(n) = \overline{t_2(n)}$ for all but finitely many values of n , we may divide the resulting equation by $\overline{\lambda^{2n}}$ and take a limit, to see that

$$\lim_{n \rightarrow \infty} \left(\frac{\lambda^2}{\overline{\lambda^2}} \right)^n a_1^2 = \overline{a_2^2}.$$

Since $\lambda^2 \neq \overline{\lambda^2}$, the limit does not exist unless $a_1 = 0$, which again contradicts Lemma 2.2.

Therefore, if $\beta \in W(\alpha)$ intersects α once, then $\lambda_{\rho_1}^2(\beta) = \lambda_{\rho_2}^2(\beta)$.

Now suppose that β is a simple non-separating curve on S which does not intersect α . We choose β' to be a simple non-separating curve intersecting both α and β once.

For all n , $\alpha^n \beta' \in W(\alpha)$ and intersects α once. By the first part of the argument,

$$\lambda_{\rho_1}^2(\alpha^n \beta') = \lambda_{\rho_2}^2(\alpha^n \beta')$$

for all n . If there exists n_0 so that $\lambda_{\rho_1}^2(\alpha^{n_0} \beta')$ is not real, then since $\beta \in W(\alpha^{n_0} \beta')$ and intersects $\alpha^{n_0} \beta'$ exactly once, we may apply the above argument to show that $\lambda_{\rho_1}^2(\beta) = \lambda_{\rho_2}^2(\beta)$.

It remains to consider the case that $\lambda_{\rho_1}^2(\alpha^n \beta')$ is real for all n . Suppose that

$$\rho_i(\beta') = \pm \begin{pmatrix} a'_i & b'_i \\ c'_i & d'_i \end{pmatrix}.$$

Again, by Lemma 2.2 all the matrix coefficients must be non-zero. Since $\Im(\lambda_{\rho_1}^2(\alpha^n \beta')) = 0$ for all n , $\Im(\text{Tr}^2(\rho_1(\alpha^n \beta'))) = 0$ for all n , so, after dividing the resulting equation by $|\lambda|^{2n}$, for all n , and passing to the limit we see that

$$\lim_{n \rightarrow \infty} \Im \left(\left(\frac{\lambda^2}{|\lambda|^2} \right)^n a_1'^2 \right) = 0.$$

Since $\frac{\lambda^2}{|\lambda|^2} \notin \mathbb{R}$, this implies that $a'_1 = 0$, which is again a contradiction. Therefore, if $\beta \in W(\alpha)$ does not intersect α , then $\lambda_{\rho_1}^2(\beta) = \lambda_{\rho_2}^2(\beta)$ which completes the proof. \square

2.5. Assembly. We can now easily assemble the proof of Proposition 2.1.

Proof of Proposition 2.1. If there exists a simple, non-separating curve α on S so that $\lambda_{\rho_1}^2(\alpha)$ is not real, then Proposition 2.1 follows immediately from Lemma 2.6. If $\lambda_{\rho_1}^2(\alpha)$ is real for every simple, non-separating curve α on S , then Lemma 2.5 implies that ρ_1 and ρ_2 have the same marked complex length spectrum on $W(\alpha)$ for any non-separating simple closed curve α . A result of Sullivan [30] implies that there are only finitely many simple curves γ on S so that $\rho_1(\gamma)$ is parabolic, so we may always choose α so that $\rho_1(\alpha)$ is hyperbolic. \square

Remark. An examination of the proofs reveals that Proposition 2.1 holds whenever the length spectra of ρ_1 and ρ_2 agree on all simple, non-separating curves.

3. Simple marked length spectrum rigidity

We are now ready for the proof of our main result.

Theorem 1.1 (Simple length rigidity for Kleinian surface groups). *If S is a closed, connected, orientable surface of genus at least two, and $\rho_1 : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(S) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ are Kleinian surface groups with the same simple marked length spectrum, then ρ_1 is conjugate to either ρ_2 or $\bar{\rho}_2$.*

We begin with a brief sketch of the proof. It follows from Lemma 2.1 that, perhaps after replacing ρ_2 with a complex conjugate representation, there exists a simple, non-separating curve α so that $\rho_1(\alpha)$ is hyperbolic and ρ_1 and ρ_2 have the same marked complex length spectrum on $W(\alpha)$. We then lift ρ_1 and ρ_2 to representations into $\mathrm{SL}(2, \mathbb{C})$ which have the same trace on a standard set of generators $\{\alpha_1, \beta_1, \dots, \alpha_g, \beta_g\}$ where $\alpha = \alpha_1$, see Lemma 3.1. An analysis of the asymptotic behavior of the traces of $\alpha_j^n \beta$ allows us to conclude that the restriction of the lifts to any subgroup of the form $G_j = \langle \alpha_j, \beta_j \rangle$ are conjugate, see Lemma 3.2. A more intricate analysis of the same type is then applied to show that if we conjugate the lifts to agree on G_j , then, for any k , they either agree on G_k or differ by conjugation by a lift of the rotation of order two in the axis of the image of the commutator of $[\alpha_j, \beta_j]$, see Lemmas 3.3 and 3.4. The proof is then easily completed when the genus is greater than two, see Lemma 3.5, but a separate analysis is required when the genus is two, see Lemma 3.6.

Proof of Theorem 1.1. Proposition 2.1 implies that there exists a simple non-separating curve α such that $\rho_1(\alpha)$ is hyperbolic and ρ_1 and ρ_2 have the same marked complex length spectrum on $W(\alpha)$. If ρ_1 and ρ_2 have conjugate complex marked length spectrum on $W(\alpha)$, then ρ_1 and $\bar{\rho}_2$ have the same marked complex length spectrum on $W(\alpha)$. Therefore, we may assume that ρ_1 and ρ_2 have the same marked complex length spectrum on $W(\alpha)$.

We begin by choosing lifts whose traces agree on a standard set of generators which includes α . We will call $\mathcal{S} = \{\alpha_1, \beta_1, \dots, \alpha_g, \beta_g\}$ a *standard set of generators* for $\pi_1(S)$ if each α_j and β_j is non-separating,

$$\pi_1(S) = \langle \alpha_j, \beta_j \mid \prod_{j=1}^g [\alpha_i, \beta_i] = \mathrm{id} \rangle \quad \text{and} \quad i(\alpha_j, \beta_j) = 1$$

for all j and if $j \neq k$ then

$$i(\alpha_j, \alpha_k) = i(\beta_j, \beta_k) = i(\alpha_j, \beta_k) = 0,$$

see Figure 1. We say that two lifts $\tilde{\rho}_1$ and $\tilde{\rho}_2$ of ρ_1 and ρ_2 are *trace normalized* with respect to \mathcal{S} if

- (1) $\rho_1(\delta)$ is hyperbolic for all $\delta \in \mathcal{S}$, and
- (2) $\text{Tr}(\tilde{\rho}_1(\delta)) = \text{Tr}(\tilde{\rho}_2(\delta))$ for all $\delta \in \mathcal{S}$.




Figure 1. Generators of $\pi_1(S, p)$.

Lemma 3.1. Suppose that $\rho_1 : \pi_1(S) \rightarrow \text{PSL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(S) \rightarrow \text{PSL}(2, \mathbb{C})$ are Kleinian surface groups with the same marked complex length spectrum on $W(\alpha)$ for some simple, non-separating curve α . If $\rho_1(\alpha)$ is hyperbolic, then there exists a standard set of generators \mathcal{S} , so that $\alpha_1 = \alpha$, and lifts $\tilde{\rho}_1 : \pi_1(S) \rightarrow \text{SL}(2, \mathbb{C})$ and $\tilde{\rho}_2 : \pi_1(S) \rightarrow \text{SL}(2, \mathbb{C})$ of ρ_1 and ρ_2 which are trace normalized with respect to \mathcal{S} .

Proof. Let $\alpha_1 = \alpha$. Choose a simple non-separating curve β which intersects α_1 exactly once so that $\rho_1(\beta_1)$ is hyperbolic. (We may do so, since, by a result of Sullivan [30], there are only finitely many simple curves γ such that $\rho_1(\gamma)$ is parabolic and there are infinitely many possibilities for β_1 .) Extend $\{\alpha_1, \beta_1\}$ to a standard set of generators $\{\alpha_1, \beta_1, \dots, \alpha_g, \beta_g\}$. We may assume that $\rho_1(\alpha_j)$ is hyperbolic for all $j \geq 2$, by replacing α_j by $\alpha_j \beta_j^n$ for some n if necessary. We may then assume that $\rho_1(\beta_j)$ is hyperbolic for all $j \geq 2$ by replacing β_j by $\beta_j \alpha_j^n$ for some n if necessary. Notice that $\mathcal{S} \subseteq W(\alpha)$.

Since each ρ_i is discrete and faithful, each ρ_i lifts to a representation $\rho'_i : \pi_1(S) \rightarrow \text{SL}(2, \mathbb{C})$ (see Culler [12] or Kra [20]). Let

$$\tilde{\rho}_2(\delta) = \begin{cases} \rho'_2(\delta) & \text{if } \text{Tr}(\rho'_1(\delta)) = \text{Tr}(\rho'_2(\delta)), \\ -\rho'_2(\delta) & \text{if } \text{Tr}(\rho'_1(\delta)) = -\text{Tr}(\rho'_2(\delta)), \end{cases}$$

for all $\delta \in \mathcal{S}$. Notice that $\tilde{\rho}_i(\delta)$ is a lift of $\rho_i(\delta)$ for all $\delta \in \mathcal{S}$ and that $\tilde{\rho}_i(\prod_{j=1}^g [\alpha_j, \beta_j]) = I$, since $\rho'_i(\prod_{j=1}^g [\alpha_j, \beta_j]) = I$. Therefore, $\tilde{\rho}_1$ and $\tilde{\rho}_2$ are lifts of ρ_1 and ρ_2 which are trace normalized with respect to \mathcal{S} . \square

We next show that the trace normalized lifts are conjugate on the subgroups $G_j = \langle \alpha_j, \beta_j \rangle$.

Lemma 3.2. *Suppose that $\tilde{\rho}_1 : \pi_1(S) \rightarrow \mathrm{SL}(2, \mathbb{C})$ and $\tilde{\rho}_2 : \pi_1(S) \rightarrow \mathrm{SL}(2, \mathbb{C})$ are trace normalized lifts, with respect to a standard generating set \mathcal{S} , of Kleinian surface groups with the same marked complex length spectrum on $W(\alpha_1)$. If $j \in \{1, \dots, g\}$, then there exists $K_j \in \mathrm{SL}(2, \mathbb{C})$ such that $\tilde{\rho}_2|_{G_j} = (K_j \tilde{\rho}_1 K_j^{-1})|_{G_j}$. In particular, if $\gamma \in G_j$, then $\mathrm{Tr}(\tilde{\rho}_1(\gamma)) = \mathrm{Tr}(\tilde{\rho}_2(\gamma))$.*

Proof. Fix j for the remainder of the proof of the lemma and assume that $\tilde{\rho}_1$ and $\tilde{\rho}_2$ are (α_j, β_j) -normalized, so

$$\tilde{\rho}_1(\alpha_j) = \tilde{\rho}_2(\alpha_j) = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix},$$

where $|\lambda| > 1$, and

$$\tilde{\rho}_i(\beta_j) = \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix},$$

where $a_i d_i - b_i c_i = 1$ and all the co-efficients are non-zero, by Lemma 2.2. Since $\mathrm{Tr}(\tilde{\rho}_1(\beta_j)) = \mathrm{Tr}(\tilde{\rho}_2(\beta_j))$,

$$a_1 + d_1 = a_2 + d_2.$$

The curve $\alpha_j^n \beta_j \in W(\alpha_1)$, for all n , since it is non-separating, simple and disjoint from α if $j > 1$ and intersects α exactly once if $j = 1$. Therefore, $\mathrm{Tr}(\tilde{\rho}_1(\alpha_j^n \beta_j)) = \pm \mathrm{Tr}(\tilde{\rho}_2(\alpha_j^n \beta_j))$, which gives the equation

$$\lambda^n a_1 + \lambda^{-n} d_1 = \pm (\lambda^n a_2 + \lambda^{-n} d_2).$$

If $\lambda^n a_1 + \lambda^{-n} d_1 = -(\lambda^n a_2 + \lambda^{-n} d_2)$ for infinitely many values of n , then we see, by dividing by λ^n and passing to a limit, that $a_1 = -a_2$. It then follows that $d_1 = -d_2$. Thus, $a_1 + d_1 = -(a_2 + d_2)$ which contradicts the fact that the traces of $\tilde{\rho}_i$ agree (and are non-zero) on elements of \mathcal{S} .

Therefore, there are an infinitely many values of n where $\lambda^n a_1 + \lambda^{-n} d_1 = \lambda^n a_2 + \lambda^{-n} d_2$. Taking limits again, we see that $a_1 = a_2$ and $d_1 = d_2$. Therefore,

$$b_1 c_1 = 1 - a_1 d_1 = 1 - a_2 d_2 = b_2 c_2.$$

Choose $u \in \mathbb{C}$ such that $u^2 = b_2/b_1 = c_1/c_2$. Then $u^2 b_1 = b_2$ and $u^{-2} c_1 = c_2$. Let

$$K_j = \begin{pmatrix} u & 0 \\ 0 & u^{-1} \end{pmatrix}.$$

Notice that $K_j \tilde{\rho}_1(\alpha_j) K_j^{-1} = \tilde{\rho}_1(\alpha_j)$ and that

$$\begin{aligned} K_j \tilde{\rho}_1(\beta_j) K_j^{-1} &= \begin{pmatrix} u & 0 \\ 0 & u^{-1} \end{pmatrix} \begin{pmatrix} a_1 & b_1 \\ c_1 & d_1 \end{pmatrix} \begin{pmatrix} u^{-1} & 0 \\ 0 & u \end{pmatrix} \\ &= \begin{pmatrix} a_1 & u^2 b_1 \\ u^{-2} c_1 & d_1 \end{pmatrix} = \tilde{\rho}_2(\beta_j). \end{aligned}$$

Therefore, K_j conjugates the restriction of $\tilde{\rho}_1$ to G_j to the restriction of $\tilde{\rho}_2$ to G_j . \square

We begin our analysis of trace normalized lifts which agree on G_j by examining the relationship between the images of elements of \mathcal{S} . Lemma 3.3 is the crucial step in the proof of our main result. The additional information concerning matrix co-efficients in case (2) will only be used when S has genus two.

Lemma 3.3. *Suppose that $\rho_1^j : \pi_1(S) \rightarrow \mathrm{SL}(2, \mathbb{C})$ and $\rho_2^j : \pi_1(S) \rightarrow \mathrm{SL}(2, \mathbb{C})$ are trace normalized lifts, with respect to a standard generating set \mathcal{S} , of Kleinian surface groups with the same complex marked length spectrum on $W(\alpha_1)$. If ρ_1^j and ρ_2^j agree on $G_j = \langle \alpha_j, \beta_j \rangle$ and $\delta \in \mathcal{S}$, then either*

- (1) $\rho_1^j(\delta) = \rho_2^j(\delta)$, or
- (2) $\rho_1^j([\alpha_j, \beta_j])$ is hyperbolic and $R_j \rho_1^j(\delta) R_j^{-1} = \rho_2^j(\delta)$, where R_j is a lift of the rotation of order two about the axis of $\rho_1^j([\alpha_j, \beta_j])$. Moreover, if $\gamma \in G_j$ such that $\gamma, \gamma\delta \in W(\alpha_1)$, and

$$\rho_1^j([\alpha_j, \beta_j]) = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}, \quad \rho_1^j(\gamma) = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad \text{and} \quad \rho_1^j(\delta) = \begin{pmatrix} e & f \\ g & h \end{pmatrix},$$

where $|\lambda| > 1$, then

$$\frac{e}{h} = -\frac{d}{a}.$$

Proof. The proof breaks up into two cases, depending on whether $\rho_1^j([\alpha_j, \beta_j])$ is hyperbolic or parabolic.

Case I. $\rho_1^j([\alpha_j, \beta_j])$ is hyperbolic. We may assume, by conjugating, that

$$\rho_1^j([\alpha_j, \beta_j]) = \rho_2^j([\alpha_j, \beta_j]) = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix},$$

where $|\lambda| > 1$. As ρ_1^j and ρ_2^j agree on G_j , we can assume that δ is not either α_j or β_j . Let

$$\rho_i^j(\delta) = \begin{pmatrix} e_i & f_i \\ g_i & h_i \end{pmatrix}$$

where $e_i h_i - f_i g_i = 1$. Moreover, since the traces agree on generators,

$$e_1 + h_1 = e_2 + h_2.$$

Suppose that $\gamma \in G_j$ and $\gamma, \gamma\delta \in W(\alpha_1)$. (We can choose either $\gamma = \beta_j$ or $\gamma = \beta_j^{-1}$ depending on the orientations on the curves, see Figure 2). Let

$$\rho_1^j(\gamma) = \rho_2^j(\gamma) = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

where $ad - bc = 1$.

Let $\gamma_n = [\alpha_j, \beta_j]^n \gamma [\alpha_j, \beta_j]^{-n} \delta$, which also lies in $W(\alpha_1)$, since it is the image of $\gamma\delta$ under the n -fold Dehn twist about $[\alpha_j, \beta_j]$. So, by assumption,

$$\text{Tr}(\rho_1^j(\gamma_n)) = \pm \text{Tr}(\rho_2^j(\gamma_n)).$$

By expanding, we see that, for all n ,

$$ae_1 + dh_1 + bg_1\lambda^{2n} + cf_1\lambda^{-2n} = \pm(ae_2 + dh_2 + bg_2\lambda^{2n} + cf_2\lambda^{-2n}).$$

Thus there exists an increasing subsequence where the traces either all agree or all differ up to sign.




Figure 2. Curves γ, δ and $[\alpha_j, \beta_j]$.

If there exists an increasing sequence $\{n_j\}$ such that the traces all agree, then

$$ae_1 + dh_1 + bg_1\lambda^{2n_j} + cf_1\lambda^{-2n_j} = ae_2 + dh_2 + bg_2\lambda^{2n_j} + cf_2\lambda^{-2n_j},$$

for all n_j . Dividing the above equation by λ^{2n_j} and taking a limit as $j \rightarrow \infty$, we see that $bg_1 = bg_2$. Since $b \neq 0$, by Lemma 2.2, $g_1 = g_2$. Thus,

$$ae_1 + dh_1 + cf_1\lambda^{-2n_j} = ae_2 + dh_2 + cf_2\lambda^{-2n_j}$$

and we see, by taking the limit as $n_j \rightarrow \infty$, that $ae_1 + dh_1 = ae_2 + dh_2$. Finally, since $cf_1\lambda^{-2n_j} = cf_2\lambda^{-2n_j}$ and λ and c are non-zero, $f_1 = f_2$. Summarizing, we have

$$ae_1 + dh_1 = ae_2 + dh_2, \quad f_1 = f_2, \quad \text{and} \quad g_1 = g_2. \quad (3.1)$$

Similarly, if there exists an increasing sequence $\{n_j\}$ such that the traces all disagree then,

$$ae_1 + dh_1 + bg_1\lambda^{2n_j} + cf_1\lambda^{-2n_j} = -(ae_2 + dh_2 + bg_2\lambda^{2n_j} + cf_2\lambda^{-2n_j})$$

for all n_j . Taking limits as above, we conclude that

$$ae_1 + dh_1 = -(ae_2 + dh_2), \quad f_1 = -f_2, \quad \text{and} \quad g_1 = -g_2. \quad (3.2)$$

Thus given any $\gamma \in G_j$ such that $\gamma, \gamma\delta \in W(\alpha_1)$, then γ either satisfies equation (3.1) or (3.2). Since f_i and g_i are non-zero, we conclude that, with the above normalization, either

- (a) Equation (3.1) holds for all $\gamma \in G_j$ such that $\gamma, \gamma\delta \in W(\alpha_1)$, or
- (b) Equation (3.2) holds for all $\gamma \in G_j$ such that $\gamma, \gamma\delta \in W(\alpha_1)$.

Case Ia. Equation (3.1) holds for all $\gamma \in G_j$ such that $\gamma, \gamma\delta \in W(\alpha_1)$. Choose $\gamma \in G_j$ such that $\gamma, \gamma\delta \in W(\alpha_1)$ and let

$$\rho_1^j(\gamma) = \rho_2^j(\gamma) = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

where $ad - bc = 1$. Then, by equation (3.1),

$$ae_1 + dh_1 = ae_2 + dh_2, \quad f_1 = f_2, \quad \text{and} \quad g_1 = g_2.$$

Since $e_i h_i - f_i g_i = 1$, we conclude that $e_1 h_1 = e_2 h_2$, so, since we also have $e_1 + h_1 = e_2 + h_2$,

$$(x - e_1)(x - h_1) = x^2 - 2(e_1 + h_1)x + e_1 h_1 = (x - e_2)(x - h_2),$$

which implies that either (i) $e_1 = e_2$ and $h_1 = h_2$; or (ii) $e_1 = h_2$ and $h_1 = e_2$.

If $e_1 = e_2$ and $h_1 = h_2$, then, since we already know that $f_1 = f_2$ and $g_1 = g_2$, we may conclude that $\rho_1^j(\delta) = \rho_2^j(\delta)$ and we are in case (1).

If $e_1 = h_2$ and $h_1 = e_2$, then, since $ae_1 + dh_1 = ae_2 + dh_2$, we conclude that

$$(a - d)(e_1 - h_1) = 0.$$

If $a \neq d$, then this implies that $e_1 = h_1$, so $e_1 = e_2$. Since we already know that $f_1 = f_2$ and $g_1 = g_2$, and all the matrix co-efficients are non-zero, we may conclude that $\rho_1^j(\delta) = \rho_2^j(\delta)$, so we are again in case (1).

In order to conclude that we are in case (1), it only remains to check that we can choose $\gamma_0 \in G_j$, so that $\gamma_0, \gamma_0\delta \in W(\alpha_1)$ and

$$\rho_1^j(\gamma_0) = \rho_2^j(\gamma_0) = \begin{pmatrix} a_0 & b_0 \\ c_0 & d_0 \end{pmatrix},$$

where $a_0 \neq d_0$.

We may assume that we have initially chosen γ to be either β_j^{-1} or β_j , and that, with the above notation, $a = d$. We may then choose γ_0 to be the element in $\{[\alpha_j, \beta_j]\gamma, [\alpha_j, \beta_j]^{-1}\gamma\}$ which is simple and has the property that $\gamma_0, \gamma_0\delta \in W(\alpha_1)$ (see Figure 3). Observe that either $a_0 = \lambda a$ and $d_0 = \lambda^{-1}d$ or $a_0 = \lambda^{-1}a$ and $d_0 = \lambda d$, so $a_0 \neq d_0$. This choice of γ_0 allows us to complete the proof.




Figure 3. The curve γ_0 .

Case Ib. Equation (3.2) holds for all $\gamma \in G_j$ such that $\gamma, \gamma\delta \in W(\alpha_1)$. Choose $\gamma \in G_j$ so that $\gamma, \gamma\delta \in W(\alpha_1)$ and let

$$\rho_1^j(\gamma) = \rho_2^j(\gamma) = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$

where $ad - bc = 1$. Then, by equation (3.2).

$$ae_1 + dh_1 = -(ae_2 + dh_2), \quad f_1 = -f_2, \quad \text{and} \quad g_1 = -g_2.$$

As $e_i h_i - f_i g_i = 1$, we conclude that $e_1 h_1 = e_2 h_2$. Since $e_1 h_1 = e_2 h_2$ and $e_1 + h_1 = e_2 + h_2$, we may conclude, just as in Case Ia, that either (i) $e_1 = h_2$ and $h_1 = e_2$; or (ii) $e_1 = e_2$, and $h_1 = h_2$.

If $e_1 = h_2$ and $h_1 = e_2$, then, since $ae_1 + dh_1 = -(ae_2 + dh_2)$, we see that

$$\text{Tr}(\rho_1^j(\gamma))\text{Tr}(\rho_1^j(\delta)) = (a + d)(e_1 + h_1) = ae_1 + dh_1 + ae_2 + dh_2 = 0,$$

which is impossible since both traces are non-zero. Therefore, $e_1 = e_2$ and $h_1 = h_2$. Since $f_1 = -f_2$ and $g_1 = -g_2$,

$$\rho_2^j(\delta) = \begin{pmatrix} e_1 & -f_1 \\ -g_1 & h_1 \end{pmatrix} = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \begin{pmatrix} e_1 & f_1 \\ g_1 & h_1 \end{pmatrix} \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix} = R_j \rho_1^j(\delta) R_j^{-1},$$

so $\rho_1^j(\delta) = R_j \rho_2^j(\delta) R_j^{-1}$ and R_j is a lift of the rotation of order two about the axis of $\rho_1^j([\alpha_j, \beta_j])$.

Moreover, since $ae_1 + dh_1 = -(ae_2 + dh_2) = -(ae_1 + dh_1)$, we see that $ae_1 + dh_1 = 0$, so

$$\frac{e_1}{h_1} = -\frac{d}{a},$$

and we are in case (2).

Case II. $\rho_1^j([\alpha_j, \beta_j])$ is parabolic. Choose $\gamma \in G_j$ so that $\gamma, \gamma\delta \in W(\alpha_1)$. We may conjugate so that

$$\rho_1^j([\alpha_j, \beta_j]) = \rho_2^j([\alpha_j, \beta_j]) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

and 0 is a fixed points of $\rho_1^j(\gamma) = \rho_2^j(\gamma)$, so

$$\rho_1^j(\gamma) = \rho_2^j(\gamma) = \begin{pmatrix} a & 0 \\ c & d \end{pmatrix},$$

where $ad = 1$. So, a and d are non-zero and c is non-zero, since otherwise $\rho_1^i([\alpha_i, \beta_i])$ and $\rho_1^i(\gamma)$ would have a common fixed point.

Let

$$\rho_i^j(\delta) = \begin{pmatrix} e_i & f_i \\ g_i & h_i \end{pmatrix},$$

where $e_i h_i - f_i g_i = 1$ and all the matrix coefficients are non-zero. Moreover, since the lifts are trace normalized with respect to \mathcal{S} ,

$$e_1 + h_1 = e_2 + h_2.$$

Again, let $\gamma_n = [\alpha_i, \beta_i]^n \gamma [\alpha_i, \beta_i]^{-n} \delta$, which lies in $W(\alpha_1)$, so

$$\text{Tr}(\rho_1^i(\gamma_n)) = \pm \text{Tr}(\rho_2^i(\gamma_n)).$$

Expanding, we obtain

$$\begin{aligned} (a + nc)e_1 + (n(d - a) - n^2c)g_1 + cf_i + (d - nc)h_1 \\ = \pm((a + nc)e_2 + (n(d - a) - n^2c)g_2 + cf_2 + (d - nc)h_2). \end{aligned}$$

If there are infinitely many n where the traces differ, then, by dividing by n^2 and taking limits, we conclude that $cg_1 = -cg_2$, so, since $c \neq 0$, $g_1 = -g_2$. By successively taking limits, we further conclude that

$$c(e_1 - h_1) = -c(e_2 - h_2) \quad \text{and} \quad ae_1 + cf_1 + dh_1 = -ae_2 - cf_2 - dh_2.$$

Since $c \neq 0$, we see that $e_1 - h_1 = -e_2 + h_2$. As $e_1 + h_1 = e_2 + h_2$, this implies that $e_1 = h_2$ and $h_1 = e_2$. Therefore,

$$f_1g_1 = e_1h_1 - 1 = e_2h_2 - 1 = f_2g_2 = -f_2g_1.$$

As $\rho_i^j(\delta)$ does not have a fixed point at ∞ , g_1 must be non-zero, so $f_1 = -f_2$. Since $ae_1 + cf_1 + dh_1 = -ae_2 - cf_2 - dh_2$, this implies that

$$\text{Tr}(\gamma)\text{Tr}(\delta) = (a + d)(e_1 + h_1) = 0$$

which is a contradiction, since both traces are non-zero.

Therefore, there are infinitely many n where the traces agree, so, taking limits as above, we see that

$$g_1 = g_2, \quad c(e_1 - h_1) = c(e_2 - h_2), \quad \text{and} \quad ae_1 + cf_1 + dh_1 = ae_2 + cf_2 + dh_2.$$

Since $c \neq 0$, $e_1 - h_1 = e_2 - h_2$, which we may combine with the fact that $e_1 + h_1 = e_2 + h_2$, to conclude that $e_1 = e_2$ and $h_1 = h_2$. Therefore,

$$f_1g_1 = e_1h_1 - 1 = e_2h_2 - 1 = f_2g_2 = f_2g_1$$

which implies that $f_1 = f_2$, so $\rho_1^j(\delta) = \rho_2^j(\delta)$, which implies that we are in case (1). \square

We now refine our analysis of trace normalized lifts which agree on G_j to show that, for all k , they either agree on G_k or differ by a rotation in the axis of the image of the commutator in G_j .

Lemma 3.4. *Suppose that $\rho_1 : \pi_1(S) \rightarrow \text{SL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(S) \rightarrow \text{SL}(2, \mathbb{C})$ are trace normalized lifts, with respect to a standard generating set \mathcal{S} , of Kleinian surface groups with the same marked complex length spectrum on $W(\alpha_1)$. If $j, k \in \{1, \dots, g\}$ and ρ_1^j and ρ_2^j agree on $G_j = \langle \alpha_j, \beta_j \rangle$, then either*

- (1) ρ_1^j and ρ_2^j agree on G_k , or
- (2) $\rho_1^j([\alpha_j, \beta_j])$ is hyperbolic and $\rho_1^j|_{G_k} = (R_j \rho_2^j R_j^{-1})|_{G_k}$ where R_j is a lift of the rotation of order two about the axis of $\rho_1^j([\alpha_j, \beta_j])$.

Proof. If neither (1) or (2) holds, then Lemma 3.3 implies that, perhaps after switching α_k and β_k , $\rho_1^j([\alpha_j, \beta_j])$ is hyperbolic,

$$\rho_2^j(\alpha_k) = \rho_1^j(\alpha_k) \quad \text{and} \quad \rho_2^j(\beta_k) = R_j \rho_1(\beta_k) R_j^{-1}$$

where R_j is a lift of the rotation of order two about the axis of $\rho_1^j([\alpha_j, \beta_j])$.

Lemma 3.2 implies that there exists $K \in \mathrm{SL}(2, \mathbb{C})$ so that ρ_1^j and $K\rho_2^j K^{-1}$ agree on G_k . If $K = \pm I$, then ρ_1^j and $\rho_2^j = K\rho_2^j K^{-1}$ agree on G_k and we are in case (1). So, we may assume that $K \neq \pm I$ and

$$\rho_1^j(\alpha_k) = K\rho_2^j(\alpha_k)K^{-1} = K\rho_1^j(\alpha_k)K^{-1}$$

and
$$\rho_1^j(\beta_k) = K\rho_2^j(\beta_k)K^{-1} = KR_j\rho_1^j(\beta_k)R_j^{-1}K^{-1}.$$

It follows that K fixes the fixed points of $\rho_1^j(\alpha_k)$ and that KR_j fixes the fixed points of $\rho_1^j(\beta_k)$.

If ρ_1^j and $K\rho_2^j K^{-1}$ agree on either α_j or β_j , then, since ρ_1^j and ρ_2^j agree on G_j , either

$$\rho_1^j(\alpha_j) = K\rho_2^j(\alpha_j)K^{-1} = K\rho_1^j(\alpha_j)K^{-1}$$

or
$$\rho_1^j(\beta_j) = K\rho_2^j(\beta_j)K^{-1} = K\rho_1^j(\beta_j)K^{-1}$$

which implies that K fixes the fixed points of either $\rho_1^j(\alpha_j)$ or $\rho_1^j(\beta_j)$. But, since K fixes the fixed points of $\rho_1^j(\alpha_k)$, and the fixed points of $\rho_1^j(\alpha_k)$, $\rho_1^j(\alpha_j)$ and $\rho_1^j(\beta_j)$ are all distinct, this implies that $K = \pm I$, which is a contradiction.

Therefore, ρ_1^j and $K\rho_2^j K^{-1}$ disagree on both α_j and β_j . Lemma 3.3 then implies that $\rho_1^j([\alpha_k, \beta_k])$ is hyperbolic and

$$\rho_1^j(\alpha_j) = R_k K \rho_2^j(\alpha_j) K^{-1} R_k^{-1} \quad \text{and} \quad \rho_1^j(\beta_j) = R_k K \rho_2^j(\beta_j) K^{-1} R_k^{-1}$$

where R_k is a lift of the rotation of order two about the axis of $\rho_1^j([\alpha_k, \beta_k])$. Therefore, ρ_1^j and $R_k K \rho_2^j K^{-1} R_k^{-1}$ agree on G_j . Since ρ_1^j and ρ_2^j agree on G_j this implies that $R_k K = \pm I$, so $K = \pm R_k^{-1}$. Therefore, K fixes the fixed points of $\rho_1^j([\alpha_k, \beta_k])$. However, since we already know that K fixes the fixed points of $\rho_1^j(\alpha_k)$, this implies that $K = \pm I$, which is again a contradiction. \square

It is now relatively simple to use Lemma 3.4 to complete the proof when S has genus at least three.

Lemma 3.5. *Suppose that S has genus $g \geq 3$ and that $\tilde{\rho}_1 : \pi_1(S) \rightarrow \mathrm{SL}(2, \mathbb{C})$ and $\tilde{\rho}_2 : \pi_1(S) \rightarrow \mathrm{SL}(2, \mathbb{C})$ are trace normalized lifts, with respect to a standard generating set \mathcal{S} , of Kleinian surface groups with the same marked complex length spectrum on $W(\alpha_1)$. Then, $\tilde{\rho}_1$ is conjugate to $\tilde{\rho}_2$ in $\mathrm{SL}(2, \mathbb{C})$.*

Proof. Lemma 3.2 implies that we may choose conjugates ρ_1^1 and ρ_2^1 of $\tilde{\rho}_1$ and $\tilde{\rho}_2$ which agree on $G_1 = \langle \alpha_j, \beta_j \rangle$. Lemma 3.4 implies that for all $j > 1$, the restrictions of ρ_1^1 and ρ_2^1 to G_j either agree or are conjugate by R_1 , where R_1 is a lift of the rotation of order two about the axis of $\rho_1^1([\alpha_1, \beta_1])$. If $\rho_1^1 \neq \rho_2^1$, we may assume without loss of generality that $\rho_1^1|_{G_2} = (R_1 \rho_2^1 R_1^{-1})|_{G_2}$.

Lemma 3.4 implies that either (1) ρ_1^1 and $R_1\rho_2^1R_1^{-1}$ agree on G_1 ; or (2) $\rho_2^1([\alpha_2, \beta_2])$ is hyperbolic and ρ_1^1 and $R_1\rho_2^1R_1^{-1}$ are conjugate by R_2 on G_1 where R_2 is a lift of the rotation of order two about the axis of $\rho_1^1([\alpha_2, \beta_2])$.

If ρ_1^1 and $R_1\rho_2^1R_1^{-1}$ agree on G_1 , then, since ρ_1^1 and ρ_2^1 agree on G_1 , R_1 commutes with every element of $\rho_1^1(G_1)$. Since $\rho_1^1(G_1)$ is non-elementary, this implies that $R_1 = \pm I$, which is a contradiction.

If ρ_1^1 and $R_2(R_1\rho_2^1R_1^{-1})R_2^{-1}$ agree on G_1 , then since ρ_1^1 and ρ_2^1 agree on G_1 , we similarly conclude that $R_2R_1 = \pm I$. So, $R_2 = \pm R_1^{-1}$. This would imply that $\rho_1^1([\alpha_1, \beta_1])$ and $\rho_1^1([\alpha_2, \beta_2])$ have the same axis, so share fixed points, which is a contradiction unless S has genus 2 and $[\alpha_1, \beta_1] = [\alpha_2, \beta_2]^{-1}$. \square

We now complete the proof by establishing our result in the genus two case.

Lemma 3.6. *Suppose that S has genus $g = 2$ and that $\tilde{\rho}_1 : \pi_1(S) \rightarrow \mathrm{SL}(2, \mathbb{C})$ and $\tilde{\rho}_2 : \pi_1(S) \rightarrow \mathrm{SL}(2, \mathbb{C})$ are trace normalized lifts, with respect to a standard generating set \mathcal{S} , of Kleinian surface groups with the same marked complex length spectrum on $W(\alpha_1)$. Then, $\tilde{\rho}_1$ is conjugate to $\tilde{\rho}_2$ in $\mathrm{SL}(2, \mathbb{C})$.*

Proof. In this case $\pi_1(S) = \langle \alpha_1, \beta_1, \alpha_2, \beta_2 \rangle$ given by the standard oriented curves as in Figure 4. Lemma 3.2 implies that we may conjugate $\tilde{\rho}_i$ to ρ_i^1 so that ρ_1^1 and ρ_2^1 agree on G_1 .

Suppose that $\rho_1^1 \neq \rho_2^1$. Lemma 3.4 implies that $\rho_1^1([\alpha_1, \beta_1])$ is hyperbolic and ρ_1^1 and $R_1\rho_2^1R_1^{-1}$ agree on G_2 where R_1 is a lift of the rotation of order two about the axis of $\rho_1^1([\alpha_1, \beta_1])$. We may normalize so that

$$\rho_1^1([\alpha_1, \beta_1]) = \rho_2^1([\alpha_1, \beta_1]) = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}$$

where $|\lambda| > 1$.




Figure 4. Surface of genus 2 with standard generators.

Let $\gamma = \alpha_1^{-1}$. Notice that, assuming that elements of \mathcal{S} are oriented as in Figure 4, $\gamma\alpha_2$ and $\gamma\beta_2$ are simple and lie in $W(\alpha_1)$. Define $A = \rho_1^1(\alpha_1^{-1})$, $C = \rho_1^1(\alpha_2)$, and $D = \rho_1^1(\beta_2)$, and adapt the convention that if a matrix is denoted by capital letter X , then it has coefficients (x_{ij}) . Since ρ_1^1 and ρ_2^1 differ on both α_2 and β_2 , Lemma 3.3 implies that

$$\frac{c_{11}}{c_{22}} = \frac{d_{11}}{d_{22}} = -\frac{a_{22}}{a_{11}}.$$

We now consider the element $\alpha_2\beta_2 \in G_2$. We replace generators $\{\alpha_2, \beta_2\}$ by $\{\alpha_2\beta_2, \alpha_2^{-1}\}$ in \mathcal{S} to form another standard generating set $\mathcal{S}' = \{\alpha_1, \beta_1, \alpha_2\beta_2, \alpha_2^{-1}\}$ (we note that $[\alpha_2\beta_2, \alpha_2^{-1}] = [\alpha_2, \beta_2]$). The representations ρ_1^1 and ρ_2^1 agree on G_1 and their traces agree on \mathcal{S}' . Notice that $\gamma(\alpha_2\beta_2) \in W(\alpha_1)$ (see Figure 4) and let $E = CD = \rho_1(\alpha_2\beta_2)$. Since ρ_1^1 and ρ_2^1 agree on G_1 and disagree on $\alpha_2\beta_2$, Lemma 3.3 implies that

$$\frac{e_{11}}{e_{22}} = -\frac{a_{22}}{a_{11}}.$$

If $M = CDC^{-1}D^{-1} = \rho_1^1([\alpha_2, \beta_2]^{-1}) = \rho_1^1([\alpha_1, \beta_1])$, then

$$M = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}.$$

Since $DC = M^{-1}CD = M^{-1}E$ and $C^{-1}(CD)C = DC$,

$$\text{Tr}(DC) = \text{Tr}(CD) = \text{Tr}(E) = \text{Tr}(M^{-1}E),$$

so $e_{11} + e_{22} = \lambda^{-1}e_{11} + \lambda e_{22}$. Therefore

$$\frac{c_{11}}{c_{22}} = \frac{d_{11}}{d_{22}} = \frac{e_{11}}{e_{22}} = \frac{(\lambda - 1)}{(1 - \lambda^{-1})} = \lambda.$$

Since $CD = MDC$,

$$(CD)_{12} = c_{11}d_{12} + c_{12}d_{22} = (MDC)_{12} = \lambda(DC)_{12} = \lambda(d_{11}c_{12} + d_{12}c_{22}),$$

so, since $c_{11} = \lambda c_{22}$ and $d_{11} = \lambda d_{22}$,

$$\lambda c_{22}d_{12} + c_{12}d_{22} = \lambda(\lambda d_{22}c_{12} + d_{12}c_{22}),$$

which implies that

$$(\lambda^2 - 1)c_{12}d_{22} = 0.$$

Thus, since d_{22} and c_{12} are both non-zero, $\lambda^2 = 1$ which is a contradiction. Therefore, it must be the case that $\rho_1^1 = \rho_2^1$. \square

Remark. Observe that in the proof we only assume that there exists a simple non-separating curve α such that ρ_1 and ρ_2 have the same or conjugate marked complex length spectrum on $W(\alpha)$. It then follows from the remark at the end of Section 2 that ρ_1 and ρ_2 are conjugate in the isometry group of \mathbb{H}^3 if they have the same marked length spectrum on all simple, non-separating curves.

4. Acylindrical 3-manifolds

We now use our main result to prove our length rigidity theorem for acylindrical 3-manifolds.

Theorem 1.2. *If M is a compact, acylindrical, hyperbolizable 3-manifold, and $\rho_1 : \pi_1(M) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(M) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ are two discrete faithful representations, such that $\ell_{\rho_1}(\alpha) = \ell_{\rho_2}(\alpha)$ if $\alpha \in \pi_1(M)$ is represented by a simple closed curve on ∂M , then ρ_1 is conjugate to either ρ_2 or $\bar{\rho}_2$.*

In order to motivate the more general proof, we offer a sketch of the proof in the case that M has connected boundary and ρ_1 and ρ_2 are both convex cocompact, i.e. each $N_i = \mathbb{H}^3 / \rho_i(\pi_1(M))$ admits a conformal compactification with boundary $\partial_c N_i$. After perhaps replacing ρ_2 with a complex conjugate representation, we may assume by Johansson's Classification Theorem [16], that there exists an orientation-preserving homeomorphism $j : N_1 \rightarrow N_2$ in the homotopy class determined by $\rho_2 \circ \rho_1^{-1}$. In this case, work of Bers [2] implies that ρ_1 is conjugate to ρ_2 if and only if j extends, up to isotopy, to a conformal homeomorphism $\hat{j} : \partial_c N_1 \rightarrow \partial_c N_2$.

If the length spectra of ρ_1 and ρ_2 agree on simple curves on the boundary, our main result implies that the restrictions of ρ_1 and ρ_2 to the fundamental group of the boundary agree up to conjugacy in the isometry group of \mathbb{H}^3 . It follows that the conformal compactifications N_1^S and N_2^S of the covers of N_1 and N_2 associated to $\pi_1(S)$ are either conformal or anti-conformally homeomorphic (by a homeomorphism in the homotopy class consistent with the identifications of their fundamental groups with $\pi_1(S)$.) Notice that $\partial_c N_i$ is identified with one component of the conformal boundary of N_i^S . If the conformal compactifications of N_1^S and N_2^S are conformally homeomorphic, then j extends to a conformal homeomorphism between $\partial_c N_1$ and $\partial_c N_2$, so Bers' result implies that ρ_1 and ρ_2 are conjugate in $\mathrm{PSL}(2, \mathbb{C})$. If not, then one may use the Klein–Maskit combination theorems (see Maskit [22]) to combine ρ_1 and (a conjugate of) ρ_2 to produce the holonomy representation of a hyperbolic structure on the double of M . Mostow's Rigidity theorem [27] then implies that the involution of the double is an isometry, so the restriction of ρ_1 and ρ_2 to the boundary is Fuchsian, and the conformal structures on the boundary must agree.

In the general case, we must organize the components of the boundary into those where the conformal structures agree and those where they disagree and replace the use of Bers' theorem and Mostow's Rigidity theorem with applications of the Ending Lamination theorem [8,26].

Proof of Theorem 1.2. Suppose that $\rho_1 : \pi_1(M) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(M) \rightarrow \mathrm{PSL}(2, \mathbb{C})$ are two discrete faithful representations, such that $\ell_{\rho_1}(\alpha) = \ell_{\rho_2}(\alpha)$ if $\alpha \in \pi_1(M)$ is represented by a simple closed curve on ∂M . Let $N_i = \mathbb{H}^3 / \rho_i(\pi_1(M))$.

There exists a homotopy equivalence $h_i : M \rightarrow N_i$ in the homotopy class determined by ρ_i (where we regard ρ_i as an isomorphism between $\pi_1(M)$ and $\pi_1(N_i) = \rho_i(\pi_1(M))$). Bonahon's Tameness theorem [3] implies that N_i may be identified with the interior of a compact 3-manifold M_i . Since M is acylindrical, Johansson's Classification theorem [16] implies that h_i is homotopic to a homeomorphism. Therefore, there exists a homeomorphism $g : M_1 \rightarrow M_2$ so that $g \circ h_1$ is homotopic to h_2 , i.e. g is in the homotopy class determined by $\rho_2 \circ \rho_1^{-1}$. If g is not orientation-preserving then we replace ρ_2 with $\bar{\rho}_2$. So, we may assume that g is orientation-preserving.

Thurston associates to each ρ_i a family of ending invariants, i.e. a multicurve p_i , called the parabolic locus, in the collection $\partial_0 M_i$ of non-toroidal components of ∂M_i , and on each component of $\partial_0 M_i - p_i$, either a hyperbolic structure of finite area (in which case the component is called geometrically finite) or a filling, geodesic lamination which admits a transverse measure of full support, called the ending lamination (in which case the component is called geometrically infinite), see [8, 26] for details. The Ending Lamination theorem of Minsky [26] and Brock–Canary–Minsky [8] implies that ρ_1 is conjugate to ρ_2 in $\mathrm{PSL}(2, \mathbb{C})$ if and only if the ending invariants of ρ_1 agree with the ending invariants of ρ_2 , i.e., up to isotopy of g , $g(p_1) = p_2$, the restriction of g to each geometrically finite component of $\partial_0 M_1 - p_1$ is an isometry onto a geometrically finite component of $\partial_0 M_2 - p_2$, and if a geometrically infinite component of $\partial_0 M_1 - p_1$ has ending lamination λ , then $g(\lambda)$ is the ending lamination of a geometrically infinite component of $\partial_0 M_2 - p_2$.

Let $\{S_1, \dots, S_r\}$ be the components of $\partial_0 M_1$. Since $\ell_{\rho_1}(\alpha) = \ell_{\rho_2}(\alpha)$ if $\alpha \in \pi_1(M)$ is represented by a simple closed curve on ∂M , our main result, Theorem 1.1, implies that, for each j , $\rho_1|_{\pi_1(S_j)}$ is conjugate to either ρ_2 or $\bar{\rho}_2$. If each $\rho_1|_{\pi_1(S_j)}$ is conjugate to $\rho_2|_{\pi_1(S_j)}$, then the ending invariants of ρ_1 agree with the ending invariants of ρ_2 , so ρ_1 is conjugate to ρ_2 in $\mathrm{PSL}(2, \mathbb{C})$ and we are done.

If there exists S_j so that $\rho_1|_{\pi_1(S_j)}$ is conjugate to $\bar{\rho}_2|_{\pi_1(S_j)}$, then we may re-order the components of ∂M , so that $\rho_1|_{\pi_1(S_j)}$ is conjugate to $\rho_2|_{\pi_1(S_j)}$ if and only if $j > q$ for some $q \in \{1, \dots, r-1\}$. Let N_i^j be the cover of N_i associated to $\pi_1(S_j)$. Then, N_i^j may be identified with the interior of $S_j \times [0, 1]$ so that the end invariants of N_i^j on $S_j \times \{1\}$ agree with the restriction of the end invariants of N_i restricted to S_j . The Covering theorem (see [10]) may be used to show that the restriction of the ending invariant of N_i^j to $S_j \times \{0\}$ is a conformal structure on all of $S_j \times \{0\}$ (see the discussion in Kent's proof of Thurston's Bounded Image theorem [17, Thm. 41]). If $j \leq q$, then the ending invariant of N_1^j on $S_j \times \{1\}$ agree with the end invariants of N_2^j on $S_j \times \{0\}$, while the ending invariant of N_1^j on $S_j \times \{0\}$ agree with the end invariants of N_2^j on $S_j \times \{1\}$. It follows that, if $j \leq q$, then the ending invariants of N_i^j are a pair of hyperbolic structures on $S_j \times \{0\}$ and $S_j \times \{1\}$, so $\rho_i|_{\pi_1(S_j)}$ is quasifuchsian, and the map from $S_j \times \{0, 1\}$ to itself given by taking (x, t) to $(x, 1-t)$

is isotopic to an isometry from the ending invariants of N_1^j to the ending invariants of N_2^j .

Let \hat{M} be the manifold obtained by doubling M along $S_1 \cup \cdots \cup S_{q-1}$. We first conjugate $\bar{\rho}_2$ so that $\rho_1(\pi_1(S_1)) = \bar{\rho}_2(\pi_1(S_1))$. We may use the Klein–Maskit Combination theorems [22] to see that the combination of $\rho_1(\pi_1(M))$ and $\bar{\rho}_2(\pi_1(M))$ gives rise to a hyperbolic structure on the double of M along S_1 . For each $j = 2, \dots, q$, let $A_j \in \mathrm{PSL}(2, \mathbb{C})$ be chosen so that A_j conjugates $\rho_1|_{\pi_1(S_j)}$ to $\rho_2|_{\pi_1(S_j)}$. Then, the Klein–Maskit Combination theorems give that the group generated by $\rho_1(\pi_1(M))$, $\rho_2(\pi_1(M))$ and $\{A_2, \dots, A_q\}$ gives rise to a hyperbolic structure on \hat{M} and so a representation $\hat{\rho} : \pi_1(\hat{M}) \rightarrow \mathrm{PSL}(2, \mathbb{C})$. (See Sections 8 and 9 of Morgan [25], in particular Theorem 9.4, for a discussion of the Klein–Maskit Combination theorems in a topological phrasing compatible with our application.) The obvious involution of \hat{M} preserves the ending invariants of $\hat{N} = \mathbb{H}^3 / \hat{\rho}(\pi_1(\hat{M}))$, so, by the Ending Lamination theorem, there is an isometry of \hat{N} realizing this involution. By restriction, ρ_1 is conjugate to ρ_2 . \square

Remark. In the case that $\partial_0 M$ is connected, Ian Agol astutely pointed out that one may rephrase this proof as an application of the fact that the square of the skinning map has a unique fixed point, see Morgan [25] and Kent [17] for a discussion of the skinning map.

5. Renormalized pressure intersection

We next show that the isometry group of the renormalized pressure intersection is generated by the (extended) mapping class group and complex conjugation. We begin by reinterpreting our renormalized pressure intersection in terms of the Patterson–Sullivan geodesic current, following Bridgeman [5] and Bridgeman–Taylor [7]. Since isometries of the renormalized pressure intersection are isometries of the pressure metric and the only degenerate vectors for the pressure metric are at points on the Fuchsian locus (see Bridgeman [5]), any isometry f of the renormalized pressure intersection must preserve the Fuchsian locus. Since the restriction of the pressure metric to the Fuchsian locus is a multiple of the Weil–Petersson metric, we may apply Masur and Wolf’s classification of isometries of the Weil–Petersson metric [23], to conclude that the restriction of f to the Fuchsian locus agrees with the action of a mapping class g . We then use Bonahon’s interpretation (see [4]) of the Thurston compactification of Teichmüller space in terms of geodesic currents, to show that if $\rho \in \mathcal{QF}(S)$, then $f(\rho)$ and $g(\rho)$ have proportional simple marked length spectrum. The proof is completed by showing that any two Kleinian surface groups with proportional simple marked length spectrum are conjugate in the isometry group of \mathbb{H}^3 .

5.1. Geodesic currents. Identify the space $G(\mathbb{H}^2)$ of geodesics in \mathbb{H}^2 with the open Möbius band $(\mathbb{S}_\infty^1 \times \mathbb{S}_\infty^1 - \text{diagonal})/\mathbb{Z}_2$ by identifying a geodesic with its endpoints. A *geodesic current* on a hyperbolic surface $X = \mathbb{H}^2/\Gamma$ is a positive Borel measure on $G(\mathbb{H}^2)$ that is invariant under the action of Γ . For example, if α is a primitive closed geodesic on X , we may consider the geodesic current C_α given by taking the Dirac measure on the pre-image of α in $G(\mathbb{H}^2)$. Let $\mathcal{C}(X)$ be the space of geodesic currents defined for $X = \mathbb{H}^2/\Gamma$ with the weak*-topology. If S is a closed surface of genus at least two, we identify S with a fixed hyperbolic surface X_0 and let $\mathcal{C}(S) = \mathcal{C}(X_0)$.

Following Bonahon [4], Bridgeman and Taylor [7, Sec. 5] showed that there is a natural continuous, linear function

$$L : \mathcal{C}(S) \times QF(S) \rightarrow \mathbb{R},$$

called the *length function*, with the property that if rC_α is a positive multiple of C_α , then $L(rC_\alpha, \rho) = r\ell_\rho(\alpha)$. Since multiples of closed geodesics are dense in $\mathcal{C}(S)$ (see [3, Prop. 4.2]), this property completely determines L .

Given $\rho \in QF(S)$, Sullivan [31, Prop. 11, Thm. 21], following work of Patterson [28] in the Fuchsian case, used the Poincaré series to define a non-atomic Γ -invariant measure μ_ρ on $\partial_\infty \mathbb{H}^3 \times \partial_\infty \mathbb{H}^3$, called the *Patterson–Sullivan measure*, which is supported on $\Lambda(\rho) \times \Lambda(\rho)$, where $\Lambda(\rho)$ is the limit set of $\rho(\pi_1(S))$, such that Γ acts ergodically on $\Lambda(\rho) \times \Lambda(\rho)$ with respect to μ_ρ (see also Sullivan [32, Thm. 3]). One may push forward μ_ρ to obtain a Γ -invariant measure $\hat{\mu}_\rho$ on $\Lambda(\rho) \times \Lambda(\rho)/\mathbb{Z}_2$ (where \mathbb{Z}_2 acts by interchanging the factors). The representation ρ induces a homeomorphism $f_\rho : S^1 \rightarrow \Lambda(\rho)$, where we use our identification of S with X_0 to identify S^1 with $\partial_\infty \mathbb{H}^2$, so one obtains a homeomorphism $\widehat{f_\rho \times f_\rho} : G(\mathbb{H}^2) \rightarrow \Lambda(\rho) \times \Lambda(\rho)/\mathbb{Z}_2$. One then defines the *Patterson–Sullivan current* as the scalar multiple $\psi(\rho) \in \mathcal{C}(S)$ of $\widehat{(f_\rho \times f_\rho)_*}(\hat{\mu}_\rho)$ so that $L(\psi(\rho), \rho) = 1$. If ρ is Fuchsian, then $\psi(\rho)$ is the Liouville geodesic current constructed by Bonahon [4] (see [15, Thm. 1]). Hamenstadt [15, Thm. 1] showed that the associated map

$$\psi : QF(S) \rightarrow \mathcal{C}(S)$$

is continuous and injective. See Bridgeman [5], Bridgeman–Taylor [7] or Hamenstadt [15] for further discussion of the Patterson–Sullivan geodesic current.

Let $F(S)$ denote the space of Fuchsian representations in $QF(S)$ and let $\mathcal{PC}(S)$ denote the space of projective classes of non-zero currents in $\mathcal{C}(S)$. Bonahon [4, Thm. 18] showed that if one considers the associated map into the space of projective geodesic currents $\bar{\psi} : F(S) \rightarrow \mathcal{PC}(S)$, then the closure of $\bar{\psi}(F(S))$ is $\bar{\psi}(F(S)) \cup PML(S)$ where $PML(S)$ is the space of projective classes of measured laminations. (We recall that the set $ML(S)$ of measured laminations in $\mathcal{C}(S)$ is exactly the closure of the set of positive multiples of currents associated to simple curves, see Bonahon [3, Sec. 4.3] for more details.)

We note that $L(\psi(\rho_1), \rho_2)$ can also be defined to be the length in ρ_2 of a random geodesic in ρ_1 , i.e.

$$L(\psi(\rho_1), \rho_2) = \lim_{T \rightarrow \infty} \frac{1}{\#\{[\gamma] \mid \ell_{\rho_1}(\gamma) \leq T\}} \sum_{\{[\gamma] \mid \ell_{\rho_1}(\gamma) \leq T\}} \frac{\ell_{\rho_2}(\gamma)}{\ell_{\rho_1}(\gamma)}. \quad (5.1)$$

It follows that

$$\mathbf{J}(\rho_1, \rho_2) = \frac{h(\rho_2)}{h(\rho_1)} L(\psi(\rho_1), \rho_2), \quad (5.2)$$

where the the renormalized intersection number \mathbf{J} and the entropy h are as defined in Section 1.3.

In analogy with Thurston's formulation of the Weil–Petersson metric on Teichmüller space, and Bonahon's re-formulation in terms of geodesic currents [4, Thm. 19], Bridgeman and Taylor [7] consider, for each $\rho \in QF(S)$, the function $\mathbf{J}_\rho : QF(S) \rightarrow \mathbb{R}$ given by $\mathbf{J}_\rho(\sigma) = \mathbf{J}(\rho, \sigma)$ and define a non-negative symmetric two-tensor G , called the *pressure form*, on the tangent bundle $TQF(S)$, where G_ρ on $T_\rho QF(S)$ is the Hessian of \mathbf{J}_ρ . By construction, the pressure form is invariant under the action of the (extended) mapping class group $\text{Mod}^*(S)$ of S . It follows from the work of Wolpert [33] and Bonahon [4], that the restriction of the pressure form to the Fuchsian locus $F(S)$ is a multiple of the Weil–Petersson metric. Motivated by the work of McMullen [24] in the setting of Teichmüller space, Bridgeman [5] showed that the pressure form is non-degenerate except along pure bending vectors based at points in the Fuchsian locus. Moreover, the pressure form gives rise to a path metric on $QF(S)$, see [6, Cor. 1.7].

Remark. In the proof of Theorem 6.1 in [5], Bridgeman gives an expression for $L(\psi(\rho_1), \rho_2)$ in terms of equilibrium measures. Our equation (5.1) then follows from equations (12) and (13) in [6], see also the discussion in Section 8 of [6]. If one prefers that the proof of Theorem 1.3 be self-contained, one can take J to be defined by our equation (5.2) as is done in Bridgeman–Taylor [7] and Bridgeman [5].

5.2. Isometries of renormalized pressure intersection. We use the interpretation of renormalized pressure intersection in terms of geodesic currents to show that the restriction of an isometry f of the renormalized pressure intersection to the Fuchsian locus $F(S)$ agrees with an element g of the (extended) mapping class group. We further show that $f(\sigma)$ and $g(\sigma)$ have closely related length spectrum whenever $\sigma \in QF(S)$.

Proposition 5.1. *If $f : QF(S) \rightarrow QF(S)$ is a smooth isometry of \mathbf{J} , then f preserves the Fuchsian locus $F(S)$ and there exists $g \in \text{Mod}^*(S)$ such that f and g agree on the Fuchsian locus.*

Moreover, if α is a simple curve on S and $\sigma \in QF(S)$, then

$$h(g(\sigma))\ell_{g(\sigma)}(\alpha) = h(f(\sigma))\ell_{f(\sigma)}(\alpha).$$

Proof. Since f is smooth and preserves \mathbf{J} , one sees by differentiating that f also preserves the pressure form G .

If $\sigma \in F(X)$ and v is a pure-bending vector, then $G_\sigma(v, v) = 0$, so $G_{f(\sigma)}(Df_\sigma(v), Df_\sigma(v)) = 0$. Since, f is an immersion, $Df_\sigma(v) \neq 0$ which implies that $f(\sigma) \in F(S)$.

Since the restriction of G to $F(S)$ is a multiple of the Weil–Petersson metric, the restriction of f to the Fuchsian locus is an isometry of the Weil–Petersson metric. Masur and Wolf [23] proved every isometry of the Weil–Petersson metric is an element of $\text{Mod}^*(S)$. So we may choose $g \in \text{Mod}^*(S)$ so that f and g agree on the Fuchsian locus $F(S)$.

If $\rho \in F(S)$ and $\sigma \in QF(S)$, then since

$$\mathbf{J}(\rho, \sigma) = \mathbf{J}(f(\rho), f(\sigma)) = \mathbf{J}(g(\rho), g(\sigma)),$$

and h is constant on $F(S)$, we may conclude that

$$\frac{L(\psi(g(\rho)), g(\sigma))}{L(\psi(f(\rho), f(\sigma)))} = \frac{h(f(\sigma))}{h(g(\sigma))}.$$

If α is a simple curve, let $\{\rho_n\}$ be a sequence in $F(S)$ so that $\{\bar{\psi}(f(\rho_n))\} = \{\bar{\psi}(g(\rho_n))\}$ converges to $[C_\alpha]$ in $\mathcal{PC}(S)$. Since L is continuous,

$$\frac{\ell_{g(\sigma)}(\alpha)}{\ell_{f(\sigma)}(\alpha)} = \lim \frac{L(\psi(g(\rho_n)), g(\sigma))}{L(\psi(f(\rho_n)), f(\sigma))} = \frac{h(f(\sigma))}{h(g(\sigma))}$$

which establishes our claim. \square

5.3. Simply k -related Kleinian surface groups. We say that two Kleinian surface groups $\rho_1 : \pi_1(S) \rightarrow \text{PSL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(S) \rightarrow \text{PSL}(2, \mathbb{C})$ are *simply k -related*, for some $k > 0$, if

$$\ell_{\rho_1}(\alpha) = k \ell_{\rho_2}(\alpha)$$

whenever α is a simple closed curve on S .

Proposition 5.1 implies that if f is an isometry of the renormalized pressure intersection, then there exists $g \in \text{Mod}^*(S)$, such that $f(\sigma)$ and $g(\sigma)$ are simply k -related whenever $\sigma \in QF(S)$. Theorem 1.3 will thus follow from the following strengthening of our simple length rigidity theorem.

Theorem 1.4. *If S is a closed, connected, orientable surface of genus at least two and $\rho_1 : \pi_1(S) \rightarrow \text{PSL}(2, \mathbb{C})$ and $\rho_2 : \pi_1(S) \rightarrow \text{PSL}(2, \mathbb{C})$ are simply k -related Kleinian surface groups, then ρ_1 is conjugate to either ρ_2 or $\bar{\rho}_2$.*

Proof of Theorem 1.4. We first choose non-separating simple curves α and β on S which intersect exactly once, so that $\rho_1(\alpha)$, $\rho_1(\beta)$, $\rho_2(\alpha)$ and $\rho_2(\beta)$ are all hyperbolic.

(One may do so, since, again by a result of Sullivan [30], only finitely many simple curves have parabolic images for ρ_1 or ρ_2 .)

We can assume that ρ_1 and ρ_2 are (α, β) -normalized, so

$$\rho_i(\alpha) = \pm \begin{pmatrix} \lambda_i & 0 \\ 0 & \lambda_i^{-1} \end{pmatrix}$$

where $|\lambda_i| > 1$, and

$$\rho_i(\beta) = \pm \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix}$$

where all the matrix coefficients are non-zero (see Lemma 2.2). Since ρ_1 and ρ_2 are simply k -related,

$$|\lambda_1| = |\lambda_2|^k.$$

Notice that if $k = 1$, then Theorem 1.4 follows immediately from our main result, Theorem 1.1. So we may assume, without loss of generality, that $k > 1$.

Let $\mu_i(n)$ be the modulus of the eigenvalue of $\rho_i(\alpha^n \beta)$ with largest modulus. Since $\alpha^n \beta$ is simple for all n , $\mu_1(n) = \mu_2(n)^k$ for all n . Lemma 2.2 implies that

$$\log \mu_i(n) = n \log |\lambda_i| + \log |a_i| + \Re \left(\lambda_i^{-2n} \frac{b_i c_i}{a_i^2} \right) + O(|\lambda_i|^{-4n}).$$

Since $\log \mu_1(n) = k \log \mu_2(n)$, after eliminating the leading terms (which are equal), we obtain

$$\begin{aligned} \log |a_1| + \Re \left(\lambda_1^{-2n} \frac{b_1 c_1}{a_1^2} \right) + O(|\lambda_1|^{-4n}) \\ = k \log |a_2| + k \Re \left(\lambda_2^{-2n} \frac{b_2 c_2}{a_2^2} \right) + O(|\lambda_2|^{-4n}). \end{aligned}$$

Therefore, by considering the limits as $n \rightarrow \infty$, we see that

$$\log |a_1| = k \log |a_2|.$$

So, after subtracting the equal leading order terms,

$$\Re \left(\lambda_1^{-2n} \frac{b_1 c_1}{a_1^2} \right) + O(|\lambda_1|^{-4n}) = k \Re \left(\lambda_2^{-2n} \frac{b_2 c_2}{a_2^2} \right) + O(|\lambda_2|^{-4n}).$$

for all n . If we multiply both sides of the above equation by $|\lambda_2|^{2n}$, then since $\frac{|\lambda_1|}{|\lambda_2|} > 1$, the left hand side converges to 0, and we see that

$$\lim_{n \rightarrow \infty} \Re \left(\left(\frac{\lambda_2}{|\lambda_2|} \right)^{-2n} \frac{b_2 c_2}{a_2^2} \right) = 0. \quad (5.3)$$

If we let

$$u = \left(\frac{\lambda_2}{|\lambda_2|} \right)^{-2} = e^{i\theta} \quad \text{and} \quad v = \frac{b_2 c_2}{a_2^2} \neq 0.$$

Then we may rewrite (5.3) as

$$\lim_{n \rightarrow \infty} \Re(u^n v) = 0.$$

As we can always choose a sequence $\{n_k\}$ so that $\lim_{k \rightarrow \infty} u^{n_k} = 1$, we see that

$$\Re(v) = 0.$$

If θ is an irrational multiple of 2π , then for all $z \in S^1$ there exists a sequence $\{n_k\}$ so that $\lim e^{n_k \theta} = z$, which implies that $\Re(zv) = 0$. Therefore, in this case, as $v \neq 0$ then choosing $z = |v|/v$, we get $|v| = 0$ a contradiction.

If θ is a rational multiple of 2π and $u \neq \pm 1$, then there exists $\{n_k\}$ so that $u^{n_k} = u$ for all k , so $\Re(uv) = 0$. Since $\Re(v) = 0$, $v = iy$ from some $y \in \mathbb{R}$, so,

$$\Re(e^{i\theta} v) = -\sin(\theta)y = 0.$$

Thus, $y = 0$, since $\sin(\theta) \neq 0$. It again follows that $v = 0$ and we obtain a contradiction.

Therefore, we can assume that $u = \pm 1$, so λ_2^2 is real. It follows that if η is a simple, non-separating curve on S , then $\lambda_{\rho_2}^2(\eta) \in \mathbb{R}$, so

$$\text{Tr}^2(\rho_2(\eta)) = \lambda_{\rho_2}^2(\eta) + 2 + \lambda_{\rho_2}^{-2}(\eta) \in \mathbb{R}.$$

In particular, since $\alpha^n \beta$ is simple and non-separating for all n , we see that

$$\text{Tr}^2(\rho_2(\alpha^n \beta)) = \lambda_2^{2n} a_2^2 + 2a_2 d_2 + \lambda_2^{-2n} d_2^2 \in \mathbb{R},$$

so,

$$\Im(\text{Tr}^2(\rho_2(\alpha^n \beta))) = \lambda_2^{2n} \Im(a_2^2) + 2\Im(a_2 d_2) + \lambda_2^{-2n} \Im(d_2^2) = 0$$

for all n . Therefore,

$$\lim_{n \rightarrow \infty} \frac{\Im(\text{Tr}^2(\rho_2(\alpha^n \beta)))}{\lambda_2^{2n}} = \Im(a_2^2) = 0,$$

so

$$\lim_{n \rightarrow \infty} \Im(\text{Tr}^2(\rho_2(\alpha^n \beta))) = 2\Im(a_2 d_2) = 0.$$

Since a_2^2 and $a_2 d_2$ are real, $v = \frac{b_2 c_2}{a_2^2} = \frac{a_2 d_2 - 1}{a_2^2}$ is real. Therefore, since we have already shown that $\Re(v) = 0$, we see that $v = 0$, which contradicts the fact that a_2 , b_2 and c_2 are all non-zero. Since we have now achieved a contradiction in all cases where $k > 1$, this completes the proof. \square

5.4. Proof of Theorem 1.3. If f is an isometry of the renormalized pressure intersection, then Proposition 5.1 implies that there exists $g \in \text{Mod}^*(S)$ such that f and g agree on $F(S)$ and $f(\sigma)$ and $g(\sigma)$ are simply k -related, where $k = \frac{h(f(\sigma))}{h(g(\sigma))}$, whenever $\sigma \in QF(S)$. Theorem 1.4 implies that if $\sigma \in QF(S)$, then either $f(\sigma) = g(\sigma)$ or $f(\sigma) = \tau(g(\sigma))$. Since f is continuous, this implies that either $f = g$ or $f = \tau \circ g$. The result follows. \square

References

- [1] L. Bers, Simultaneous uniformization, *Bull. Amer. Math. Soc.*, **66** (1960), 94–97. Zbl 0090.05101 MR 111834
- [2] L. Bers, On moduli of Kleinian groups, *Russian Math. Surveys*, **29** (1974), 88–102. Zbl 0304.30013 MR 422691
- [3] F. Bonahon, Bouts des variétés hyperboliques de dimension 3, *Annals of Math.*, **124** (1986), 71–158. Zbl 0671.57008 MR 847953
- [4] F. Bonahon, The geometry of Teichmüller space via geodesic currents, *Invent. Math.* **92**(1988), 139–162. Zbl 0653.32022 MR 931208
- [5] M. Bridgeman, Hausdorff dimension and the Weil–Petersson extension to quasifuchsian space, *Geom. Top.*, **14** (2010), 799–831. Zbl 1200.30037 MR 2602852
- [6] M. Bridgeman, R. Canary, F. Labourie, and A. Sambarino, The pressure metric for Anosov representations, *Geom. Funct. Anal.*, **25** (2015), no. 4, 1089–1179. Zbl 1360.37078 MR 3385630
- [7] M. Bridgeman and E. Taylor, An extension of the Weil–Petersson metric to quasi-fuchsian space, *Math Ann.*, **341** (2008), 927–943. Zbl 1176.30095 MR 2407332
- [8] J. F. Brock, R. D. Canary and Y. N. Minsky, The Classification of Kleinian Surface Groups II: The Ending Lamination Conjecture, *Annals of Math.*, **176** (2012), 1–149. Zbl 1253.57009 MR 2925381
- [9] M. Burger, Intersection, the Manhattan curve and Patterson–Sullivan theory in rank 2, *Internat. Math. Res. Notices*, **7** (1993), 217–225. Zbl 0829.57023 MR 1230298
- [10] R. D. Canary, A covering theorem for hyperbolic 3-manifolds and its applications, *Topology*, **35** (1996), 751–778. Zbl 0863.57010 MR 1396777
- [11] L. Charles and J. Marché, Multicurves and regular functions on the representation variety of a surface in $SU(2)$, *Comm. Math. Helv.*, **87** (2012), 409–431. Zbl 1246.57022 MR 2914854
- [12] M. Culler, Lifting representations to covering groups, *Adv. Math.*, **59** (1986), 64–70. Zbl 0582.57001 MR 825087
- [13] M. Culler and P. Shalen, Varieties of group representations and splittings of 3-manifolds, *Annals of Math.*, **117** (1983), 109–146. Zbl 0529.57005 MR 683804
- [14] F. Dal’Bo and I. Kim, Marked length rigidity for symmetric spaces, *Comm. Math. Helv.*, **77** (2002), 399–407. Zbl 1002.22005 MR 1915048
- [15] U. Hamenstädt, Ergodic properties of function groups, *Geom. Ded.*, **93** (2002), 163–176. Zbl 1014.37021 MR 1934696

- [16] K. Johansson, *Homotopy Equivalences of 3-manifolds with Boundary*, Lecture Notes in Mathematics, 761, Springer-Verlag, 1979. Zbl 0412.57007 MR 551744
- [17] A. E. Kent, Skinning maps, *Duke Math. J.*, **151** (2010), 279–336. Zbl 1193.30062 MR 2598379
- [18] I. Kim, Marked length rigidity of rank one symmetric spaces and their product, *Topology*, **40** (2001), 1295–1323. Zbl 0997.53034 MR 1867246
- [19] C. Kourouniotis, Complex length coordinates for quasi-fuchsian groups, *Mathematika*, **41** (1994), 173–188. Zbl 0801.30036 MR 1288062
- [20] I. Kra, On lifting Kleinian groups to $SL(2, \mathbb{C})$, in *Differential geometry and complex analysis*, 181–193, Springer-Verlag, 1985. Zbl 0571.30037 MR 780044
- [21] J. Marché and M. Wolff, The modular action on $PSL(2, \mathbb{R})$ -characters in genus 2, *Duke Math. J.*, **165** (2016), 371–412. Zbl 1353.37056 MR 3457677
- [22] B. Maskit, On Klein’s combination theorem. III, in *Advances in the Theory of Riemann Surfaces (Proc. Conf., Stony Brook, N.Y., 1969)*, 297–316, Ann. of Math. Studies, 66, Princeton Univ. Press, Princeton, N.J., 1971. Zbl 0222.30029 MR 289768
- [23] H. Masur and M. Wolf, The Weil–Petersson isometry group, *Geom. Ded.*, **93** (2002), 177–190. Zbl 1014.32008 MR 1934697
- [24] C. McMullen, Thermodynamics, dimension and the Weil–Petersson metric, *Invent. Math.*, **173** (2008), 365–425. Zbl 1156.30035 MR 2415311
- [25] J. Morgan, On Thurston’s uniformization theorem for three dimensional manifolds, in *The Smith conjecture (New York, 1979)*, 37–125, Pure Appl. Math., 112, Academic Press, Orlando, FL, 1984. Zbl 0599.57002 MR 758464
- [26] Y. Minsky, The classification of Kleinian surface groups. I: Models and bounds, *Annals of Math.*, **171** (2010), 1–107. Zbl 1193.30063 MR 2630036
- [27] G. D. Mostow, Quasiconformal mappings in n -space and the rigidity of hyperbolic space forms, *Publ. I.H.E.S.*, **34** (1968), 53–104. Zbl 0189.09402 MR 236383
- [28] S. Patterson, The limit set of a fuchsian group, *Acta Math.*, **136** (1976), 241–273. Zbl 0336.30005 MR 450547
- [29] H. Royden, Automorphisms and isometries of Teichmüller spaces, in *Advances in the Theory of Riemann Surfaces (Proc. Conf., Stony Brook, N.Y., 1969)*, 369–383, Ann. of Math. Studies, 66, Princeton Univ. Press, Princeton, N.J., 1971. Zbl 0218.32011 MR 288254
- [30] D. Sullivan, A finiteness theorem for cusps, *Acta Math.*, **147** (1981), 289–299. Zbl 0502.57004 MR 639042
- [31] D. Sullivan, The density at infinity of a discrete group of hyperbolic motions, *Publ. Math. I.H.E.S.*, **50** (1979), 171–202. Zbl 0439.30034 MR 556586
- [32] D. Sullivan, Entropy, Hausdorff measures old and new, and limit sets of geometrically finite Kleinian groups, *Acta Math.*, **153** (1984), 259–277. Zbl 0566.58022 MR 766265
- [33] S. Wolpert, Thurston’s Riemannian metric for Teichmüller space, *J. Diff. Geom.*, **23** (1986), 143–174. Zbl 0592.53037 MR 845703

Received September 29, 2015

M. Bridgeman, Department of Mathematics, Boston College,
Chestnut Hill, MA 02467, USA

E-mail: bridgem-at-bc.edu

R. Canary, Department of Mathematics, University of Michigan,
Ann Arbor, MI 48109-1043, USA

E-mail: canary@umich.edu