

The generating identity of Cauchy-Schwarz-Bunyakovsky inequality

Autor(en): **Sommariva, Antonino M.**

Objekttyp: **Article**

Zeitschrift: **Elemente der Mathematik**

Band (Jahr): **63 (2008)**

PDF erstellt am: **25.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-99055>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

The generating identity of Cauchy-Schwarz-Bunyakovsky inequality

Antonino M. Sommariva

Antonino M. Sommariva received a “Laurea in Ingegneria Elettronica” degree from the University of Palermo in 1977. Currently, he is a full professor in the “Dipartimento di Elettronica per l’Automazione” at the University of Brescia (Italy). His main research interests are perturbation methods, and the fundamental problems of classical circuit theory.

1 Introduction

The Cauchy-Schwarz-Bunyakovsky inequality (CSB-inequality for short) is well-known to pure and applied mathematicians for its application in different areas of mathematics (algebra, analysis, geometry, probability theory, etc.). A complete account of its both intricate and intriguing history can be found in a well documented paper by Schreiber [3] or in the delightful book by Steele [4].

This important relationship can be formulated as follows:

CSB-inequality – *Let V be a real or complex inner product vector space, and let x, y be elements of V . Then*

$$|x \cdot y| \leq \|x\| \|y\|$$

where equality holds if and only if x and y are linearly dependent.

Was Sie immer schon über die Schwarzsche Ungleichung wissen wollten, sich aber nie zu fragen getrauteten: Wo bleibt eigentlich die Differenz? Die Beweise in den Lehrbüchern der linearen Algebra oder der Funktionalanalysis sind auf Knappheit getrimmt und geben darüber keine Auskunft. Es gibt allerdings eine Lagrange zugeschriebene Identität, die diese Differenz als Quadrat eines gewissen schiefsymmetrischen Ausdrucks darstellt. Der Autor der vorliegenden Arbeit ist der Sache auf den Grund gegangen und findet im Rahmen des Tensorprodukts $V \otimes V$ eine einleuchtende und allgemeingültige Erklärung.

To this author's best knowledge, identities giving rise to the CSB-inequality have been obtained in the past, but for particular cases only. Thus, Cauchy derived his version of the inequality by deleting some summands in an arithmetical identity ascribed to Lagrange, i.e.

$$(a\alpha + a'\alpha' + a''\alpha'' + \dots)^2 + (a\alpha' - a'\alpha)^2 + (a\alpha'' - a''\alpha)^2 + (a'\alpha'' - a''\alpha')^2 + \dots \\ = (a^2 + a'^2 + a''^2 + \dots)^2 (\alpha^2 + \alpha'^2 + \alpha''^2 + \dots)^2$$

where $a, a', a'', \dots, \alpha, \alpha', \alpha'', \dots$ are two sequences of n real numbers each. The same identity appears (adopting the self-evident notation of [2, pp. 33, 62]) in exterior algebra or the theory of spatial vectors as

$$(\mathbf{a} \cdot \mathbf{b})^2 + |\mathbf{a} \wedge \mathbf{b}|^2 = \mathbf{a}^2 \mathbf{b}^2 \quad \text{or} \quad (\mathbf{a} \cdot \mathbf{b})^2 + |\mathbf{a} \times \mathbf{b}|^2 = \mathbf{a}^2 \mathbf{b}^2$$

respectively.

In the context of functional analysis, Courant and Hilbert [1, p. 49] reported (without reference) an identity which can be rewritten as

$$(f, g)^2 + \frac{1}{2} \int \int (f(x)g(\xi) - f(\xi)g(x))^2 dx d\xi = (f, f)(g, g)$$

where f and g are real-valued functions which are piecewise continuous in some finite interval $G \subset \mathbb{R}$.

This paper proposes an identity which gives rise to the CSB-inequality for the general case, based on the concept of tensor product of vector spaces. That is, first, a fundamental relationship is established between the inner product in a vector space V and a particular inner product in the tensor product of V by itself. Then, a Pythagorean relation, involving symmetric and antisymmetric parts of a tensor, is derived and manipulated to obtain the desired identity. Finally, two examples are presented.

2 The identity behind the CSB-inequality

Let V be a real or complex inner product vector space, and let $V \otimes V$ be the tensor product of V by itself. If $\{h_m\}$ is any basis of V , then $\{h_m \otimes h_n\}$ is a basis of $V \otimes V$, and it is referred to as the basis induced by $\{h_m\}$. An inner product can be introduced also in $V \otimes V$ by specifying proper values for the inner products of the induced basis elements. In particular, if the basis $\{h_m\}$ is orthonormal, i.e.,

$$h_p \cdot h_q = \delta_{pq},$$

and the basis $\{h_m \otimes h_n\}$ is assumed as orthonormal, i.e.,

$$(h_p \otimes h_q) \cdot (h_r \otimes h_s) = \delta_{pr} \delta_{qs},$$

then it is easily seen that for any $a, b, c, d \in V$ the following formula holds:

$$(a \otimes b) \cdot (c \otimes d) = (a \cdot c)(b \cdot d).$$

In fact, having set

$$a = \sum a_p h_p, \quad b = \sum b_q h_q, \quad c = \sum c_r h_r, \quad d = \sum d_s h_s,$$

it turns out that

$$\begin{aligned}
(a \otimes b) \cdot (c \otimes d) &= \left(\sum a_p h_p \otimes \sum b_q h_q \right) \cdot \left(\sum c_r h_r \otimes \sum d_s h_s \right) \\
&= \sum a_p b_q (h_p \otimes h_q) \cdot \sum c_r d_s (h_r \otimes h_s) \\
&= \sum a_p b_q \bar{c}_r \bar{d}_s (h_p \otimes h_q) \cdot (h_r \otimes h_s) \\
&= \sum a_p b_q \bar{c}_r \bar{d}_s \delta_{p,r} \delta_{q,s} = \sum a_p b_q \bar{c}_p \bar{d}_q = \sum a_p \bar{c}_p \sum b_q \bar{d}_q \\
&= (a \cdot c)(b \cdot d).
\end{aligned}$$

Under these assumptions, the symmetric and the antisymmetric part of the tensor product $x \otimes y$, denoted $\text{Sym}(x \otimes y)$ and $\text{Asym}(x \otimes y)$ respectively, are orthogonal:

$$\begin{aligned}
\text{Sym}(x \otimes y) \cdot \text{Asym}(x \otimes y) &= \frac{1}{2} (x \otimes y + y \otimes x) \cdot \frac{1}{2} (x \otimes y - y \otimes x) \\
&= \frac{1}{4} \left(\|x\|^2 \|y\|^2 - \|y\|^2 \|x\|^2 - (x \cdot y)(y \cdot x) + (y \cdot x)(x \cdot y) \right) = 0
\end{aligned}$$

and thus the Pythagorean relationship holds:

$$\|\text{Sym}(x \otimes y)\|^2 + \|\text{Asym}(x \otimes y)\|^2 = \|x \otimes y\|^2. \quad (1)$$

Moreover, one can obtain

$$\|x \otimes y\|^2 = (x \otimes y) \cdot (x \otimes y) = \|x\|^2 \|y\|^2 \quad (2)$$

and

$$\begin{aligned}
\|\text{Sym}(x \otimes y)\|^2 &= \frac{1}{2} (x \otimes y + y \otimes x) \cdot \frac{1}{2} (x \otimes y + y \otimes x) \\
&= \frac{1}{4} \left(\|x\|^2 \|y\|^2 + \|y\|^2 \|x\|^2 + (x \cdot y)(y \cdot x) + (y \cdot x)(x \cdot y) \right) \\
&= \frac{1}{2} (\|x\|^2 \|y\|^2 + |x \cdot y|^2). \quad (3)
\end{aligned}$$

Hence, substitution of (2) and (3) into (1) yields the wanted identity:

$$|x \cdot y|^2 + 2 \|\text{Asym}(x \otimes y)\|^2 = \|x\|^2 \|y\|^2. \quad (4)$$

According to (4), the term which completes the (squared) CSB-inequality can be interpreted and calculated as twice the squared norm of the antisymmetric part of the tensor product of the involved elements, provided that the basis induced by an orthonormal basis of V is assumed as an orthonormal basis of $V \otimes V$.

Sometimes, $V \otimes V$ has some given inner product, and it is desirable to use it. In this case, one must verify that the induced basis $\{h_m \otimes h_n\}$ is orthonormal with respect to this inner product.

3 Examples

In this section, two examples are considered which extend the results (Lagrange's identity, Courant-Hilbert's identity) mentioned in the introduction. In both of them, the tensor product $V \otimes V$ has a given usual inner product, and the basis of $V \otimes V$ induced by an orthonormal basis of V is orthonormal with respect to this inner product.

Let $\mathbb{C}^{M \times N}$ be the space of $M \times N$ matrices over \mathbb{C} , and let $\mathbb{C}^{M \times N}$ be endowed with its usual inner product, i.e., if $\mathbf{A}, \mathbf{B} \in \mathbb{C}^{M \times N}$,

$$\mathbf{A} \cdot \mathbf{B} := \text{trace}(\tilde{\mathbf{A}} \overline{\mathbf{B}})$$

where the tilde stands for transpose. For any $\mathbf{A}, \mathbf{B} \in \mathbb{C}^{M \times N}$, consider the column partitions

$$\mathbf{A} = [\mathbf{a}_1 \quad \mathbf{a}_2 \quad \cdots \quad \mathbf{a}_N], \quad \mathbf{B} = [\mathbf{b}_1 \quad \mathbf{b}_2 \quad \cdots \quad \mathbf{b}_N],$$

then set

$$\mathbf{A} \otimes \mathbf{B} := \begin{bmatrix} \mathbf{a}_1 \tilde{\mathbf{b}}_1 & \mathbf{a}_1 \tilde{\mathbf{b}}_2 & \cdots & \mathbf{a}_1 \tilde{\mathbf{b}}_N \\ \mathbf{a}_2 \tilde{\mathbf{b}}_1 & \mathbf{a}_2 \tilde{\mathbf{b}}_2 & \cdots & \mathbf{a}_2 \tilde{\mathbf{b}}_N \\ \dots & \dots & \dots & \dots \\ \mathbf{a}_N \tilde{\mathbf{b}}_1 & \mathbf{a}_N \tilde{\mathbf{b}}_2 & \cdots & \mathbf{a}_N \tilde{\mathbf{b}}_N \end{bmatrix}$$

and observe that $\mathbb{C}^{M \times N} \otimes \mathbb{C}^{M \times N} = \mathbb{C}^{(MN) \times (MN)}$. Therefore, according to (4), if $\mathbf{A} = (a_{pq})$ and $\mathbf{B} = (b_{rs})$, the term which completes the (squared) CSB-inequality in this example amounts to

$$\frac{1}{2} \sum_{p,q,r,s} |a_{pq} b_{rs} - b_{pq} a_{rs}|^2.$$

The real case of the above formula with $N = 1$ corresponds to Lagrange's identity.

Let $C_{\text{pw}}(G, \mathbb{C})$ be the space of complex-valued functions which are piecewise continuous in some finite interval $G \subset \mathbb{R}^N$, and let $C_{\text{pw}}(G, \mathbb{C})$ be endowed with its usual inner product, i.e. if $f, g \in C_{\text{pw}}(G, \mathbb{C})$

$$f \cdot g := \int \cdots \int f(x_1, \dots, x_N) \overline{g}(x_1, \dots, x_N) dx_1 \cdots dx_N.$$

For any $f, g \in C_{\text{pw}}(G, \mathbb{C})$, set

$$f \otimes g : G^2 \rightarrow \mathbb{C}$$

$$(f \otimes g)(x_1, \dots, x_N, \xi_1, \dots, \xi_N) := f(x_1, \dots, x_N) g(\xi_1, \dots, \xi_N)$$

and observe that $C_{\text{pw}}(G, \mathbb{C}) \otimes C_{\text{pw}}(G, \mathbb{C}) = C_{\text{pw}}(G^2, \mathbb{C})$. Therefore, according to (4), the term which completes the (squared) CSB-inequality in this example amounts to

$$\frac{1}{2} \int \cdots \int |f(x_1, \dots, x_N) g(\xi_1, \dots, \xi_N) - g(x_1, \dots, x_N) f(\xi_1, \dots, \xi_N)|^2 dx_1 \cdots d\xi_N.$$

The real case of the above formula with $N = 1$ corresponds to Courant-Hilbert's identity.

References

- [1] Courant, R.; Hilbert, D.: *Methods of Mathematical Physics I*. Wiley-Interscience, 1989.
- [2] Gröbner, W.: *Matrizenrechnung*. Bibliographisches Institut, Mannheim 1966.
- [3] Schreiber, P.: *The Cauchy-Bunyakovsky-Schwarz Inequality*. In: Hermann Graßmann – Werk und Wirkung. Ernst-Moritz-Arndt-Universität, Greifswald 1995, 64–70.
- [4] Steele, J.M.: *The Cauchy-Schwarz Master Class: An Introduction to the Art of Mathematical Inequalities*. MAA Problem Book Series, Cambridge 2004.

Antonino M. Sommariva
Università degli Studi di Brescia
Via Branze, 38
I-25123 Brescia, Italy
e-mail: sommariv@ing.unibs.it