

A new class of permutation polynomials of \mathbb{F}_q

Autor(en): **Ayad, Mohamed / Kihel, Omar**

Objekttyp: **Article**

Zeitschrift: **Elemente der Mathematik**

Band (Jahr): **68 (2013)**

PDF erstellt am: **26.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-515895>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

A new class of permutation polynomials of \mathbb{F}_q

Mohamed Ayad and Omar Kihel

Mohamed Ayad received his Ph.D. from Université de Caen in 1992. He taught in Algerian and French universities. Since 1994 he is Maître de Conférences at Université du Littoral (Calais, France). His main fields of research are number theory, arithmetic geometry, finite fields, and algorithmic algebra.

Omar Kihel received his Ph.D. from Université Laval (Canada) in 1996. He then had a postdoctoral position at CICMA (McGill University and Concordia University) and another postdoctoral position at Université Laval. Professor Omar Kihel joined Brock University (Ontario, Canada) in July 2002. His areas of research are algebraic number theory, elliptic curves and finite fields.

Let \mathbb{F}_q be the finite field of characteristic p containing $q = p^r$ elements. A polynomial $f(x) \in \mathbb{F}_q[x]$ is called a permutation polynomial of \mathbb{F}_q if the induced map $f : \mathbb{F}_q \rightarrow \mathbb{F}_q$ is one to one. The study of permutation polynomials goes back to Hermite [2] for \mathbb{F}_p and to Dickson [1] for \mathbb{F}_q . One of the open problems proposed by Lidl and Mullen [3], is to find new classes of permutation polynomials of \mathbb{F}_q . We refer to [4] or [5] for the basic results on permutation polynomials. Wan and Lidl [7], gave conditions on a polynomial of the form $x^r f(x^{(q-1)/d})$ to be a permutation polynomial. The conditions are not explicitly given in terms of q and r , and may be difficult to verify in general. In the present note, without using the characterization of Wan and Lidl [7], but using only an elementary method, we exhibit a new class of permutation polynomials. We prove the following:

Theorem 1 *Let $q = p^r$, where p is a prime number and r is a positive integer. Let u be a positive integer and let*

$$f(x) = x^u \left(x^{\frac{q-1}{2}} + x^{\frac{q-1}{4}} + 1 \right). \quad (1)$$

Unter einem Permutationspolynom des kommutativen Rings R mit Einselement versteht man ein Polynom $p \in R[x]$, für welches die durch $\pi\alpha := p(\alpha)$ definierte Abbildung von R nach R eine Permutation der Ringelemente ist. Permutationspolynome sind beliebte Studienobjekte der Zahlentheorie, der Algebra und der Kombinatorik. Am besten untersucht ist wohl der Fall, wenn R ein endlicher Körper ist. Die Autoren der vorliegenden Arbeit steuern zu dieser Theorie eine neue, einfache Klasse von Permutationspolynomen bei.

Assume that the following conditions hold:

- (i) $\gcd(u, q - 1) = 1$.
- (ii) $q \equiv 1 \pmod{8}$.
- (iii) $3^{\frac{q-1}{4}} \equiv 1 \pmod{p}$.

Then $f(x)$ is a permutation polynomial of \mathbb{F}_q .

Proof. We will prove that under the above conditions, the polynomial f induces a one-to-one application on \mathbb{F}_q . Suppose that $f(a) = f(b)$ for some elements a and b of \mathbb{F}_q . If one of them, say a , is 0, then $b^u(b^{\frac{q-1}{2}} + b^{\frac{q-1}{4}} + 1) = 0$. Suppose that $b \neq 0$, then $b^{\frac{q-1}{2}} + b^{\frac{q-1}{4}} + 1 = 0$. Set $c = b^{\frac{q-1}{4}}$, then $c^2 + c + 1 = 0$ and c is a cubic root of unity. Condition (iii) implies $c \neq 1$. We have $c = c^4 = b^{q-1} = 1$, which is a contradiction. It follows that $b = 0 = a$.

From now on we may suppose that $ab \neq 0$. It is clear that $a^{\frac{q-1}{2}} = \pm 1$ and $b^{\frac{q-1}{2}} = \pm 1$. By symmetry, we have to consider only the following three cases:

Case 1: If $a^{\frac{q-1}{2}} = b^{\frac{q-1}{2}} = 1$. If $a^{\frac{q-1}{4}} = b^{\frac{q-1}{4}} = 1$, then $a^u = b^u$, hence $(\frac{a}{b})^u = 1$. Therefore $a = b$ by (i). To complete Case 1, we may suppose that $a^{\frac{q-1}{4}} = 1$ and $b^{\frac{q-1}{4}} = -1$. From equation (1) we have $3a^u = b^u$ or $(\frac{b}{a})^u = 3$. We deduce that $\left(\frac{b^{\frac{q-1}{4}}}{a^{\frac{q-1}{4}}}\right)^u = 3^{\frac{q-1}{4}}$, hence $(-1)^u = 1$ by (iii). By (i), u is odd and we reached a contradiction.

Case 2: If $a^{\frac{q-1}{2}} = b^{\frac{q-1}{2}} = -1$. From equation (1) we get: $a^{u+\frac{q-1}{4}} = b^{u+\frac{q-1}{4}}$, hence $(b/a)^{u+\frac{q-1}{4}} = 1$. The order δ of b/a in \mathbb{F}_q divides $q - 1$ and $u + \frac{q-1}{4}$. Let l be a prime factor of δ . Because u is odd and by (ii), we may exclude the case $l = 2$. It follows that l is odd and $l \mid \frac{q-1}{4}$, therefore $l \mid u$, which contradicts (i).

Case 3: If $a^{\frac{q-1}{2}} = -b^{\frac{q-1}{2}} = 1$. Here we have $a^{\frac{q-1}{4}} = \pm 1$ and $b^{\frac{q-1}{4}} = \zeta$, where ζ is a primitive quartic root of unity.

- If $a^{\frac{q-1}{4}} = -1$ and $b^{\frac{q-1}{4}} = \zeta$, then by equation (1), we have $a^u = \zeta b^u$. We deduce that $(a/b)^u = \zeta$, therefore $(a/b)^{4u} = 1$. Using (i), we conclude that $(a/b)^4 = 1$. If $a/b = -1$, then $a^{\frac{q-1}{2}} = (-1)^{\frac{q-1}{2}} b^{\frac{q-1}{2}} = b^{\frac{q-1}{2}}$, which is a contradiction. Suppose next that $a/b = \pm\zeta$, then $a^{\frac{q-1}{2}} = (\pm\zeta)^{\frac{q-1}{2}} b^{\frac{q-1}{2}}$. Hence $1 = (\zeta^4)^{\frac{q-1}{8}}(-1) = -1$, which is a contradiction. We conclude that $a = b$.
- Suppose now that $a^{\frac{q-1}{4}} = 1$ and $b^{\frac{q-1}{4}} = \zeta$, then by equation (1) we have $3a^u = \zeta b^u$. By condition (iii), the characteristic of the field is $\neq 3$, hence we may write this equation in the form: $(a/b)^u = \zeta/3$. It follows that $\left(\frac{a^{\frac{q-1}{2}}}{b^{\frac{q-1}{2}}}\right)^u = \frac{\zeta^{\frac{q-1}{2}}}{3^{\frac{q-1}{2}}}$, hence $3^{\frac{q-1}{2}} = -1$, contradicting (iii). \square

Remark 1 The minimal example for Theorem 1 is when $p = 7$ and $q = 7^2$.

Example 1 Let p be a prime number such that $p \equiv 1 \pmod{8}$ and $p \equiv 1 \pmod{3}$ and let $q = p^r$ where r is positive and even. It is clear that condition (ii) of Theorem 1 is satisfied. Euler criteria gives that $3^{\frac{p-1}{2}} = \left(\frac{3}{p}\right) = 1$ (see [6]). It follows that $3^{\frac{p-1}{4}} = \pm 1$, hence $3^{\frac{q-1}{4}} = (3^{\frac{p-1}{4}})^{(1+p+\dots+p^{r-1})} = 1$ and condition (iii) of Theorem 1 is fulfilled. By Dirichlet's theorem (see [6]), there exist infinitely many prime numbers $p \equiv 1 \pmod{8}$ and $p \equiv 1 \pmod{3}$. The smallest such prime is 73. Any polynomial $f(x)$ of the form (1) such that u satisfies condition (i) of Theorem 1 induces a permutation of \mathbb{F}_q . We may put $u = 1$ for example.

Acknowledgement

The second author was supported in part by NSERC.

References

- [1] Dickson, L.E.: The analytic representation of substitutions on a power of a prime number of letters with a discussion of the linear group. *Ann. of Math.* 11 (1896/97) 16, 161–183.
- [2] Hermite, C.: Sur les fonctions de sept lettres. *C.R. Acad. Sci. Paris* 57 (1863) 750–757.
- [3] Lidl, R.; Mullen, G.L.: When does a polynomial over a finite field permute the elements of the field? *Amer. Math. Monthly* 95 (1988) 243–246.
- [4] Lidl, R.; Niederreiter, H.: *Finite fields (Encyclopedia of Mathematics and its Applications)*. Cambridge Univ. Press, 2008.
- [5] Small, C.: *Arithmetic of finite fields*. Marcel Dekker, Inc., 1991.
- [6] Strayer, J.K.: *Elementary Number Theory*. PWS Publishing Company, Boston 1994.
- [7] Wan, D.Q.; Lidl, R.: Permutation polynomials of the form $x^r f(x^{(q-1)/d})$ and their group structure. *Monatsh. Math.* 112 (1991) 2, 149–163.

Mohamed Ayad
 Laboratoire de Mathématiques Pures et Appliquées
 Université du Littoral
 F-62228 Calais, France
 e-mail: ayad@lmpa.univ-littoral.fr

Omar Kihel
 Department of Mathematics
 Brock University
 Ontario, Canada L2S 3A1
 e-mail: okihel@brocku.ca