

# Triangles with sides in arithmetic progression

Autor(en): **Mikami, Kentaro / O'Hara, Jun / Sugawara, Kunio**

Objekttyp: **Article**

Zeitschrift: **Elemente der Mathematik**

Band (Jahr): **72 (2017)**

Heft 2

PDF erstellt am: **25.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-730834>

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, [www.library.ethz.ch](http://www.library.ethz.ch)

<http://www.e-periodica.ch>

## Triangles with sides in arithmetic progression

Kentaro Mikami, Jun O’Hara and Kunio Sugawara

Kentaro Mikami is a differential geometer and former professor of the Graduate School of Engineering and Resource Science, Akita University, Japan. He graduated from Tohoku University in 1970, where he also earned his Ph.D.

Jun O’Hara is a geometer and professor of the Department of Mathematics and Informatics, Faculty of Science, Chiba University, Japan. He was born in 1963 and graduated from the University of Tokyo where he obtained his Ph.D. in 1991.

Kunio Sugawara is a differential geometer and former professor of the Department of Mathematical Sciences, Osaka Kyoiku University, Japan. He was born in 1960 and graduated from Kyoto University, where he earned his Ph.D.

In this note we study geometric properties of triangles  $\triangle ABC$  the sides of which are in arithmetic progression,  $CA - BC = AB - CA$ . As far as the authors know, this topic does not seem to have been in the spotlight<sup>1</sup>. This is why we would like to integrate some known results and (possibly) new results here.

**Theorem 1.** *Let  $\triangle ABC$  be a triangle with edge lengths  $a = BC, b = CA, c = AB$ . Then, each of the following three conditions is a necessary and sufficient condition so that the sequence of sides  $a, b, c$  is an arithmetic progression.*

(i) *The line joining the centroid  $G$  and the incenter  $I$  is parallel to the edge  $AC$ .*

<sup>1</sup>with the exception of the study in the case of Heronian triangles, i.e., triangles such that the side lengths and area are all integers ([BG]).

Im Zentrum dieses Artikels stehen Dreiecke, deren Seiten  $a, b, c$  eine arithmetische Progression bilden, d.h.  $b = (a + c)/2$ . Derartige Dreiecke sind weder besonders selten noch besonders häufig anzutreffen: In der Menge der Äquivalenzklassen ähnlicher Dreiecke bilden sie einen Raum der Kodimension eins, genau wie gleichschenklige oder rechtwinklige Dreiecke. Im Gegensatz zu den eben genannten Klassen scheinen aber Dreiecke, deren Seiten eine arithmetische Progression bilden, wenig untersucht worden zu sein, obwohl sie, wie die Autoren der vorliegenden Arbeit zeigen, hübsche geometrische Eigenschaften aufweisen, über die sie sich charakterisieren lassen.




Fig. 1




Fig. 2

- (ii) *The line joining the incenter  $I$  and the circumcenter  $O$  is perpendicular to the line  $BI$  (see Figure 1).*
- (iii) *The midpoint of  $B$  and the excenter opposite to  $B$  lies on the edge  $AC$  (see Figure 2).*

The condition (i) is given in an exercise in Section 8.6.7 on page 120 of a note by Paul Yiu which is available through the web ([Y]). The authors found the conditions (ii) and (iii) accidentally when they were drawing figures with softwares such as GeoGebra, Maple, and Mathematica. Later, they got informed that the condition (ii) has been given in Problem 1 of the Indian National Mathematical Olympiad 2006. Three proofs of (ii), using Stewart's theorem, Ptolemy's theorem, and Euler's theorem and the formula of  $\cos(B/2)$  respectively, can be found in [I]. We give a (possibly) new proof of (ii), a little bit more elementary, but still using Euler's theorem. As for the condition (iii), the authors do not know references.

Let us first prepare formulae which are needed in our proof. Let  $r, R$  denote the radii of the incircle and the circumcircle respectively,  $S$  the area of  $\triangle ABC$ , and  $s = (a+b+c)/2$ . Then, Heron's formula states

$$S = \sqrt{s(s-a)(s-b)(s-c)}. \quad (1)$$

Since  $|\triangle ABC| = |\triangle IAB| + |\triangle IBC| + |\triangle IAC|$  we have

$$r = \frac{2S}{a+b+c} = \frac{S}{s}. \quad (2)$$

The law of sines,  $R = a/(\sin A)$ , implies

$$R = \frac{abc}{4S}. \quad (3)$$

Finally, Euler's theorem (see, for example, [J, p. 186]) states

$$IO^2 = R^2 - 2rR. \quad (4)$$

*Proof.* (i) The line  $GI$  is parallel to the line  $AC$  if and only if the heights of  $G$  and  $I$  above the line  $AC$  are the same, which should be one third of the height of the vertex  $B$ . Therefore the condition (i) is equivalent to saying that  $|\triangle IAC| = |\triangle ABC|/3$ , which, by (2), is equivalent to  $br = (a + b + c)r/3$ .

(ii) Let  $D$  be the foot of the perpendicular to the edge  $BC$  from  $I$  (see Figure 1). Since  $D$  is the tangent point of the incircle with the side  $BC$ , we have  $BD = (a - b + c)/2 = s - b$ . Therefore,

$$\begin{aligned} OI \perp BI &\iff BI^2 + IO^2 = OB^2 \\ &\stackrel{(4)}{\iff} (BD^2 + r^2) + (R^2 - 2rR) = R^2 \\ &\stackrel{(2)}{\iff} 2rR = (s - b)^2 + \frac{s(s - a)(s - b)(s - c)}{s^2} \\ &\stackrel{(2),(3)}{\iff} \frac{abc}{2s} = \frac{s - b}{s} (s(s - b) + (s - a)(s - c)) \\ &\iff abc = 2(s - b)(2s^2 - (a + b + c)s + ac) = 2(s - b)ac \\ &\iff b = a - b + c, \end{aligned}$$

which completes the proof.

(iii) Let  $J$  be the excenter opposite to  $B$ ,  $M$  the intersection of  $BJ$  and  $AC$ . We show that  $2b = a + c$  if and only if  $BJ = 2BM$ .

Let us compute  $BJ$  first. Let  $H$  be the foot of perpendicular to the line  $BC$  from  $J$  (see Figure 1). Since  $H$  is the tangent point of the excircle and the line  $BC$ , we have  $BH = (a + b + c)/2 = s$ . On the other hand, since  $JH$  is the radius of the excircle,  $JH = 2S/(a - b + c) = S/(s - b)$ . Therefore,

$$\begin{aligned} BJ^2 &= BH^2 + JH^2 \\ &\stackrel{(1)}{=} s^2 + \frac{s(s - a)(s - b)(s - c)}{(s - b)^2} \\ &= \frac{s}{(s - b)} (s(s - b) + (s - a)(s - c)) \\ &= ac \frac{a + b + c}{a - b + c}. \end{aligned}$$

Next we compute  $BM$ . Let  $E$  be the intersection of the circumcircle of the triangle  $ABC$  and the line  $BM$ . Put  $m = CM, n = AM, x = BM$ , and  $y = ME$ . Since the triangle




Fig. 3

$BEC$  is similar to the triangle  $BAM$  (see Figure 3), we have  $(x + y) : a = c : x$ , i.e.,  $x^2 = ac - xy$ . On the other hand, the secant theorem implies  $xy = mn$ . Since  $m : n = a : c$ ,

$$x^2 = ac - mn = ac - \frac{ab}{a+c} \cdot \frac{cb}{a+c} = \frac{ac}{(a+c)^2} ((a+c)^2 - b^2),$$

which implies

$$BM^2 = \frac{ac(a+b+c)(a-b+c)}{(a+c)^2}. \quad (5)$$

Therefore,

$$\begin{aligned} BJ = 2BM &\iff (a+c)^2 = 4(a-b+c)^2 \\ &\iff 2b = a+c, \end{aligned}$$

which completes the proof.  $\square$

**Remark 2.** We can also show the formula (5) by applying the law of cosines to  $\triangle BAC$  and  $\triangle BAM$ .

### Acknowledgement

The authors thank the anonymous referee of the previous version of this note for the references [CG] and [I].

### References

- [BG] H. Bailey and W. Gosnell, *Heronian triangles with sides in arithmetic progression: An inradius perspective*, Mathematics Magazine 85 (2012), 290–294.

- [CG] D. Coxeter and S. Greitzer, *Geometry Revisited*, Mathematical Association of America, Washington, DC, USA, (1967).
- [I] <http://olympiads.hbcse.tifr.res.in/uploads/inmo-2006>
- [J] Roger A. Johnson, Advanced Euclidean Geometry, Dover Publ., (2007) [1929]
- [Y] P. Yiu, <http://math.fau.edu/Yiu/EuclideanGeometryNotes.pdf>

Kentaro Mikami  
Graduate School of Engineering and Resource Science  
Akita University  
1-1 Tegata Gakuen-machi  
Akita City, 010-8502 JAPAN  
e-mail: mikami@math.akita-u.ac.jp

Jun O'Hara  
Department of Mathematics and Informatics  
Faculty of Science  
Chiba University  
1-33 Yayoi-cho, Inage  
Chiba, 263-8522, JAPAN  
e-mail: ohara@math.s.chiba-u.ac.jp

Kunio Sugawara  
Department of Mathematical Sciences  
Osaka Kyoiku University  
4-698-1 Asahigaoka  
Kashiwara, Osaka 582-8582 JAPAN  
e-mail: sugahara@cc.osaka-kyoiku.ac.jp