Radiolarian faunal turnover through the Paleocene-Eocene transition, Mead Stream, New Zealand

Autor(en): Hollis, Christopher J.

Objekttyp: Article

Zeitschrift: Eclogae Geologicae Helvetiae

Band (Jahr): 99 (2006)

Heft [1]: Radiolaria : siliceous plankton through time : proceedings of the tenth meeting of the International Association of Radiolarian Palaeontologists INTERRAD X

PDF erstellt am: 20.05.2024

Persistenter Link: https://doi.org/10.5169/seals-169256

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek* ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

http://www.e-periodica.ch

Radiolarian faunal turnover through the Paleocene-Eocene transition, Mead Stream, New Zealand

CHRISTOPHER J. HOLLIS¹

Key words: South Pacific, Paleogene, Radiolaria, diatoms, paleoecology, global change

ABSTRACT

The Mead Stream section, northern Clarence Valley, is the most complete Paleocene-early Eocene record of pelagic sedimentation in the mid-latitude (~55° S paleolatitude) Pacific Ocean. Integrated studies of sediments, siliceous and calcareous microfossils and carbon isotopes have shown that major global climate events are recorded by distinct changes in lithofacies and biofacies. The consistent and often abundant occurrence of siliceous microfossils in the section provides a rare opportunity to undertake quantitative analysis of highlatitude radiolarian population changes through the late Paleocene and early Eocene. Late Paleocene assemblages are dominated by spumellarians, although the nassellarian species Buryella tetradica is the most abundant species. The Paleocene-Eocene boundary (= base of Paleocene-Eocene thermal maximum) in the Mead Stream section is marked by major faunal turnover, including an abrupt decrease in B. tetradica, first occurrences of several low-latitude species (e.g. Amphicraspedum prolixum s.s., Lychnocanium auxilla, Podocyrtis papalis, Phormocyrtis turgida, Theocorys? phyzella) and increased abundance of large, robust spumellarians relative to small actinommids. Above an 18-m thick, lowermost Eocene interval in which radiolarians are abundant to common, radiolarian abundance declines progressively, falling to <10 individuals per gram in the marl-dominated unit that is correlated with the early Eocene climatic optimum. These trends in siliceous microfossil populations signal major changes in watermass characteristics along the northeastern New Zealand margin in the earliest Eocene. Assemblages typical of cool, eutrophic, watermasses that dominated the Marlborough Paleocene were replaced in the early Eocene by assemblages more characteristic of oligotrophic, stratified, subtropical-tropical watermasses.

ZUSAMMENFASSUNG

Das Mead Stream Profil, nördliches Clarence Tal, ist die vollständigste Paläozän-Früh Eozän Aufzeichnung pelagischer Sedimentation in mittleren Breiten (~55° S Paläobreite) des Pazifischen Ozeans. Integrierte Studien von Sedimenten, kieseligen und kalkigen Mikrofossilien und Kohlenstoffisotopen haben gezeigt, dass bedeutende globale Klimaereignisse durch deutliche Veränderungen in Litho- und Biofazies gekennzeichnet sind. Das beständige und oft sehr häufige Vorkommen kieseliger Mikrofossilien im Profil bietet eine seltene Gelegenheit, quantitative Studien von Veränderungen innerhalb von Radiolarienpopulationen hoher Breiten während des späten Paläozäns und frühen Eozäns durchzuführen. Spätpaläozäne Vergesellschaftungen werden von Spumellarien dominiert, obwohl die Nassellarienart Buryella tetradica die häufigste Art ist. Die Paläozän-Eozän Grenze (=Basis des Paläozän-Eozän Wärmemaximums) im Mead Stream Profil ist durch bedeutende faunistische Erneuerungen gekennzeichnet, einschließlich der abrupten Abnahme von B. tetradica, dem ersten Auftreten zahlreicher Arten niederer Breiten (z. B. Amphicraspedum prolixum s.s., Lychnocanium auxilla, Podocyrtis papalis, Phormocyrtis turgida, Theocorys? phyzella) und der zunehmenden Häufigkeit von großen, robusten Spumellarien im Verhältnis zu kleinen Actinommida. Oberhalb eines 18 Meter mächtigen Intervals des untersten Eozäns, in dem Radiolarien sehr zahlreich bis häufig sind, nimmt die Radiolarienhäufigkeit progressive ab, bis auf eine Anzahl von weniger als zehn Individuen pro Gramm innerhalb der Mergel-dominierten Einheit, die dem früheozänen Klimaoptimum entspricht. Diese Trends innerhalb von Populationen kieseliger Mikrofossilien signalisieren bedeutende Veränderungen in Wassermassencharakteristika entlang des nordöstlichen neuseeländischen Kontinentalrandes im frühesten Eozän. Vergesellschaftungen typisch für kühle, eutrophische Wassermassen, die das Marlborough Paläozän dominierten, wurden im frühen Eozän durch Vergesellschaftungen ersetzt, die eher für oligotrophische, stratifizierte, subtropische-tropische Wassermassen charakteristisch sind.

1. Introduction

Radiolaria are the most diverse and widely distributed group of marine plankton preserved in the fossil record (Lipps 1993; De Wever et al. 2001). High evolutionary turnover and considerable geographic variation in populations make radiolarian fossils invaluable research tools in biostratigraphy and paleoceanography. Consequently, detailed census studies through sedimentary successions have the potential to reveal much about biosphere-geosphere interactions in the marine realm during critical episodes in Earth history. Whereas quantitative studies are common for late Cenozoic radiolarians (e.g. Boltovskoy 1987; Caulet et al. 1992), very few such studies have been made of early Cenozoic or Mesozoic assemblages.

Paleocene-Eocene radiolarians, Mead Stream S79

¹ Institute of Geological and Nuclear Sciences, PO Box 30-368, Lower Hutt, New Zealand. Email: c.hollis@gns.cri.nz



Fig. 1. Simplified geological map of eastern Marlborough (after Crampton et al. 2003) showing location of Mead Stream and other stratigraphic sections mentioned in the text.

Notable exceptions are studies of Cretaceous-Tertiary boundary radiolarians in New Zealand and Ecuador (Hollis 1996; Keller et al. 1997; Hollis et al. 2003a, b). In this article we report the results of the first quantitative study of radiolarian assemblage variation through the Paleocene-Eocene (P-E) transition (~60-50 Ma).

In addition, this is the first study of radiolarian faunal change through the P-E transition in the mid-latitude South Pacific; eastern Marlborough being situated at $\sim 55^{\circ}$ S in the latest Paleocene (King et al. 1999; Sutherland et al. 2001). Sanfilippo & Nigrini (1998a) reviewed DSDP and ODP records of late Paleocene-early Eocene radiolarians and established a composite biostratigraphy from well-preserved, low-latitude assemblages in 12 sites between 40°N and 30°S. Most of these sites are in the western central and North Atlantic, with two in the South Atlantic, one in the western central Indian Ocean and one in the North Pacific. Subsequently, Nigrini & Sanfilippo (2000) and Sanfilippo & Blome (2001) reported on radiolarian assemblages in well-constrained P-E boundary records in Caribbean and western North Atlantic ODP holes, respectively.

The section exposed in the southern branch of Mead Stream, Clarence River valley (Fig. 1), is the most complete known record of South Pacific pelagic-hemipelagic sedimentation from Late Cretaceous to middle Eocene (Strong et al. 1995; Hollis et al. 2005). Rich radiolarian assemblages make it a key reference section for early Paleogene radiolarian biostratigraphy and it is the type section for eight zones (RP6-13) in the South Pacific radiolarian zonation (Strong et al. 1995; Hollis 1997, 2002; Hollis et al. 2005). Detailed stratigraphic and paleoenvironmental studies have been completed on the Cretaceous/Tertiary (K/T) boundary (Hollis et al. 2003a) and the

S80 C. J. Hollis



Fig. 2. Lithostratigraphy, biostratigraphy and carbon isotope stratigraphy of Upper Cretaceous-lower Eocene strata at Mead Stream (after Hollis et al. 2005). For age and biostratigraphy columns, intervals of uncertain age or biozone are shaded. Three episodes of major global change are identified from bulk carbonate δ^{13} C stratigraphy: late Paleocene carbon isotope maximum (PCIM), Paleocene-Eocene thermal maximum (PETM) and early Eocene climatic optimum (EECO). For comparison, the global compilation of benthic foraminiferal δ^{13} C data of Zachos et al. (2001) is also shown; the plotted values are running five-point averages calibrated to the geochronometric time scale of Berggren et al. (1995) following Hollis et al. (2005).

Paleocene-Eocene (P-E) transition (Hollis et al. 2005). In the latter study, carbon isotope stratigraphy was integrated with radiolarian, foraminiferal and calcareous nannofossil biostratigraphy in order to relate lithofacies changes to major perturbations in the early Paleogene global carbon cycle and climate system. Three of these episodes of perturbations have been identified in upper Paleocene and lower Eocene strata at Mead Stream (Fig. 2): the late Paleocene carbon isotope maximum (PCIM – Corfield & Cartlidge 1992, Thompson & Schmitz 1997; Kurtz et al. 2003), the Paleocene-Eocene thermal maximum (PETM – Zachos et al. 2001, 2003), and the early Eocene climatic optimum (EECO – Zachos et al. 2001). The base of the PETM corresponds with the Paleocene-Eocene boundary (Zachos et al. 2003). The aim of the current study is to use changes in radiolarian assemblages through the P-E transition at Mead Stream to further elucidate the influence of these global events on oceanic conditions in the Southwest Pacific.

2. Material and Methods

This study is based on 42 radiolarian assemblages recovered from rock samples between 100 and 250 m above the well-constrained K/T boundary (Hollis et al. 2003a) in the southern and main branch of Mead Stream. Sampling methods are described by Hollis et al. (2005). For radiolarian microfossil extraction, samples were crushed into 5–10 mm chips and leached in 15% hydrochloric acid (HCl) until reaction ceased. Siliceous rocks with weak reaction to HCl digestion were then rinsed and leached in 5% hydrofluoric acid for 2–4 hours. Samples were then washed through a 63 μ m screen and the residues cleaned by gentle heating in a 1:1 solution of 10% hydrogen peroxide and calgon, (NaPO₃)₆.

Taxonomic nomenclature is summarised in Table 1. Selected species are illustrated in Plates 1-3. Census data (Hollis et al. 2005, supplementary data) have been derived from vertical traverses of strewn slides under transmitted light. For sparse samples (<300 radiolarians per slide), all radiolarians on one, two or more slides were counted. For richer samples, the following double count method was utilised. First, all specimens were counted until a total count of ~300 was achieved. The proportion of the slide examined to this point was determined and the abundance of common taxa (>15 radiolarians or >5% of the total fauna) was estimated for the remainder of the slide. The remainder of the slide was then examined and occurrences of rare taxa (<5% in initial count) recorded. This method has the advantage of establishing reliable abundance estimates for rare taxa without expending great amounts of time counting many hundreds of specimens. One disadvantage is that the total count is highly variable, ranging in this dataset from <60 to >7000 individuals. Abundance of larger ($>63 \mu m$) diatoms was also recorded during the initial count (Hollis et al. 2005).

In order to obtain an accurate estimate of relative abundance, all specimens have been assigned a taxonomic category. Although these counting groups range from the level of subspecies to family, they are given equal weighting in diversity analysis.

3. Stratigraphy

The lithostratigraphy and foraminiferal, radiolarian and dinoflagellate cysts biostratigraphy of Upper Cretaceous to middle Eocene strata at Mead Stream was described by Strong et al. (1995). Subsequently, the radiolarian biostratigraphy was reviewed and revised by Hollis (1997). Hollis et al. (2003a) described the stratigraphy of the K/T boundary transition in detail, as part of a paleoenvironmental study of this interval at Mead Stream and nearby Branch Stream (Fig. 1). The K/T boundary is defined as zero datum in the current study. As a forerunner to the present study, Hollis et al. (2005) completed a detailed stratigraphic study of upper Paleocene-lower Eocene strata at Mead Stream, incorporating lithostratigraphy, foraminiferal, calcareous nannofossil and radiolarian biostratigraphy and carbon isotope stratigraphy (Fig. 2).

Paleocene-Eocene strata exposed along Mead Stream consist of well-bedded units of chert, siliceous limestone with chert nodules, siliceous mudstone, limestone and marl, which form three formations within the Muzzle Group (Reay 1993). The Mead Hill Formation consists of dm-bedded chert and siliceous limestone. It extends from a faulted base 170 m below the K/T boundary to a sharp contact with overlying the Waipawa Formation at 108 m above the K/T boundary. The Waipawa Formation is a 7.5 m thick interval (108–115.5 m) that consists of lower and upper siliceous mudstone units with an intervening unit of siliceous limestone. The two mudstone units carry the same distinctive geochemical signature of the Waipawa Formation as in its type area in Hawkes Bay, North Island (Killops et al. 2000).

The overlying formation, Amuri Limestone, has been subdivided into several lithotypes and members (Reay 1993; Hollis et al. 2005). The basal lithotype is Lower Limestone, which includes two members, Dee Marl and Upper Ribbon Chert. Lower Limestone (116-192 m) consists of dm-bedded siliceous limestone with thin marl interbeds and rare chert nodules. One unusually thick marl bed provides a useful reference point (136.5-136.6 m) within Lower Limestone and is referred to as Mudstone C (following Hollis et al. 2005). Dee Marl is a 2.4 m thick unit (157-159.4 m) of dm-bedded alternating marl and marly limestone. This unit marks the base of the Eocene at Mead Stream and at the type section, Dee Stream (Hancock et al. 2003). Upper Ribbon Chert, the highest occurrence of chert in the section, is a 12.6 m thick interval (159.4-172 m) of dmbedded siliceous limestone with narrow chert stringers in the centre of the beds. The abundance of chert decreases gradually up-section, whereas the abundance and thickness of marl interbeds increases. The top of Lower Limestone is placed at the base of the first of a series of thick marl beds intercalated with limestone beds. The overlying lithotype, Lower Marl, consists of 110 m of alternating marl and marly limestone beds, similar to the lithology of Dee Marl. The stratigraphy of the upper part of Amuri Limestone, which is not considered here, is described by Strong et al. (1995).

4. Siliceous microfossil occurrence

Of 134 samples processed for radiolarians in upper Paleocene and lower Eocene strata at Mead Stream (~75–300 m), 42 samples contained radiolarian assemblages of adequate preservation and abundance for species level identification and quanti-



Fig. 3. Variation (a) radiolarian preservation. (b) radiolarian abundance, (c) diatom/radiolarian ratio, (d) radiolarian diversity (Fisher α Index), (e) relative abundance of major radiolarian groups, and (f) radiolarian first and last occurrences in upper Paleocene-lower Eocene siliceous microfossil assemblages at Mead Stream. Radiolarian preservation is recorded as Barren, Poor (internal and surface features obscured by recrystallisation). Moderate (internal features obscured by infilling or recrystallisation, surface features preserved), Good (test morphology not affected by recrystallisation, no infilling, breakage minor to moderate). Shaded intervals represent two distinctive units (WF = Waipawa Formation, Mst C = Mudstone C) and a 21 m-thick lowermost Eocene interval characterised by peak radiolarian abundances (rad acme).

tative analysis (Fig. 3). Radiolarians are generally common but poorly preserved in the upper Paleocene where strong recrystallisation, including siliceous infilling and overgrowth, is evident in siliceous limestone samples (Fig. 3a–b). In the first 21 m of Eocene strata, radiolarians are generally abundant and moderately preserved. Recrystallisation is less pronounced in this interval. This probably reflects reduced silica content in Dee Marl. In the overlying Upper Ribbon Chert, chertification is restricted to a central zone in each bed and samples taken from the outer parts of beds contained relatively wellpreserved siliceous microfossils. In the overlying marl-rich Eocene interval, radiolarians are increasingly rare but preservation is moderate to good.

Diatoms are common in radiolarian residues in the Paleocene but are rare to very rare in the Eocene (Fig. 3c). The original change in diatom abundance across the P/E boundary is likely to have been even more pronounced than is described here. Owing to their smaller average size and more delicate tests, diatoms are more prone to diagenetic destruction than radiolarians. Therefore, diatoms are likely to be under-represented in Paleocene siliceous limestone samples relative to the better preserved Eocene material.

Radiolarian diversity is highly variable. Most assemblages contain 20–50 taxa whereas three assemblages have <20 taxa and seven assemblages have >50 taxa. The Fisher α Index is used to assess diversity trends in census data (Fig. 3d). The diversity curve for all taxa is considered to be less reliable than the curve for nassellarians. Most nassellarians have been identified to species level, whereas many spumellarians remain undifferentiated below family level. This results in an underestimate of overall diversity in spumellarian-dominated samples, notably in the earliest Eocene. Nassellarian α values indicate relatively low diversity in the Paleocene, apart from a short-

S82 C. J. Hollis

lived increase in the marl bed (Mudstone C) near the base of Zone RP7. A rapid increase in diversity in the earliest Eocene is followed by a gradual decline with relatively low α values through the Lower Marl.

Paleocene radiolarian assemblages are dominated by spumellarians (Fig. 3e) and, in order of abundance, the most common species and species groups are: *Buryella tetradica*, *Haliomma* spp. gr. b, *Amphisphaera coronata* gr, *A. goruna*, *Lithelius minor* gr. and *Buryella granulata*. In the lowermost 21 m of Eocene strata, spumellarian dominance is greater than in the Paleocene and the most common species are: *Haliomma* gr. b., *Amphisphaera coronata* gr. *Lithelius minor* gr. and *Theocorys*? cf. *phyzella*. In overlying Eocene strata, spumellarians dominate to a similar extent as in the Paleocene and the most common species are: *Haliomma* gr. b, *Amphisphaera coronata* gr. *Lithelius minor* gr., *Calocycloma ampulla*, *Phormocyrtis striata striata*, *Podocyrtis papalis*, *Sethocytris babylonis*, *Axoprunum pierinae* and *Periphaena heliasteriscus*.

5. Radiolarian faunal turnover through the Paleocene-Eocene transition

In contrast to low-latitude regions where there is little change in radiolarian assemblages across the Paleocene-Eocene boundary (Sanfilippo & Nigrini 1998a; Nigrini & Sanfilippo 2000; Sanfilippo & Blome 2001), four episodes of significant faunal turnover are observed in the upper Paleocene-lower Eocene succession at Mead Stream (Fig. 3f, Table 2).

The first episode occurs in the lower Waipawa Formation and is associated with the RP5/RP6 zone boundary. The first occurrences (FOs) of 22 species and the last occurrences (LOs) of 12 species are recorded within 3 m of strata (~108–111 m). Some of the LOs are isolated records of Cretaceous-early PaTable 1. Annotated list of radiolarian species or species groups encountered in this study. Species ranges are based on the South Pacific zonation for Mead Stream and other South Pacific ("Sth Pac") sections (Hollis, 1997, 2002; Hollis et al. 1997), and on the tropical zonation (Sanfilippo and Nigrini, 1998a) for low latitude ("Low Lat") records; LK = Late Cretaceous, UZP = unzoned Paleocene, EOC = Eocene, MIO = Miocene. Reference to taxon concept: CC42 = Clark and Campbell, 1942; F73 = Foreman, 1973; H02 = Hollis, 2002; H97a = Hollis, 1997; H97b = Hollis et al. 1997; K99 = Kozlova, 1999; N92 = Nishimura, 1992; S85 = Sanfilippo et al. 1985; SR92 = Sanfilippo and Riedel, 1992; S95 = Strong et al. 1995; SR73 = Sanfilippo & Riedel, 1973.

Taxon name	Zonal range			Taxon co	ncept
	Mead	Sth Pac	Low Lat.	Ref.	Strong et al. 1995:
Amphicraspedum murrayanum HAECKEL	RP8	RP8	RP6-8	SR73	
Amphicraspedum prolixum form. gracilis (LIPMAN)	RP8-9	RP8-9	?	K99	
Amphicraspedum prolixum SANFILIPPO & RIEDEL gr.	RP7-10	RP7-10	RP7-13	SR73	
Amphicraspedum prolixum	RP8-10	RP8-10	RP7-9	RP7-9	208
Amphinternis alamedaensis (CAMPB & CL)	RP3-9	RK9-RP9	LK-RP7	H97a	
Amphipynday stocki (CAMPB & CLARK)	RK9-RP7	RK9-RP7	LK-UZP	H97a	
Amphisphaera aff. magnaporulosa CL. & CAMPB.	RP5-9	RP4-6	?	H02	
Amphisphaera coronata s.l. (EHRENBERG)	RP4-9	RP4-6	UZP-EOC	H97a	208 (A. radiosa)
Amphisphaera goruna (SANFILIPPO & RIEDEL)	RP3-8	RP2-6	UZP-RP7	H97a	208, fig. 8G, 9A
Amphisphaera kina Hollis	RP2-7	RP2-6	?	H97a	208, fig. 8F
Amphisphaera macrosphaera (NISHIMURA)	RP3-9	RP3-14	UZP-RP6	H97a	208 (A. radiosa)
Amphymenium splendiarmatum CLARK & CAMPB.	RP6-8	RP2-6	RP8-20	H97a	
Artostrobus pusillus (EHRENBERG)	RK9-RP8	RK9-RP8	?	H97a	
Aspis murus NISHIMURA	RP6	RP5-6	RP6	N92	
Axoprunum pierinae (CLARK & CAMPB.)	RP5-9	RP6-14	UZP-EOC	SR73	208, fig. 10C
Axoprunum? aff. bispiculum (POPOFSKY)	RP6-9	RP4-6	?	H02	
Bathropyramis magnifica (CLARK & CAMPB.)	RP5-13	RP4-13	?	H02	
Bathropyramis sanjoaquinesis (CAMPB. & CLARK)	RK9-RP8	RK9-RP8	LK-UZP	H97a	
Bekoma bidartensis RIEDEL & SANFILIPPO	RP8	RP8	RP7-8	S 95	208, fig. 9G
Bekoma campechensis FOREMAN	RP6	RP6	RP6-7	S95	208, fig. 9F
Bekoma divaricata FOREMAN	RP6-8	RP6-8	RP7	S 95	208, fig. 9H
Buryella dumitricai PETRUSHEVSKAYA	RP4-7	RP4-7	?	H97a	208, fig. 8L-M
Buryella foremanae PETRUSHEVSKAYA	RP4-7	RP4-7	?	H97a	208, fig. 8K
Buryella granulata (PETRUSHEVSKAYA)	RP3-8	RP4-8	?	H97a	208, fig. 8J
Buryella helenae O'CONNOR	RP6	RP4-6	?	H02	
Burvella kaikoura HOLLIS	RP5-6	RP5-6	?	H97a	
Buryella pentadica FOREMAN	RP6	RP6	UZP-RP7	N92	
Burvella petrushevskavae O'CONNOR	RP6	RP5-6	?	H02	
Burvella tetradica tetradica FOREMAN	RP5-9	RP5-8	UZP-RP9	H02	208, fig. 8N, 9O-R
Burvella tetradica tridica O'CONNOR	RP6	RP5-6	?	H02	200, 18. 01. , 7. 2
Calocycloma ampulla (EHRENBERG)	RP5-10	RP5-10	RP6-14	S95	208. fig. 9V-X
Cassideus aff. mariae NISHIMURA	RP5-9	RP5-9	?	H02	209. fig. 9C (Microsciadiocapsa ? sp.)
Clathrocyclas australis HOLLIS	RP3-6	RP2-6	?	H97a	P. 208, 8P (Clathrocycloma sp. A)
Clathrocyclas universa gr.	RP5-13	RP5-13	?	H97b	·····
Conoactinomma stilloformis (LIPMAN)	RP9	RP9	?	K99	206 (Conocaryomma sp.)
Cornutella californica CAMPB. & CLARK	RK9-RP13	RK9-RP13	LK-UZP	H97a	(
Corvthomelissa adunca (SANFILIPPO & RIEDEL)	RP5-8	RP4-8	RP6-8	H02	208, fig. 9B
Corythomelissa spp.	RP6-8	RP6-8	?	H02	
Cryptocarpium? cf. ornatum (EHRENBERG)	RP6-13	RK9-RP13	?	H97a	
Cycladophora aff. cosma LOMBARI AND & LAZARUS	RP6-8	RP4-8	?	H02	
Cycladophora cf. bicornis (POPOFSKY)	RP6-8	RP6-8	?	H02	208, fig. 9D (Clathrocycl. humerus)
Dictvomitra andersoni (CAMPB. & CLARK)	RK9-RP6	RK9-RP6	LK-UZP	H97a	,,
Haliomma gr. B	RP4-9		H97a		
Lamptonium fabaeforme fabaeforme (KRASHEN.)	RP8-9	RP8-9	RP7-9	S85	
Lamptonium pennatum FOREMAN	RP7-9	RP7-9	UZP-RP8	S 95	208, fig. 9I
Lamptonium? aff. colymbus FOREMAN	RP9		F73		
Lithelius ? minor gr. JÖRGENSEN	RK9-RP9	RK9-RP14	?	H97a	
Lithocampe wharanui Hollis	RP3-6	RK9-RP6	?	H97a	
Lithomespilus coronatus SOUINABOL	RK9RP7	RK9-RP8	UZP-RP8	H97a	
Lithostrobus longus GRIGORJEVA	RP1-7	RP1-6	?	H02	209, fig. 8H-I (Stichomitra wero)
Lophophaena mugaica (GRIGORJEVA)	RP5-8	RP2-8	?	H02	

Paleocene-Eocene radiolarians, Mead Stream $\ S83$

Table 1. (cont.)

Taxon name	Zonal range			rept									
	Mead	Sth Pac	Low Lat.	Ref.	Strong et al. 1995:								
Lychnocanium aff. carinatum EHRENBERG	RP8	RP8	RP6-7	N92	209, fig. 9M (Rhopalocanium? sp.)								
Lychnocanium aff. tripodium EHRENBERG	RP7-8	RP7-8	?	K99									
Lychnocanium amphitrite (FOREMAN)	RP9-13	RP9-14	RP8-20	F73									
Lychnocanium auxilla (FOREMAN)	RP8	RP8	RP6-7	F73									
Lychnocanium bellum CLARK & CAMPB.	RP9-13	RP9-14	RP8-14	F73	209, fig. 11I-J (as Lychnocanoma)								
Lychnocanium satelles (KOZLOVA)	RP5-9	RP5-9	?	H02	208, fig. 9L (L. auxilla)								
Lychnocanium sp. B	RP7-8		?	S95	209, fig. 9J								
Lychnocanium zhamoidai KOZLOVA	RP7-9		?	K99	209, fig. 9K (Lychnocanoma sp. A)								
Middourium regulare	RP6-8	RP6-8	RP6-7	K99									
Mita regina (CAMPB. & CLARK)	RK9-RP6	RK9-RP6	LK-UZP	H97a									
Mita sp. A (? = Siphocampe? "eilizabethae")	RP9-10	RP9-RP15	?	S95	209, fig. 10X								
Monobrachium irregulare	RP7-8	RP7-8	RP6-7	K99									
Myllocercion acineton FOREMAN	RK9-RP6	RK9-RP4	LK-UZP	H97a									
Orbiculiforma renillaeformis gr. (CAMPB. & CL.)	RK9-RP6	RK9-RP6	LK-UZP	H02									
Patagospyris confluens (EHRENBERG)	RP4-RP8	RP4-8	UZP-RP10	K99									
Periphaena alveolata (LIPMAN)	RP5-9	RP5-9	?	K99									
Periphaena heliasteriscus (CLARK & CAMPB.)	RP8-9	RP6-9	UZP-EOC	SR73									
Peritiviator? dumitricai NISHIMURA	RP5-7	RP5-6	UZP-RP6	N92									
Petalospyris fovealata EHRENBERG	RP5-9	RP5-9	?	K99	209, fig. 8Q-R								
Petalospyris senta (KOZLOVA)	RP6-9	RP6-9	?	K99	209, fig. 8S								
Phacostaurus? quadratus NISHIMURA	RP6-7	RP6	RP6-7	N92									
Phormocyrtis cubensis (RIEDEL & SANFILIPPO)	RP8	RP8	RP6-8	F73									
Phormocyrtis striata exquisita (KOZLOVA)	RP7-8	RP7-8	RP6-9	S 95	209, fig. 9N								
Phormocyrtis striata striata (BRANDT)	RP7-10	RP7-10	RP7-14	S95	209, fig. 9O-P								
Phormocyrtus turgida (KRASHENINNIKOV)	RP8	RP8	RP6-8	F73									
Plectodiscus circularis (CLARK & CAMPB.)	RP6-8	RP6-8	RP6-EOC	PK72									
Podocyrtis acalles SANFILIPPO & RIEDEL	RP8-9	RP8-9	RP8-9	S92	209, fig. 9T-U (P. aphorma)								
Podocyrtis papalis EHRENBERG	RP8-10	RP8-14	RP7-18	S92	209, fig. 9S								
Pterocodon poculum NISHIMURA	RP6-7	RP6-7	RP6-7	H02									
Saturnalis kennetti DUMITRICA	RP3-8	RP1-8	?	H97a									
Sethochytris babylonis (CLARK & CAMPB.) gr.	RP6-13	RP6-15	RP6-20	H02	208, fig. 11A-B (as Lychnocanoma)								
Siphocampe nodosaria (HAECKEL)	RP6-13	RP4-15	RP6-14	H02	209 (S. arachnaea)								
Siphocampe quadrata (PETRUSHEV. & KOZLOVA)	RP8-13	RP5-15	RP6-20	H02	209, fig. 10R								
Spongurus bilobatus gr. CLARK & CAMPB.	RP5-9	RP4-9	?	H02									
Stichomitra carnegiense CAMPB. & CLARK	RK9-RP6	RK9-RP6	LK-UZP	H97a									
Stylodictya targaeformis CLARK & CAMPB.	RP5-9	RP5-9	?	CC42									
Stylosphaera minor CLARK & CAMPB.	RP3-9	RP3-6	UZP-MIO	H97a									
Theocampe urceolus (CAMPB. & CLARK)	RP9-13	RP9-15	RP8-20	S95	209, fig. 10Q								
Theocampe vanderhoofi (CAMPB. & CLARK)	RK9-RP6	RK9-RP6	LK-UZP	H97a									
Theocorys? aff. phyzella FOREMAN	RP6-8	RP6-8	?	H02	208, fig. 9E (Calocyclas asperum)								
Theocorys? cf. phyzella FOREMAN	RP7-8	?	RP7	SB01	(as Theocorys aff. phyzella)								
Theocorys? phyzella FOREMAN	RP8		RP7-8	F73									
Tholodiscus densus (KOZLOVA)	RP6-8	RK9-RP8	?	H97a									

S84 C. J. Hollis



Fig. 4. Ranges of radiolarian index species in upper Paleocene-lower Eocene strata at Mead Stream. For age and biostratigraphy columns, intervals of uncertain age or biozone are shaded. Low-latitude ranges are based on (Sanfilippo & Nigrini 1998a, b) and Sanfilippo & Blome (2001).

leocene species (e.g. Lithocampe wharanui, Mita regina, Myllocercion acineton) that may be reworked into the siliciclastic Waipawa Formation. Other events are isolated records of taxa that are largely restricted to two particularly rich and diverse assemblages (P30/f318, f452), e.g. Buryella helenae, B. petrushevskayae, Cryptocarpium cf. ornatum. As most of the FOs recorded in this interval relate to species that are known to have earlier local FOs (Hollis 1997, 2002), this episode is not thought to reflect a time of high evolutionary turnover. Instead, the large number of events at the base of the Waipawa Formation suggests that either the unit is highly condensed or that there is a basal unconformity.

The second episode occurs within or directly below the 10 cm thick marl bed (Mudstone C). The FOs of 6 species and the LOs of 5 species occur within 5 m of strata. Most of the

LOs are the last record of Cretaceous-Paleocene species that are generally very rare in the upper Paleocene but are present in one or more of three relatively rich samples in this interval (P20/f1072, f1076, f1081); two of these species may be reworked into Mudstone C (*Amphipyndax stocki, Theocampe* vanderhoofi). Three species with FOs in this interval are key index species that have much earlier FOs in low latitudes (*Lamptonium pennatum, Phormocyrtis striata exquisita, P. s.* striata – Fig. 4). All are very rare in overlying Paleocene strata but occur consistently in the lowermost Eocene.

The third faunal turnover episode occurs at the base of the Dee Marl (Fig. 3f, Table 2). The FOs of 13 species are recorded within the first 3 m of Eocene strata at Mead Stream, with 9 of these species in the basal Eocene sample. Four of these species appear to be global markers for the P/E boundary

Table 2. Radiolarian species in order of (A) first and (B) last occurrence at Mead Stream. Species with first occurrences lower in the section (Hollis et al. 2003a) or last occurrences higher in the section (Strong et al. 1995) are not shown. (MHF = Mead Hill Formation, WF = Waipawa Formation, DM = Dee Marl, URC = Upper Ribbon Chert)

A. IN ORDER OF FIRST OCCURREN	CE																																			
International Age		Late Paleocene											Early Eocene																							
South Pacific Radiolarian Zone	4	5			RF	° 6									RP7	7										RP	8						F	₹ Р9		
Lithologic Unit	N	ИH	F		WF							Lo	wer	Lir	nest	tone	•					D	М	ι	JRC	2	I	lowe	r Ls	t		I	low	er M	arl	
	_	6	7	8	3	5	100		72	76	81	94	66	00	0 10	10	05	15	1	22	23 25	29	32	28	3 6	46	47	51	4	5	6	9	5	~ ~	0 0	20
NZ Fossil Record No. P30/f	45	52	52	31	50	45	10	35	10	10	10	10	2 :	Ξ 3	501	1	: =	Ξ	53	= :	= =	Ξ	Ξ	53	53	Ξ	=	12	53	53	49	53	53	46	5 6	64
	0	1.7	22	3.3	.3		4. 4	0	9.6	6.9	7.2	9.6	6.1	2.7	0.4			<u>.</u>	0.4	× 1.00	0.0	3.3	0.6	9.6	25	7.1	8.0	5.5	6.0	0.0	4.5	0.0	0.0	5.0	2.0	2.5
DEPTH (metres above KTB)	99.	10	10	108	110	Ξ:	121	1	136	136	13,	136	4	4	4 4	140	150	15	154	15	15:1	158	15	15	16	16	16	11	17	18	19.	22	23	24	20	29
Axoprunum pierinae	*	*	*			*	* *	• *	*	*	*	*		* :	* *	* *	•	*	*	*			*	* *	* *	*	*	* *	*	*	*	*	*	*	* *	* *
Axoprunum? aff. bispiculum	*		*	*	*	*	* *	*	*	*	*	*	*	* :	* *	* *	¢	*	*	*		*	*	* *	* *	*	*	* *	*		*					
Bathropyramis magnifica	*	*	*	*	*	*	* *	•	*	*	*	*		*	* *	* *	* *	*	*	*	*		*	* *	* *	*	*	*	*	*	*			*	*	* *
Buryella tetradica	*	*	*	*	*	*	* *	• *	*	*	*	*	*	*	* *	* *	* *	*	*	*	* *	*	*	* *	* *	*	*	* *	*	*	*	*	*	2	*	
Cassideus aff. mariae	*	*	*		*	*	* *	•		*	*				*	* *	•		*		*	*	*	* *	* *	*	*	* *								
Conocaryomma stilloformis	*																																			*
Corythomelissa adunca	*	*	*			*				*	*				3	k			*				*	*	*			*	*	*						
Heliostylus spp.	*					*										*	¢												*							*
Peritiviator? dumitricai	*					*					*																									
Periphaena alveolata	*	*	*	*		*	*	*	*	*	*	*	*	*	* *	k		*	*	*		*	*	* *	* *	*	*	* *	*	*		*			*	*
Stylodictya targaeformis	*	*				*																											*	*		
Calocycloma ampulla		*	*	*		*	* *	4	*	*	*											*	*	,	* *		*	*		*	*		*	*	* *	* *
Cycladophora aff. cosma		*				*				*													*													
Lophophaena mugaica		*	*			*				*																*										
Patagospyris confluens		*	*	*					*	*		*		*								*	*	,	ĸ	*	*									
Petalospyris fovealata		*	*	*		*				*	*	*			*	* *	•					*	*	*	*			*	*		*					
Amphisph. aff. magnaporulosa			*	*	*	*	* *	4	*	*	*	*	*	*	* *	* *	¢	*	*	*	* *		*	* >	* *	*	*	*		*		*				
Amphymenium splendiarmatum			*	*		*			*	*											*			,	ĸ		*	*	*	*						
Aspis murus			*																																	
Bekoma campechensis			*			*																														
Burella pentadica			*	*																																
Buryella tridica			*	*																																
Cycladophora cf. bicornis			*						*	*					2	ĸ									*			*	*							
Petalospyris senta			*			*				*													*	* >	* *			* *	*	*	*					
Siphocampe nodosaria			*																				*	,	k		*	*	*							
Amphicraspedum prolixum gr.				*																	*	*	*	,	* *		*	* *	*	*	*		*	*		
Buryella helenae				*				1																												
Buryella petrushevskayae				*																																
Cryptocarpium cf. ornatum				*		*		L																									*	*	*	ĸ

Table 2 (cont.)

International Age	Late Paleocene									Early Eocene																									
South Pacific Radiolarian Zone	5	5 RP6 RP7								RP8 RP9																									
Lithologic Unit	Mł	ΗF	W	/F	Lower Limestone								D	М	U	JRC	2		Lo	wer	Lst			Lo	wer	Ma	rl								
NZ Fossil Record No. P30/f	451	527	318	452	1057	625	1072	1076	1081	1094	1100	530	1011	1110	1105	1115 531	100	1123	1125	1129	1132	532	533	1146	1147	1151	1223	534	535	499	527	498	538	539	497
DEPTH (metres above KTB)	0.66	108.2	108.3	110.8	116.4	0.151	136.6	136.9	137.2	139.6	141.9	144.0	145.4	149.1	150.3	151.8	0.401	155.6	156.6	158.3	159.0	159.6	166.5	167.1	168.8	171.5	175.5	176.0	180.0	194.5	0.022	245.0	250.0	280.0	292.5
Bekoma divaricata Lychnocanium zhamoidai				*	* * :	*	*	*	*											*	*		*				*	*	*	*			*	*	
Middourium regulare Phacostaurus? quadratus				*	3	ĸ	*	*	*	*	* *	* *	*			* *	*			*	*	* :	* *	*	*	*		*							
Plectodiscus circularis				*	*		*	*	*	*													*	2											
Sethochytris babylonis gr.				*	*	*	*	*	*	3	* *	k				*	* :	*	*	*	*	* :	* *		*	*	*	*	*	*	* >	* *	*	*	*
Theocorys? all. phyzella				*	* :	* *	*	*										*																	
Theocorys : CL phyzena Theolodisous densus				*	*	1*	*	*	*	Ŧ							-	*		*	*	* *	* *	*	*	*	*	*	*						
Lychnocanium sp. B				Ŧ			*	*						*		* *		4			*		* .					*							
Phormocurtis striata exavisita						*	*	*	*	*			*	Ŧ		* 1		*			*	÷.,			*	*	*	*							
Lychnocanium aff. tripodium						1	*	*	*											*	*	*		T	Ŧ	т	Ŧ	т							
Phormocyrtis striata striata							*	*	*								3	*	*	*	*	*	*	*	*	*	*	*	*	*			*	*	*
Pterocodon poculum							*	*	*																									÷	
Lamptonium pennatum								*	*												*	* :	* *	*		*		*	*	*					
Monobrachium irregulare																	ş	* *		*	*	* :	* *	*	*	*	*	*	*		*				
Amphicraspedum murrayanum																				*	*														
Amphicraspedum prolixum s.s.																				*	*	* :	* *		*		*	*	*						*
A. prolixum f. gracilis																				*	*		*					*	*	*					*
Lamptonium f. fabaeformae																				*	*		*			*		*	*	*				*	
Lychnocanium auxilla																				*	*	* *	* *	*		*	*	*	*						
Phormocyrtis cubensis																				*	*	,	*	*			*								
Phormocyrtis turgida																				*	*				*			*							
Theocorys? phyzella																				*	*	* *	* *	*	*	*	*	*	*						
Podocyrtis papilis																				*	*	* *	* *	*		*	*	*	*	*		*	*	*	*
Clathrocyclas universa gr.																					*	* *	* *	*	*	*		*	*			k			
Lychnocanium aff. carinatum																					*		*					*	- 1						
Podocyrtis acelles																					*	*	*								,	ĸ	*		*
Bekoma bidartensis																						*	*												
Siphocampe quadrata																							*			*	*								
Periphaena heliasteriscus																											*		*	*	*	*	*	*	*
Mita sp. A																												*	*						*
Lamptonium? aff. columbus																														*	* *	4			
Theocampe urceolus																															2	¢	*		*
Lychnocanium bellum																																¢			*
Lychnocanium amphitrite		1																											- 1				*	*	
First Occurrences	10	5 9	4	09	0	0 2	2 3	1	0	0	0 (0 0	0	0	0	0 (0	1 (0	9	3	1	0 1	0	0	0	1	1	0	1	0	2 0	1	0	0

Table 2 (cont.)

Table 2 (cont.) B. IN ORDER OF LAST OCCURRENCE	CE				Ľ	
International Age			Late Paleocene		Early Eocene	
South Pacific Radiolarian Zone	5	RP6		RP7	RP8	RP9
Lithologic Unit	MHF	WF	Lower L	limestone	DM URC Lower Lst Low	ver Marl
NZ Fossil Record No. P30/f	451 526 527	503 503 452 1057 528	529 529 1072 1076 1081 1094 1099 1100	530 11101 11105 11105 11105 531 1122 1123	1129 1132 532 533 1138 533 1147 1151 1151 1151 1153 534 535 535 537 537 537 537	498 538 539 497
DEPTH (metres above KTB)	99.0 107.7 * 108.2	108.3 110.3 110.8 116.4 131.5	136.6 136.6 136.6 136.9 137.2 137.2 139.6 141.9 141.9	144.0 145.4 149.1 150.3 151.8 151.8 154.8 154.8 155.6 155.6	158.3 159.0 159.6 161.2 161.2 166.5 167.1 167.1 168.8 171.5 175.5 175.5 175.5 176.0 176.0 176.0 176.0 176.0 177.5 220.0 220.0 0 220.0 0 220.0 0 220.0 0 220.0 0 220.0 0 220.0 0 20.0 159.6 16 16 16 16 16 16 16 16 16 16 16 16 16	245.0 250.0 280.0 292.5
Liphocampe wharanui Mita regina Buryella helenae Buryella kaikoura Burgella pastedica	* * * * *	* *				
Buryella pentatica Buryella petrushevskayae Buryella tetradica tridica Myllocercion acineton Bekoma campechensis	*	* * *				
Clathrocyclas australis Dictyomitra andersoni Stichomitra carnegiense Amphipyndax stocki Theosomuca yandarhoofi	* * *	* * * * * * * * * *	*	?		
Phacostaurus? quadratus Peritiviator? dumitricai Pterocodon poculum	*	* *	* * *			
Buryella dumitricai	* * *	* * * * *	* * *			
Amphisphaera kina	* * *	* * * * *	* * * * * * *	* *		
Lithomespillus coronatus Theocorys? aff. phyzella	*	* * *	* * * * * * * * *	* * * * * *		
Lithostrobus longus Amphicraspedum murrayanum Cycladophora aff. cosma Lychnocanium aff. tripodium	* * *	* * *	* *	* * * *	* * * *	
Bekoma bidartensis Lychnocanium sp. B Plectodiscus circularis Lophophaena mugaica	* *	* *	* * * * * * *	* * * *	*	
Patagospyris confluens Cassideus aff. mariae Orbiculiforma renillaeformis gr. Phormocyrtis cubensis Siphocampe quadrata	* * *	* * * *	** * *	** * *	* * * * * * * * * * * * * * * * * * * *	

Table 2 (cont.)

Table 2 (cont.)																																					
International Age	_								La	ate l	Pale	eoce	ne															Ea	ırly	Eoce	ene						
South Pacific Radiolarian Zone	5				RP6	8								F	RP7											R	P8							R	P9		
Lithologic Unit	N	1HF	Τ	V	VF	Г					- 1	Lov	ver l	Lim	nest	one						D	М		UR	С	Т	Lo	wer	Lst			L	owe	r M:	arl	_
						5			3	9	_ ,	4 0	~ 0	>	-	. 0	2	5	1	0.0	ς γ	6	5		×	4	7 0		3								_
NZ Fossil Record No. P30/f	451	526	170	318	452	105	528	529	107	107	108	601	110	530		2 =	10	Ξ	531	112	112	12	113	532	13	114	7 1	115	22	534	535	661	536	337	138	539	L6t
		-	~	m	n 00	. 4	5	0	9	•	~	0 0					~	~	~	~ `	0.10	-	_	10		0 -				41	41	4	<u>.</u>	<u></u>		47	4
DEPTH (matrix above KTP)	9.0	5	08	08	10.1	9	31	32.	36.	36.	22	39.6		44	12	6	50.	21.8	4.0	4.4	20.00	8	59.0	69.6	10	90.		11.00	15.	76.0	80.0	4	0.0	20.0 2	0.05	<u>80.0</u>	2.5
Artostrobus pusillus	6 *	-	*	= .		-	*	-	*	*	*	*	*					-	1			-	-	Ξ.	Ξ.	= =		1	-	-	12	-	20	n'n	1 0	5	29
Burvella gravulata	*	*	*	*	* *	*		*	*	*	*	*						-	4	т Т	- -		4		<u>ـ</u>	<u>ب</u>		т ь т	ىد	т _							
Cycladophora of hicornis			*						*	*	-		-		*	Ť		Ŧ	Ŧ	Ť	Ť	1	Ť	Ť	Ť	÷.		• •	÷	*		1					
Lychnocanium aff carinatum															-								*			*			Ŧ	÷							
Middourium regulare					*		*		*	*	*	* >	* *	* *				*	*				*	4	*	τ.				Ť		1					
Phormocyrtis striata avauisita								*	*	*	*	*	e					Ţ	Ŧ	4		L.	÷	÷	т т	÷.		 	4	Ť							
Phormocyrtis turaida															-			т		Ŧ		L.	т 	Ŧ	Ŧ	T '			Ť	Ť		1					
Saturnalis kannatti	*				*					*	*											1	Ŧ				1			Ť		1					
Sinhocampe nodosaria			*																				4	Ŧ	÷			6	4	÷		i i					
Tholodiscus densus					*					*													т т		÷		1		Ť	÷							
Amphisphaera goruna	*	*	*	*	* *	*	*	*	*	*	*	* >	k *	. *	. *				*	*	*		*	*	*	* .		* *	*	*	*						
Amphymenium splendiarmatum			*	*	*				*	*															*				Ŧ	-	Ţ	i i					
Bekoma divaricata					*	*																*	*			*				*	*	i i					
Corvthomelissa adunca	*	*	*		*					*	*				*				*			1	*	*		*			*	*	*	1					
Lychnocanium auxilla																						*	*	*	*	* >	k	*	*	*	*						
Theocorys? cf. phyzella					*	*		*	*	*	*	*								*		*	*	*	*	* >	k >	* *	*	*	*						
Theocorys? phyzella																						*	*	*	*	* >	k x	k *	*	*	*	1					
Amphipternis alamadaensis	*	*			*																	1								*	*	*					
Axoprunum? aff. bispiculum	*		*	*	* *	*	*	*	*	*	*	* >	* *	* *	* *	*		*	*	*		*	*	*	*	* *	k	k *	*	*		*					
Lamptonium pennatum										*	*											1	*	*	*	* *	k	*		*	*	*					
Petalospyris fovealata		*	*	*	*					*	*	*			*	*						*	*	*		*		*		*		*					
Petalospyris senta			*		*					*													*	*	*	*		*	*	*	*	*					
Amphisph. aff. magnaporulosa			*	*	* *	*	*		*	*	*	* >	k *	*	*	*		*	*	*	* *	8	*	*	*	* *	k x	k	*		*		*				
Monobrachium irregulare																				*	*	*	*	*	*	* *	k	* *	*	*	*	í –	*				
Clathrocyclas universa gr.																							*	*	*	* *	k x	* *		*	*	í –		*			
Lamptonium? aff. columbus																																*	*	*			
Lychnocanium satelles	*	*	*	*	*	*	*	*	*	*	*	* *	* *	* *	¢				*				*		*	*			*	*	*	*		*			
Stylodictya targaeformis	*	*			*																													* *	*		
Buryella tetradica	*	*	*	*	* *	*	*	*	*	*	*	* *	* *	*	* *	*	*	*	*	*	* *	*	*	*	*	* *	k x	k *	*	*	*	*	*	*	*		
Cryptocarpium cf. ornatum				*	*																											1		* *	*	*	
Lamptonium f. fabaeformae																						*	*			*		*		*	*	*				*	
Lychnocanium amphitrite																																			*	*	
Lychnocanium zhamoidai					*	*	*		*	*	*											1				*			*	*	*	*			*	*	
Periphaena alveolata	*	*	*	*	*	*		*	*	*	*	* *	* *	*	*			*	*	*		*	*	*	*	* *	k x	* *	*	*	*		*			*	
Podocyrtis acelles																						1	*	*		*						1		*	*		*
Last Occurrences	10	0	3	6	0 3	1	0	0	2	1	2	0	0 0	0 0	0 0) ()	0	1	0	2	0 1	0	2	1	0	3	1	1 0	4	10	7	5	2	3	1 1	5	1



Fig. 5. Relative abundance of common radiolarian species and species groups in upper Paleocene-lower Eocene strata at Mead Stream.

(Sanfilippo & Nigrini 1998a; Sanfilippo & Blome 2001): Lamptonium fabaeforme fabaeforme, Podocyrtis papalis, Phormocyrtis turgida and Theocorys? phyzella (Fig. 4). Five of the species are index species with late Paleocene FOs in low latitudes (Sanfilippo & Nigrini 1998a, b): Amphicraspedum murrayanum, A prolixum s.s., Bekoma bidartensis, Lychnocanium auxilla, Phormocyrtis cubensis (Fig. 4).

There are relatively few significant LOs associated with the P/E boundary (Fig. 3f). The LOs of *Lithomespilus coronatus*, *Lithostrobus longus* and *Theocorys*? aff. *phyzella* occur within the uppermost 3 m of Paleocene strata. Of the three LOs that occur within the basal Eocene Dee Marl, two are species that are restricted to the unit (*Amphicraspedum murrayanum, Lychnocanium* aff. *tripodium*) while one is the LO of a sporadically occurring late Paleocene species (*Cycladophora* aff. *cosma*). In contrast to the faunal changes observed at the base



Fig. 6. Paleogeographic reconstruction of New Zealand region during the latest Paleocene (adapted from King et al. 1999) showing major current systems based on coupled climate model simulations for the Eocene (Huber 2002)

of the Waipawa Formation, the low number of LOs relative to FOs in the P/E boundary interval indicates that the faunal turnover is not due to a stratigraphic discontinuity. The abrupt appearance of many low-latitude species indicates that warming at the PETM promoted pole-ward migration of warmwater radiolarians. Similar, but less extreme, short-lived warming may explain the FOs of several low-latitude species around the time of deposition of Mudstone C (Fig. 3f, Table 2).

The final episode of marked faunal change in the studied section occurs ~20 m above the P/E boundary (Fig. 3f, Table 2). Within the same 5 m interval in which radiolarian abundance begins to decrease (175–180 m), the LOs of 21 species are recorded. These species include many that thrived through much of the late Paleocene as well as many that appeared in the earliest Eocene (Fig. 4). Only two FOs are recorded in this interval, one of these being the FO of the common Eocene phacodiscid *Periphaena heliasteriscus*. The high number of LOs, coupled with the onset of long term decline in radiolarian abundance and diversity, indicates that this episode of faunal change reflects a oceanographic changes unfavourable to all radiolarians.

6. Radiolarian biofacies

To simplify the discussion and presentation of census data (Fig. 5) in this section, some species are combined into larger groups that reflect a close phylogenetic relationship or morphological similarity:

I. Amphisphaera coronata gr. (*A. macrosphaera*, *A. coronata* s.l, *A.* aff. *magnaporulosa*).

- 2. Amphisphaera goruna gr. (A. kina, A. goruna)
- 3. Axoprunum spp. (Axoprunum pierinae, A. aff. bispiculum)
- 4. Buryella granulata gr. (B. granulata, B. foremanae, B. dumitricai)
- Middourium/Monobrachium (Middourium regulare, Monobrachium irregulare)
- 6. Periphaena spp. (Periphaena alveolata, P. heliasteriscus)
- 7. Phormocyrtis spp. (Phormocyrtis cubensis, P. striata exquisita, P. striata striata, P. turgida)
- 8. Plectopyramidins (Bathropyramis and Cornutella spp.)
- 9. Theocorys? phyzella gr. (Theocorys? cf. phyzella, Theocorys? phyzella)

Three radiolarian biofacies are identified in the Paleocene-Eocene section at Mead Stream: (A) upper Paleocene (100–157 m), (B) lowermost Eocene (157–178 m) and (C) lower Eocene (178–250 m).

Biofacies A is characterised by abundant *Buryella tetradica*, common to abundant *Amphisphaera coronata* gr. and *Haliomma* gr. b, and few to common *Amphisphaera goruna* gr., *Buryella granulata* gr. and *Cassideus* aff. *mariae* (Fig. 5). Diatoms are common in this biofacies (Fig. 3c) and sediments are moderately siliceous, with a highly siliceous facies of the organic-rich Waipawa Formation (Killops et al. 2000) occurring in the lower part of the interval. Faunal similarities with diatom-rich, biosiliceous Paleocene lithofacies lower in the Marlborough sequence (Hollis et al. 1995, 2003a, b) and on the eastern margin of Campbell Plateau (Hollis 2002) suggest moderately eutrophic conditions, consistent with weak to moderate upwelling in outer shelf to upper slope depths (Fig. 6).

This biofacies spans an interval correlated with the PCIM, which is interpreted as a 3 m.y. episode of enhanced global carbon burial, either through increased marine or terrestrial productivity (Corfield & Cartlidge 1992; Thompson & Schmitz 1997; Kurtz et al. 2003). The regional occurrence of organicrich mudstone in the lower part of the interval (Killops et al. 2000; Hollis et al. 2005) and abundant siliceous microfossils (Hollis 2002) indicates that this event is associated with enhanced marine productivity in the New Zealand region.

Biofacies B occurs in the lowermost 21 m of Eocene strata (157-178 m). It is characterised by abundant Haliomma gr. b and few to common Phormocyrtis spp. and Theocorys? phyzella gr. (Fig. 5). The base of the biofacies coincides with the P/E boundary and the onset of the PETM. A marked increase in radiolarian abundance and diversity is associated with a sharp decline in diatom abundance and in the abundance of Buryella tetradica - the dominant radiolarian species in the upper Paleocene (Fig. 3b-d, f). This species and B. granulata s.s. are rare in this interval, apart from two samples in which both species are common (P30/f533, f1223). The high abundance of large, robust spumellarians, namely Haliomma gr. b, Middourium/Monobrachium gr. and Periphaena alveolata, is a distinctive feature of Biofacies B. These taxa tend to be most abundant in marl samples of Dee Marl and are also common to abundant in underlying Paleocene marly layers, notably at ~137 and ~142 m, and overlying Eocene marl layers (Fig. 5). Despite the significant lithologic change from Dee Marl to Upper Ribbon Chert, there is little change in siliceous microfossil assemblages. A slight increase in biosiliceous productivity in Upper Ribbon Chert may be inferred from the increase in silica content and a small increase in diatom abundance, although overall radiolarian abundance and diversity declines (Fig. 3b–d).

Biofacies B has unusual characteristics that make interpretation difficult. The presence of many low-latitude species indicates relatively warm climatic conditions. The low abundance of diatoms and of radiolarian taxa characteristic of the eutrophic Paleocene Biofacies A, suggests conditions of relatively low productivity. However, radiolarian abundance and diversity maxima are recorded in Dee Marl and in underlying Mudstone C, which also contains an assemblage similar to Biofacies B. A very similar assemblage is present in lowermost Eocene carbonate ooze at high-latitude DSDP Site 277, southwest Campbell Plateau (Hollis et al. 1997). The three features of high total abundance, high diversity and high numbers of warm-water species are consistent with late Cenozoic tropical radiolarian assemblages (Casey 1993). This suggests that the PETM may have promoted the expansion of weakly eutrophic subtropical water into the Southwest Pacific (Fig. 6). Further biogeographic study of the distinctive spumellarians that dominate this biofacies is required to confirm this interpretation.

Biofacies C occurs in the upper part of the lower Eocene sequence (178 to 250 m). It is characterised by abundant Amphisphaera coronata gr., Haliomma gr. b, common Calocycloma ampulla, few to common Phormocyrtis striata striata and few to rare Amphicraspedum prolixum gr., Axoprunum spp., Buryella tetradica, Periphaena heliasteriscus, plectopyramidins, Podocyrtis papalis and Sethochytris babylonis gr. An upwards decrease in radiolarian abundance and diversity, coupled with low diatom abundance, indicates that this biofacies represents relatively oligotrophic conditions. Similar assemblages are found in lower Eocene carbonate ooze at DSDP site 277 (Hollis et al. 1997). As this biofacies is correlated with the EECO, it seems likely that global warming associated with this event promoted the expansion of oligotrophic subtropical watermasses into the Southwest Pacific.

7. Conclusions

The first quantitative study of radiolarian assemblages from an upper Paleocene-lower Eocene bathyal succession has identified three distinct biofacies that can be used to determine the influence of major changes in the global carbon cycle and climate system on Southwest Pacific oceanic conditions.

Biofacies A is associated with the PCIM, a 3 m.y. episode of maximum δ^{13} C values that is inferred to represent high rates of carbon burial due to enhanced marine or terrestrial productivity (Corfield & Cartlidge 1992; Thompson & Schmitz 1997; Kurtz et al. 2003). Hollis et al. (2005) argued that Mead Stream lithofacies, including organic mudstone and siliceous limestone,

are indicative of enhanced marine productivity in the Marlborough basin during the PCIM. Siliceous microfossil assemblages also indicate enhanced marine productivity. Biofacies A is associated with common diatoms and is dominated by the same or similar species that are found in biogenic silica-rich lithologies in the lower Paleocene in Marlborough (Hollis et al. 2003a, b) and in the upper Paleocene of the Campbell Plateau margin (Hollis 2002). Deposition of radiolarian and diatom-rich sediments in the Marlborough basin, as well as on the northwest and southeast continental margin of New Zealand, in the late Paleocene between ~60–56 Ma indicates that the PCIM was associated with enhanced marine productivity, at least in the southern Pacific Ocean. This is consistent with ocean circulation models for the South Pacific (Huber 2002).

Biofacies B is associated with the PETM, a 100–200,000 yr episode of extreme global warmth recorded as pronounced negative δ^{13} C and δ^{18} O excursions (Zachos et al. 2001), which at Mead Stream is associated with a lithofacies change from siliceous limestone to marly limestone and marl (Dee Marl). A marl bed (Mudstone C) ~20 m below the PETM may represent a precursor event. Not only is the bed also associated with a negative δ^{13} C excursion, but radiolarian assemblages are very similar to those of Biofacies B. Biofacies B is characterised by high radiolarian abundance and diversity, but very low diatom abundance. It is dominated by large, robust smooth-walled spumellarians of uncertain biogeographic affinity, and contains the only local record of several low-latitude species. The biofacies records an abrupt warm-water incursion in the East Coast Basin. The combination of an incursion of subtropical water and waning cool-water upwelling may have promoted an unusual bloom of warm water species. The occurrence of similar earliest Eocene radiolarian assemblages at New Zealand's southern continental margin (Hollis et al. 1997) indicates that warm subtropical waters may have extended as far south as 60°.

Biofacies C is associated with the onset of the EECO, a 3 m.y. episode of extreme global warming (Zachos et al. 2001), which at Mead Stream is associated with the transition from limestone- to marl-dominated sedimentation. Biofacies C is characterised by declining radiolarian abundance and diversity and low diatom abundance, implying a marked decrease in biosiliceous productivity. Significantly, many low-latitude as well as high-latitude radiolarian species disappear at the onset of the EECO. With similar early Eocene radiolarian assemblages recorded from the southern Campbell Plateau (Hollis et al. 1997), progressive warming is inferred to have promoted increasingly oligotrophic oceanic conditions around New Zealand.

Acknowledgements

Sample collection and stratigraphic integration was carried out with the assistance of Jerry Dickens, Brad Field, Craig Jones and Percy Strong. Samples were processed and slides were prepared by Kellee Anderson, Neville Orr and John Simes. The manuscript benefited for careful reviews by Annika Sanfilippo, Akiko Nishimura, James Crampton, Erica Crouch and Percy Strong. This research was supported by the Foundation for Research Science and Technology (GNS Global Change Through Time Programme, Contract CO5X0202).

REFERENCES

- BERGGREN W., KENT D., SWISHER C. & AUBRY M.-P. 1995: A revised Cenozoic geochronology and chronostratigraphy. In: BERGGREN, W. et al. (Eds): Geochronology, time scales and global stratigraphic correlation, 129–212, SEPM Special Publication 54.
- BOLTOVSKY D. 1987: Sedimentary record of radiolarian biogeography in the equatorial to Antarctic western Pacific Ocean. Micropaleontology 33, 267–281.
- CASEY R. 1993: Radiolaria. In: LIPPS, J.H. (Ed.): Fossil Prokaryotes and Protists, 249–284. Blackwell Scientific Publications, Oxford/London, UK.
- CAULET J., VENEC-PEYRE M.-T., VERGNAUD-GRAZZINI C. & NIGRINI C. 1992: Variation of South Somalian upwelling during the last 160 ka: radiolarian and foraminifera records in core MD 85674. In: SUMMERHAYES, C. et al. (Eds): Upwelling Systems: Evolution since the Early Miocene, Geological Society of London, Special Publication 64, 379–389.
- CORFIELD R. M. & CARTLIDGE J. E. 1992: Oceanographic and climatic implications of the Palaeocene carbon isotope maximum. Terra Nova 4, 443–455.
- CRAMPTON J., LAIRD M., NICOL A., TOWNSEND D. & VAN DISSEN R. 2003: Palinspastic reconstructions of southeastern Marlborough, New Zealand, for Late Cretaceous to Eocene times. New Zealand Journal of Geology and Geophysics 46, 153–175.
- DE WEVER P., DUMITRICA P., CAULET J.-P., NIGRINI C. & CARIDROIT M. 2001: Radiolarians in the sedimentary record. 533 p., Gordon and Breach Science Publishers, The Netherlands.
- HANCOCK H., DICKENS G., STRONG C., HOLLIS C. & FIELD B. 2003: Foraminiferal and carbon isotope stratigraphy through the Paleocene-Eocene transition at Dee Stream, Marlborough, New Zealand. New Zealand Journal of Geology and Geophysics 46, 1–19.
- HOLLIS C. 1996: Radiolarian faunal change through the Cretaceous-Tertiary transition of eastern Marlborough, New Zealand. In: MACLEOD, N. & KELLER, G. (Eds.): Cretaceous-Tertiary Mass Extinctions: Biotic and Environmental Changes, 173–204, Norton Press, New York.
- 1997: Cretaceous-Paleocene Radiolaria from eastern Marlborough, New Zealand. Institute of Geological and Nuclear-Sciences Monograph 17, 152 p.
- 2002: Biostratigraphy and paleoceanographic significance of Paleocene radiolarians from offshore eastern New Zealand. Marine Micropaleontology 46, 265–316.
- HOLLIS C., DICKENS G., FIELD B., JONES C. & STRONG C. 2005: The Paleocene-Eocene transition at Mead Stream, New Zealand: a southern Pacific record of early Cenozoic global change. Palaeogeography, Palaeoclimatology, Paleoecology 215, 313–343.
- HOLLIS C., RODGERS K. & PARKER R. 1995: Siliceous plankton bloom in the earliest Tertiary of Marlborough, New Zealand. Geology (Boulder) 23, 835–838.
- HOLLIS C., WAGHORN D., STRONG C. & CROUCH E. 1997: Integrated Paleogene biostratigraphy of DSDP site 277 (Leg 29): foraminifera, calcareous nannofossils, Radiolaria, and palynomorphs. Institute of Geological & Nuclear Sciences Science Report 97/7, 1–73.
- HOLLIS C., RODGERS K., STRONG C., FIELD B. & ROGERS K. 2003a: Paleoenvironmental changes across the Cretaceous/Tertiary boundary in the northern Clarence Valley, southeastern Marlborough, New Zealand. New Zealand Journal of Geology and Geophysics 46, 209–234.
- HOLLIS C., STRONG C., RODGERS K. & ROGERS K. 2003b: Paleoenvironmental changes across the Cretaceous/Tertiary boundary at Flaxbourne River and Woodside Creek, eastern Marlborough, New Zealand. New Zealand Journal of Geology and Geophysics 46, 177–197.
- HOLLIS C., DICKENS G., FIELD B., JONES C. & STRONG C. 2005: The Paleocene-Eocene transition at Mead Stream, New Zealand: a southern Pacific record of early Cenozoic global change. Palaeogeography, Palaeoclimatology, Paleoecology 215, 313–343.
- HUBER M. 2002: Straw man 1; preliminary view of the tropical Pacific from a global coupled climate model simulation of the early Paleogene. Proceedings of the Ocean Drilling Program, initial reports, Paleogene equatorial transect, Leg 199. Proceedings of the Ocean Drilling Program, Part A: Initial Reports 199, 30.

- KELLER G., ADATTE T., HOLLIS C., ORDONEZ M., ZAMBRANO I., JIMENEZ N., STINNESBECK E., ALEMAN A. & HALE-ERLICH W. 1997: The Cretaceous/Tertiary boundary event in Ecuador: reduced biotic effects due to eastern boundary current setting. Marine Micropaleontology 31, 97–133.
- KILLOPS S., HOLLIS C., MORGANS H., SUTHERLAND R., FIELD B. & LECKIE D. 2000: Paleoceanographic significance of Late Paleocene dysaerobia at the shelf/slope break around New Zealand. Palaeogeography, Palaeoclimatology, Palaeoecology 156, 51–70.
- KING P., NAISH T., BROWNE G., FIELD B. & EDBROOKE S. 1999: Cretaceous to Recent sedimentary patterns in New Zealand. Institute of Geological and Nuclear Sciences Folio Series 1. 35 p.
- KURTZ A., KUMP L., ARTHUR M., ZACHOS J. & PAYTAN A. 2003: Early Cenozoic decoupling of the global carbon and sulfur cycles. Paleoceanography 18(4), 1090, doi:10.1029/2003PA000908
- LIPPS J. H. 1993: Fossil prokaryotes and protists. 342 p., Blackwell Scientific, Boston.
- NIGRINI C. & SANFILIPPO A. 2000: Paleogene radiolarians from Sites 998, 999, and 1001 in the Caribbean. Proceedings of the Ocean Drilling Program, Scientific Results 165, 57–81.
- REAY M. 1993: Geology of the middle part of the Clarence Valley. Institute of Geological and Nuclear Sciences Geological Map 10. 144 p. + 1 map.
- SANFILIPPO A. & BLOME C. 2001: Biostratigraphic implications of mid-latitude Palaeocene-Eocene radiolarian faunas from Hole 1051A, ODP Leg 171B, Blake Nose, western North Atlantic. In: KROON, D. et al. (Eds.): Western North Atlantic Palaeogene and Cretaceous palaeoceanography. Geological Society Special Publication 183, 185–224.
- SANFILIPPO A. & NIGRINI C. 1998a: Upper Paleocene-Lower Eocene deep-sea radiolarian stratigraphy and the Paleocene/Eocene Series boundary. In: AUBRY, M.-P., et al. (Eds.): Late Paleocene-Early Eocene climatic and biotic events in the marine and terrestrial records, 244–276, Columbia University Press, New York.

- 1998b: Code numbers for Cenozoic low latitude radiolarian biostratigraphic zones and GPTS conversion tables. Marine Micropaleontology 33, 109–156.
- STRONG C., HOLLIS C. & WILSON G. 1995: Foraminiferal, radiolarian, and dinoflagellate biostratigraphy of Late Cretaceous to middle Eocene pelagic sediments (Muzzle Group), Mead Stream, Marlborough, New Zealand. New Zealand Journal of Geology and Geophysics 38, 171–209.
- SUTHERLAND R., KING P. & WOOD R. 2001: Tectonic evolution of Cretaceous rift basins in south-eastern Australia and New Zealand: implications for exploration risk assessment. In: HILL K. & BERNECKER T. (Eds.): Eastern Australasian basins symposium: a refocused energy perspective for the future. Petroleum Exploration Society of Australia Special Publication 1, 3–13.
- THOMPSON E. & SCHMITZ B. 1997: Barium and the late Paleocene δ^{13} C maximum: evidence of increased marine surface productivity. Paleoceanography 12, 239–254.
- ZACHOS J., PAGANI M., SLOAN L., THOMAS E. & BILLUPS K. 2001: Trends, rhythms, and aberrations in global climate 65 Ma to present. Science 292, 686–693.
- ZACHOS J., WARA M., BOHATY S., DELANEY M., PETRIZZO M., BRILLA., BRALOWER T. & PREMOLI-SILVA I. 2003: A transient rise in tropical sea surface temperature during the Paleocene-Eocene thermal maximum. Science 302, 1551–1554.

Manuscript received January 2004 Revision accepted February 2005

Paleocene-Eocene radiolarians, Mead Stream S93

Plate 1

Scanning electron micrographs of spumellarian radiolarians from lower Eocene strata, Mead Stream. Scale bars = $100 \,\mu$ m.

1. Amphisphaera coronata Ehrenberg gr. CH12076, P30/f534, 176 m, RP8.

2. Axoprunum aff. bispiculum (POPOFSKY). CH12086, P30/f534, 176 m, RP8.

3. Axoprunum pierinae (CLARK AND CAMPBELL). CH12079, P30/f534, 176 m, RP8.

4. Stylosphaera minor CLARK AND CAMPBELL. CH12078, P30/f534, 176 m, RP8.

5. Haliomma gr. b. (cf. Theocosphaerella rotunda (BORISSENKO)). CH12005, P30/f1129, 158.3 m, RP8.

6. Haliomma gr. b. CH12001, P30/f1129, 158.3 m, RP8.

7. Periphaena? duplus (KOZLOVA). CH12020, P30/f1129, 158.3 m, RP8.

8. Periphaena? duplus (KOZLOVA). CH12019, P30/f1129, 158.3 m, RP8.

9. Periphaena? duplus (KOZLOVA). CH12016, P30/f1129, 158.3 m, RP8.

10. Periphaena? duplus (KOZLOVA). CH12018, P30/f1129, 158.3 m, RP8.

11. Middourium regulare (BORISSENKO). CH12301, P30/f1129, 158.3 m, RP8.

12. Monobrachium irregulare (NISHIMURA). CH12012, P30/f1129, 158.3 m, RP8.

13. Monobrachium irregulare (NISHIMURA). CH12011, P30/f1129, 158.3 m, RP8.

14. Amphiscraspedum prolixum SANFILIPPO AND RIEDEL CH12094, P30/f534, 176 m, RP8.

15. Amphicraspedum gracilis (LIPMAN). CH12093, P30/f534, 176 m, RP8.

16. Amphicraspedum gracilis (LIPMAN). CH12092, P30/f534, 176 m, RP8.

17. Amphicraspedum gracilis (LIPMAN). CH12091, P30/f534, 176 m, RP8.

18. Amphicraspedum murrayanum HAECKEL. CH12030, P30/f1129, 158.3 m, RP8.

19. Amphicraspedum murrayanum HAECKEL. CH12064, P30/f1129, 158.3 m, RP8.

20. Amphiscraspedum prolixum SANFILIPPO AND RIEDEL CH12059, P30/f1129, 158.3 m, RP8.

21. Amphiscraspedum prolixum SANFILIPPO AND RIEDEL CH12089, P30/f534, 176 m, RP8.

22. Spumellarian gen. et sp. indet. CH12085, P30/f534, 176 m, RP8.

S94 C. J. Hollis



 $Paleocene-Eocene\ radiolarians,\ Mead\ Stream\ \ S95$

Plate 2

Scanning electron micrographs of nassellarian radiolarians from upper Paleocene (Fig. 4) and lower Eocene strata, Mead Stream. Scale bar = $100 \ \mu m$.

1. Patagopsyris confluens (EHRENBERG). CH12074, P30/f1129, 158.3 m, RP8.

2. Petalospyris foveolata (EHRENBERG). CH12075, P30/f1129, 158.3 m, RP8.

3. Clathrocyclas universa CLARK AND CAMPBELL. CH12103, P30/f534, 176 m, RP8.

4. Buryella tetradica FOREMAN. CH01028, P30/f318, 108.3, RP6.

5. Buryella tetradica FOREMAN. CH12040, P30/f1129, 158.3 m, RP8.

6. Buryella tetradica Foreman. CH12065, P30/f1129, 158.3 m, RP8.

7. Bathropyramis magnifica (CLARK AND CAMPBELL). CH12101, P30/f534, 176 m, RP8.

8. Bathropyramis magnifica (CLARK AND CAMPBELL). CH12100, P30/f534, 176 m, RP8.

9. Lychnocanium ponderosa (KOZLOVA). CH12067, P30/f1129, 158.3 m, RP8.

10. Lychnocanium satelles (KOZLOVA). CH12098, P30/f534, 176 m, RP8.

11. Phormocyrtis striata exquisita (KOZLOVA). CH12070, P30/f1129, 158.3 m, RP8.

12. Phormocyrtis striata striata BRANDT. CH12038, P30/f1129, 158.3 m, RP8.

Phormocyrtis turgida (KRASHENINNIKOV). CH12023, P30/f1129, 158.3 m, RP8.
Phormocyrtis turgida (KRASHENINNIKOV). CH12056, P30/f1129, 158.3 m, RP8.

15. Phormocyrtis cubensis (RIEDEL AND SANFILIPPO). CH12057, P30/f1129, 158.3 m, RP8.

16. Phormocyrtis cubensis (RIEDEL AND SANTLERTO). CH1203, 13011129, 150.5 III, RT8.

17. Phormocyrus cubensis (Riedel and Santilino). CH12029, P30/f1129, 158.3 m, RP8.

Inomocynis Cabelas (Repel and Sannerro): CH12020, 150/1129, 150/5 m, RF8.
Lamptonium fabaeforme fabaeforme (KRASHENINNIKOV). CH12055, P30/f1129, 158.3 m, RP8.

Lamptonium fabaeforme fabaeforme (KRASHENINIKOV). CH12048, P30/f1129, 158.3 m, RP8.

20. *Theocorys? phyzella* FOREMAN. CH12073, P30/f1129, 158.3 m, RP8.

21. *Theocorys? phyzella* FOREMAN. CH12042, P30/f1129, 158.3 m, RP8.

22. Theocorys? phyzella Foreman. CH12058, P30/f1129, 158.3 m, RP8.

23. Theocorys? phyzella FOREMAN. CH12045, P30/f1129, 158.3 m, RP8.

24. Theocorys? phyzella FOREMAN. CH12043, P30/f1129, 158.3 m, RP8.

S96 C. J. Hollis



 ${\it Paleocene-Eocene\ radiolarians,\ Mead\ Stream\ }S97$

Plate 3

Transmitted light micrographs of radiolarians from upper Paleocene and lower Eocene strata, Mead Stream. Scale bars = 100 µm.

1. Haliomma gr. A-K47/4, P30/f451, 99 m, RP5.

2. Haliomma gr. A-L36/3, P30/f533, 166.5 m, RP8.

3. Haliomma gr. A-S28/0, P30/f533, 166.5 m, RP8.

4. Haliomma gr. A-D29/3, P30/f533, 166.5 m, RP8.

5. Haliomma gr. A-R46/1, P30/f533, 166.5 m, RP8.

6. Periphaena? duplus (KOZLOVA). A-032/2, P30/f451, 99 m, RP5.

7. Periphaena? duplus (KOZLOVA). A-S31/3, P30/f451, 99 m, RP5.

8. Periphaena? duplus (KOZLOVA). A-G29/1, P30/f451, 99 m, RP5.

9. Middourium regulare (BORISSENKO). A-W53/0, P30/f1076, 136.9 m, RP7.

10. Dorcadospyris foveolata (EHRENBERG). A-L49/3, P30/f527, 108.2 m, RP6.

11. Dorcadospyris foveolata (EHRENBERG). A-P42/0, P30/f527, 108.2 m, RP6.

12. Dorcadospyris confluens (EHRENBERG). A-W50/1, P30/f527, 108.2 m, RP6.

13. Lophophaena mugaica (GRIGORJEVA). A-B42/1, P30/f527, 108.2 m, RP6.

14. Corythomelissa adunca (SANFILIPPO AND RIEDEL). A-?, P30/f451, 99 m, RP5.

15. Corythomelissa sp. A. B-F36/0, P30/f527, 108.2 m, RP6.

16. Buryella granulata (PETRUSHEVSKAYA). D-P34/0, P30/f451, 99 m, RP5.

17. Buryella foremanae (PETRUSHEVSKAYA). B-U42/0, P30/f527, 108.2 m, RP6.

18. Buryella tetradica FOREMAN. A-F48/0, P30/f1076, 136.9 m, RP7.

19. Buryella tetradica FOREMAN. D-P32/0, P30/f451, 99 m, RP5.

20. Buryella pentadica FOREMAN. C-034/2, P30/f527, 108.2 m, RP6.

21. Pterocodon poculum (NISHIMURA). A-O47/2, P30/f1072, 136.6 m, RP7.

22. Pterocodon poculum (NISHIMURA). A-L37/3, P30/f1076, 136.9 m, RP7.

23. Theocorys? aff. phyzella FOREMAN. C-G39/0, P30/f452, 110.8 m, RP6.

24. Theocorys? aff. phyzella FOREMAN. C-E41/3, P30/f452, 110.8 m, RP6.

25. Theocorys? cf. phyzella FOREMAN. D-U29/3, P30/f452, 110.8 m, RP6.

26. Theocorys? phyzella FOREMAN. B-V33/1, P30/f1072, 136.6 m, RP7.

27. Theocorys? phyzella FOREMAN. A-H34/4, P30/f1072, 136.6 m, RP7.

S98 C. J. Hollis



Paleocene-Eocene radiolarians, Mead Stream $\ S99$