

SPECIAL CASE OF KUMMER'S CONGRUENCE

Autor(en): **Carlitz, L.**

Objekttyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **5 (1959)**

Heft 3: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **04.06.2024**

Persistenter Link: <https://doi.org/10.5169/seals-35487>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

<http://www.e-periodica.ch>

A SPECIAL CASE OF KUMMER'S CONGRUENCE
 (mod 2^e)

by L. CARLITZ, Duke University, Durham

(Reçu le 30 janvier 1959.)

Put

$$\frac{x}{\sinh x} = \sum_1^{\infty} D_{2n} \frac{x^{2n}}{(2n)!}, \quad D_{2n+1} = 0, \quad (1)$$

where [3, p. 27-28]

$$(D + 1)^n - (D - 1)^n = \begin{cases} 2 & (n = 1) \\ 0 & (n < 1); \end{cases}$$

also

$$D^n = (2B + 1)^n = (2 - 2^n) B_n \quad (2)$$

and B_n is defined by

$$\frac{x}{e^x - 1} = \sum_0^{\infty} B_n \frac{x^n}{n!}.$$

The first few values of D_{2n} are

$$D_0 = 1, D_2 = -\frac{1}{3}, \quad D_4 = \frac{7}{15}, \quad D_6 = \frac{31}{21}, \quad D_8 = \frac{127}{15}.$$

If we define

$$\Delta_n = (D^2 - 1)^n = \sum_{r=0}^n (-1)^{n-r} \binom{n}{r} D_{2r} \quad (3)$$

we find that

$$\Delta_0 = 1, \quad \Delta_1 = \frac{2^2}{3}, \quad \Delta_2 = \frac{2^5}{15}, \quad \Delta_3 = \frac{2^9}{105}, \quad \Delta_4 = \frac{2^{11}}{105}.$$

(Compare [1, p. 821].)

By the second part of (2), both numerator and denominator of D_{2n} (in reduced form) are odd. Moreover by the Staudt-Clausen theorem for the Bernoulli numbers B_n , we have

$$p D_{2n} \equiv \begin{cases} -1 & (p \equiv 1 \pmod{2n}) \\ 0 & (p \equiv 1 \pmod{2n}) \end{cases},$$

where p is any odd prime. Thus the denominator of Δ_n can be determined (see below). The numerators of Δ_n for $0 \geq n \geq 4$ are all powers of 2. However this is really too good to be true; indeed we find that

$$\Delta_5 = -\frac{2^{15}}{3 \cdot 7 \cdot 11}, \quad \Delta_6 = \frac{2^{17} \cdot 191}{3 \cdot 5 \cdot 7 \cdot 11 \cdot 13}.$$

It may be of interest to determine the highest power of 2 dividing Δ_n . A result of this sort is analogous to Kummer's congruences for the Bernoulli or Euler numbers [2, Chapter 14]; however the standard results on Kummer's congruences are restricted to odd moduli.

It is convenient to first transform Δ_n . We have, using the symbolic notation,

$$\Delta_n = (D^2 - 1)^n = ((2B + 1)^2 - 1)^n = (4B^2 + 4B)^n = 4^n (B^2 + B)^n$$

and therefore

$$\Delta_n = 2^{2n} \sum_{r=0}^n \binom{n}{r} B_{n+r}. \quad (4)$$

Since $2B_{2r} \equiv 1 \pmod{2}$ it follows readily that the denominator of

$$\sum_{r=0}^n \binom{n}{r} B_{n+r}$$

(in reduced form) is odd. Consequently

$$\Delta_n \equiv 0 \pmod{2^{2n}}. \quad (5)$$

Incidentally (4) is a bit more convenient for computation than (2). For example

$$\Delta_5 = 2^{10} (5B_6 + 10B_8 + B_{10}) = -\frac{2^{15}}{3 \cdot 7 \cdot 11},$$

$$\begin{aligned} \Delta_6 &= 2^{12} (B_6 + 15B_8 + 15B_{10} + B_{12}) \\ &= 2^{12} \left(\frac{1}{42} - \frac{15}{30} + \frac{75}{66} - \frac{691}{2730} \right) = \frac{2^{17} \cdot 191}{3 \cdot 5 \cdot 7 \cdot 11 \cdot 13}. \end{aligned}$$

The congruence (5) can be improved. We make use of the explicit formula

$$B_n = \sum_{r=0}^n \frac{1}{r+1} \sum_{s=0}^r (-1)^s \binom{r}{s} s^n,$$

which yields

$$(B^2 + B)^n = 2^n \sum_{r=0}^{2n} \frac{1}{r+1} \sum_{s=0}^r (-1)^s \binom{r}{s} \left(\frac{1}{2}s(s+1)\right)^n. \quad (6)$$

However allowance must be made for the denominator $r+1$. Clearly the highest power of 2 contained in $r+1$, where $0 \leq r \leq 2n$, is at most 2^e , where e is determined by

$$2^e \leq 2n < 2^{e+1}. \quad (7)$$

Hence by (4) and (6),

$$\Delta_n \equiv 0 \pmod{2^{3n-e}}. \quad (8)$$

This result also can be improved for $n \geq 2$. Suppose first that n is even and let

$$2^k \mid n, 2^{k+1} + n \quad (k \geq 1). \quad (9)$$

Then for arbitrary odd u we have

$$u^n \equiv 1 \pmod{2^{k+2}}. \quad (10)$$

Put

$$S_j = S_j^{(r)} = \sum_{s \equiv j \pmod{4}} \binom{r}{s} \quad (0 \leq j \leq 3).$$

It is easily verified that

$$2(S_1 - S_2) = (1+i)^{r-1} + (1-i)^{r-1}.$$

Hence for r odd

$$S_1 - S_2 = \begin{cases} 0 & (r \equiv 3 \pmod{4}) \\ (-1)^{1/4(r-1)} 2^{1/2(r-1)} & (r \equiv 1 \pmod{4}) \end{cases}$$

Since $s(s+1)/2$ is odd if and only if $s \equiv 1$ or $2 \pmod{4}$, it follows from (10) that ($n > 2$)

$$S = \sum_{s=0}^r (-1)^s \binom{r}{s} \left(\frac{1}{2}s(s+1)\right)^n \equiv S_2 - S_1 \pmod{2^{k+2}},$$

and therefore

$$S \equiv 0 \pmod{2^{k+2}} \quad (r \equiv 3 \pmod{4}), \quad (11)$$

$$S \equiv 0 \pmod{(2^{k+2}, 2^{1/2(r-1)})} \quad (r \equiv 1 \pmod{4}). \quad (12)$$

Now when $r \equiv 1 \pmod{4}$, $r+1$ is divisible by 2 but not by 2^2 ; hence the term $S/(r+1)$ is integral ($\pmod{2}$) provided

$r > 1$. When $r \equiv 3 \pmod{4}$, let 2^{e_r} denote the highest power of 2 dividing $r + 1$. Then it follows from (11) that $S/(r + 1)$ is of the form $2^{k+2-e_r}A$, where A is integral $(\pmod{2})$. The least favorable case arises when e_r has its maximum value e as determined by (7). We have therefore

$$\Delta_n \equiv 0 \pmod{2^{3n-e+k+2}} \quad (13)$$

provided $k + 2 > e$; when $k + 2 \leq e$ we have

$$\Delta_n \equiv 0 \pmod{2^{3n-1}} \quad (14)$$

since for example the term $r = 1$ has denominator 2.

In the next place for n odd, let

$$2^k \mid n - 1, \quad 2^{k+1} \nmid n - 1 \quad (k \geq 1);$$

Then (10) is replaced by

$$u^{n-1} \equiv 1 \pmod{2^{k+2}},$$

where u is odd. Put

$$\begin{aligned} T_j = T_j^{(r)} &= \frac{1}{2} \sum_{s \equiv j \pmod{4}} \binom{r}{s} s(s+1) \\ &= \binom{r}{2} S_{j-2}^{(r-2)} + r S_{j-1}^{(r-1)}. \end{aligned}$$

Thus

$$\begin{aligned} T_1 - T_2 &= \binom{r}{2} (S_3^{(r-2)} - S_0^{(r-2)}) + r (S_0^{(r-1)} - S_0^{(r-1)}) \\ &= \frac{i}{4} \binom{r}{2} \{(1+i)^{r-1} - (1-i)^{r-1}\} + \frac{r}{4} \{(1+i)^r + (1-i)^r\}. \end{aligned}$$

Simplifying we get (for $r \geq 3$)

$$T_1 - T_2 = \begin{cases} \frac{1}{4}r(r+1)2^{\frac{1}{2}(r-1)}i^{\frac{1}{2}(r+1)} & (r \equiv 3 \pmod{4}) \\ \frac{1}{2}r(2i)^{\frac{1}{2}(r-1)} & (r \equiv 1 \pmod{4}). \end{cases}$$

Also ($n > 3$)

$$T = \sum_{s=0}^r (-1)^s \binom{r}{s} \left(\frac{1}{2}s(s+1)\right)^n \equiv T_2 - T_1 \pmod{2^{k+2}}$$

and therefore in particular

$$T \equiv 0 \pmod{(2^{\frac{1}{2}(r-1)}, 2^{k+2})} \quad (r \equiv 3 \pmod{4}). \quad (15)$$

As in the case n even, it will suffice to take $r \equiv 3 \pmod{4}$. Let 2^{e_r} denote the highest power of 2 dividing $r + 1$, and consider $2^{-e_r} T$. For $e_r \leq 2$, we get the exponent -1 ; for $e_r \geq 3$ it follows that $\frac{1}{2}(r - 1) \geq e_r$. Hence if $k + 2 < e_r$, we get the exponent $k + 2 - e_r$, while if $k + 2 \geq e_r$ we again get -1 (at most). Consequently

$$\Delta_n \equiv 0 \pmod{2^{3n-e+k+2}} \quad (k + 2 < e_r), \quad (16)$$

$$\Delta_n \equiv 0 \pmod{2^{3n-1}} \quad (k + 2 \geq e_r). \quad (17)$$

Comparing (16) and (17) with (13) and (14) we may accordingly state the following

THEOREM. *Let $2^e \leq 2n < 2^{e+1}$ and let 2^k denote the highest power of 2 dividing n or $n - 1$ according as n is even or odd. Then Δ_n as defined by (1) and (3) satisfies (16) and (17).*

Applying the Staudt-Clausen theorem, it follows from (3) that

$$p\Delta_n \equiv - \sum_{r>0} (-1)^{n-rm} \binom{n}{rm} \pmod{p},$$

where $p = 2m + 1$ is an odd prime. It can be shown that

$$\sum_{r>0} (-1)^{n-rm} \binom{n}{rm} \equiv \begin{cases} 2 & (p - 1 \mid n) \\ \binom{m}{k} & (p - 1 \mid n + k, 0 < k \leq m) \\ 0 & \text{otherwise.} \end{cases}$$

In particular the prime factors of the denominator of Δ_n are simple and cannot exceed $2n + 1$.

In connection with formula (4) above it may be of interest to cite the formula [2, p. 189]

$$(-1)^n \sum_{r=0}^n \binom{n}{r} \frac{B_{n+r+1}}{n+r+1} + \frac{n! n!}{2(2n+1)!} = 0.$$

REFERENCES

1. G. FROBENIUS, Über die Bernoullischen Zahlen und die Eulerschen Polynome, *Sitzungsberichte der Preussischen Akademie der Wissenschaften* (1910), pp. 809-847.
 2. N. NIELSEN, *Traité élémentaire des nombres de Bernoulli*. Paris, 1923.
 3. N. E. NÖRLUND, *Vorlesungen über Differenzenrechnung*. Berlin, 1924.
- Duke University.