

# NEW FAMILY OF LINEAR TRANSFORMATIONS

Autor(en): **Meir, A.**

Objekttyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **13 (1967)**

Heft 1: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-41552>

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, [www.library.ethz.ch](http://www.library.ethz.ch)

# A NEW FAMILY OF LINEAR TRANSFORMATIONS

by A. MEIR

1. In 1935 J. Karamata [6] defined a linear transformation method which in 1953 was rediscovered by Lototsky [8] and was published in the survey article of R. P. Agnew [2]. Let  $\{P_{nk}\}$  ( $k=0, 1, \dots, n; n=0, 1, \dots$ ) be defined by the identity

$$(1.1) \quad x(x+1)\dots(x+n-1) = \sum_{k=0}^n P_{nk} x^k,$$

and let  $\{t_n\}_0^\infty$ , the Lototsky transform of a sequence  $\{s_n\}_0^\infty$  be

$$t_n = \frac{1}{n!} \sum_{k=0}^n P_{nk} s_k.$$

Some immediate generalizations were given by Wuckovic [11] who replaced the left hand side of (1.1) by  $(x+\alpha)\dots(x+\alpha+n-1)$  ( $\alpha > -1$ ) and by Harlestan [4] who replaced it by  $(x+q-1)\dots(x+nq-1)$ , ( $q > 0$ ). Jakimovski's result [5] included all the previous ones as special cases by defining the  $[F, d_n]$  transformation for any fixed sequence of real or complex numbers  $\{d_n\}_1^\infty$  ( $d_n \neq -1$ ) by

$$(x+d_1)(x+d_2)\dots(x+d_n) = \sum_{k=0}^n P_{nk} x^k.$$

Regularity, inclusion, shift-properties and applications to analytic continuation, Fourier series, etc., have been thoroughly investigated (see [9], [7]).

Here we give certain further possible generalizations of the above methods, some of their main properties and their relations to well-known classical summation methods.

2. For given sequences of real or complex numbers  $\{c_n\}_1^\infty$ ,  $\{d_n\}_1^\infty$  ( $c_n \neq -1, d_n \neq -1$ ) let  $C_0(x) = D_0(x) = 1$  and for  $n \geq 1$

$$(2.1) \quad \begin{cases} C_n(x) = (x+c_1)(x+c_2)\dots(x+c_n) \\ D_n(x) = (x+d_1)(x+d_2)\dots(x+d_n) \end{cases}$$

and let the matrix  $\{r_{nk}\}$  ( $n=0, 1, \dots; k=0, \dots, n$ ) be defined by  $r_{00} = 1$

and for  $n \geq 1$  by

$$(2.2) \quad \sum_{k=0}^n r_{nk} C_k(x) / C_k(1) = D_n(x) / D_n(1).$$

A sequence  $\{s_n\}_0^\infty$  will be called  $[F, c, d]$ -summable to  $\sigma$  if

$$\lim_{n \rightarrow \infty} \sum_{k=0}^n r_{nk} s_k = \sigma$$

The  $[F, d_n]$ -transformation is the special case  $c_n = 0$  for  $n = 1, 2, \dots$ . Other special examples are the following: (i) If  $d_n = 0$  ( $n=1, 2, \dots$ ) and for some fixed  $q > 0$ ,  $c_n = q^{n-1}$  ( $n=1, 2, \dots$ ) then (2.2) yields

$$r_{nk} = (-1)^{n-k} \begin{bmatrix} n \\ k \end{bmatrix} \prod_{j=1}^n (1 + q^{j-1})$$

where  $\begin{bmatrix} n \\ k \end{bmatrix}$  denotes the  $q$ -binomial coefficients (see e.g. [1]). (ii) If  $d_n = n$ ,  $c_n = q^{n-1}$  ( $n=1, 2, \dots$ ) then (2.2) is equivalent to formula (2) of Carlitz [3]. (iii) If  $d_n = 0$  ( $n=1, 2, \dots$ ) then  $\{r_{nk}\}$  is the inverse of  $[F, c_n]$ -transformation (see Formula (8.7) in [9]). (iv) If  $c_k = d_{k-1}$  ( $k=2, 3, \dots$ ),  $c_1 = 0$  then  $\{r_{nk}\}$  is the matrix transforming the  $[F, d_n]$ -sum of  $0, s_0, s_1, \dots$  into the  $[F, d_n]$ -sum of  $s_0, s_1, \dots$ .

3. Regularity. The following recursion relation is an easy conclusion from (2.2)

$$(3.1) \quad (1 + d_n) r_{nk} = (d_n - c_{k+1}) r_{n-1,k} + (1 + c_k) r_{n-1,k-1}$$

where  $r_{00} = 1$  and  $r_{nk} = 0$  if  $k > n$ , if  $k < 0$  or if  $n < 0$ . Iterating (3.1) one shows by induction that for  $n \geq k \geq 1$  we have

$$(3.2) \quad r_{nk} = \sum_{v=k}^n b_{nv} r_{v-1,k-1}$$

where for  $k \leq v \leq n$

$$(3.3) \quad b_{nv} \equiv b_{nv}^{(k)} = \frac{1 + c_k}{1 + d_v} \prod_{j=v+1}^n \left( 1 - \frac{1 + c_{k+1}}{1 + d_j} \right)$$

(an empty product being 1 by definition). We formulate now the following

**THEOREM.** Suppose  $\{c_n\}_1^\infty, \{d_n\}_1^\infty$  satisfy

$$(3.4) \quad 0 < (1 + c_k)(1 + d_{k+v}) \leq 1, \quad k = 1, 2, \dots; v = 0, 1, \dots$$

$$(3.5) \quad \sum_{n=1}^{\infty} |1 + d_n|^{-1} = +\infty.$$

*Then [F, c, d]-method of summation is regular.*

*Proof.* First we observe that from (3.4) it follows by (3.1) that  $r_{nk} \geq 0$  for all  $n$  and  $k$  and from (2.2) that  $\sum_{k=0}^n r_{nk} = 1$  for all  $n$ . Thus to prove regularity it remains to show that for fixed  $k = 0, 1, \dots$  we have  $r_{nk} = 0$  (1) as  $n \rightarrow \infty$ . Now, from (3.1)  $r_{n0} = (d_n - c_1)(1+d_n)^{-1} r_{n-1,0}$ , since  $r_{00} = 1$ , we have

$$r_{n0} = \prod_{k=1}^n \left( 1 - \frac{1+c_1}{1+d_k} \right)$$

and by (3.4)

$$\leq \exp \left\{ - |1 + c_1| \sum_{k=1}^n |1 + d_k|^{-1} \right\}.$$

By (3.5) then

$$(3.6) \quad \lim_{n \rightarrow \infty} r_{n0} = 0.$$

We assume now that for some  $k \geq 1$

$$\lim_{n \rightarrow \infty} r_{n,k-1} = 0$$

and prove that then  $\lim_{n \rightarrow \infty} r_{nk} = 0$  too. This will follow from (3.2) if we establish that, for fixed  $k$ ,  $\{b_{nv}\}$  satisfy the inequalities

$$(3.7) \quad \sum_{v=k}^n |b_{nv}| \leq H < \infty, \quad n = k, k+1, \dots$$

$$(3.8) \quad \lim_{n \rightarrow \infty} b_{nv} = 0, \quad v = k, k+1, \dots$$

Now from (3.3),  $b_{nv} \geq 0$  for all  $n, v$  and by an easy calculation

$$\begin{aligned} \sum_{v=k}^n b_{nv} &= \frac{1 + c_k}{1 + c_{k+1}} \left\{ 1 - \frac{d_k - c_{k+1}}{1 + d_k} \prod_{j=k+1}^n \left( 1 - \frac{1 + c_{k+1}}{1 + d_j} \right) \right\} \\ &\leq \left| \frac{1 + c_k}{1 + c_{k+1}} \right| \left\{ 1 + \left| \frac{d_k - c_{k+1}}{1 + d_k} \right| \right\} \end{aligned}$$

which proves (3.7). Also by (3.4) for fixed  $v \geq k$ ,

$$|b_{nv}| \leq \frac{1 + c_k}{1 + d_v} \exp \left\{ - |1 + c_{k+1}| \sum_{j=v+1}^n |1 + d_j|^{-1} \right\}$$

which by (3.5)

$$\leq 0(1), \quad \text{as } n \rightarrow \infty.$$

This proves (3.8) and thus by induction from (3.6)

$$\lim_{n \rightarrow \infty} r_{nk} = 0, \quad k = 0, 1, \dots$$

completing the proof of the Theorem.

4. The  $\{r_{nk}\}$  defined by (2.2) or (3.1) can be expressed on using divided differences (see e.g. [10] p. 8, formula 1.4 (1)) as follows

$$(4.1) \quad r_{nk} = (1 + c_1)(1 + c_2) \dots (1 + c_k) [-c_1, -c_2, \dots, -c_{k+1}; D_n]$$

where  $[x_1, x_2, \dots, x_k; f]$  denotes the usual divided differences of a function  $f(x)$  at  $x_1, x_2, \dots, x_k$ . This formula suggests another family of possible transformation methods. For a function  $f(x)$  with  $f(0) = 1$ , for a given sequence  $\{c_k\}_1^\infty$  and for  $\lambda > 0$ , let for  $k \geq 0$

$$(4.2) \quad r_k(\lambda) = (-1)^k (\lambda + c_1) \dots (\lambda + c_k) [\lambda + c_1, \dots, \lambda + c_{k+1}; f].$$

If for a sequence  $\{s_n\}_0^\infty$  and for  $\lambda \geq \lambda_0$

$$t(\lambda) = \sum_{k=0}^{\infty} r_k(\lambda) s_k$$

exists and  $\lim_{\lambda \rightarrow \infty} t(\lambda) = \sigma$ , then we say the sequence  $\{s_n\}$  is  $\{f, c\}$  summable to  $\sigma$ . In particular if  $f(x) = (1+x)^{-1}$  then

$$r_k(\lambda) = \frac{(\lambda + c_1) \dots (\lambda + c_k)}{(1 + \lambda + c_1) \dots (1 + \lambda + c_{k+1})}$$

and this transformation method reduces to the classical Abel method if  $c_k = 0$  ( $k = 1, 2, \dots$ ).

Another modification of (4.1) would suggest to define  $\{r_{nk}\}$  for a function  $f(x)$  and a given sequence  $\{c_n\}$  by

$$r_{nk} = (-1)^{n-k} c_{k+1} c_{k+2} \dots c_n [c_k, \dots, c_n; f]$$

which, if  $f(c_n) = \mu_n$ , generates the well known generalized Hausdorff  $[H, c_n; \mu_n]$ -transformation methods.

REFERENCES

- [1] AL-SALAM, W. A. Note on a  $q$ -identity. *Math. Scand.* 5 (1957) 202-204.
- [2] AGNEW, R. P. The Lototsky method for evaluation of series, *Mich. Math. J.* 4 (1957) 105-128.
- [3] CARLITZ, L. Note on a paper of Laksov. *Math. Scand.* 19 (1966) 38-40.
- [4] HARLESTAM, T. *On a family of summation-methods*. Thesis.
- [5] JAKIMOVSKI, A. A generalization of the Lototsky method of summability. *Mich. Math. J.* 6 (1959) 277-290.
- [6] KARAMATA, J. Théorèmes sur la sommabilité exponentielle et d'autres s'y rattachant, *Mathematica*, Cluj 9 (1935) 164-178.
- [7] LORCH, L. and D. J. NEWMAN. On the  $[F, d_n]$  Summation of Fourier Series. *Comm. Pure Appl. Math.* 15 (1962) 109-118.
- [8] LOTOTSKY, A. V. On a linear transformation of sequences and series. *Ivanov Gos. Ped. Inst. Uc. Zap. Fiz. Math. Nauki* 4 (1953) 61-91 (Russian).
- [9] MEIR, A. On the  $[F, d_n]$ -transformations of A. Jakimovski. *Bull. Res. Council Israel* 10 F (1962) 165-187.
- [10] MILNE-THOMSON, L. M. *The calculus of finite differences*. Macmillan Publ. New York, 1960.
- [11] VUČKOVIĆ, V. *Eine neue Klasse von Polynomen und ihre Anwendung in Theorie der Limitierungsverfahren*, Publ. L'Inst. Ac. Serbe 12 (1958) 125-136.

(Reçu le 25 avril 1968)

A. Meir  
Department of Mathematics  
The University of Alberta  
Edmonton, Canada

**vide-leer-empty**