

# **Existence of admissible refinements of measure coverings**

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **14 (1968)**

Heft 1: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **23.05.2024**

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

$= \{ \hat{U}_\iota(\rho) \}_{\iota=1}^{\iota^*}$  is a Stein covering of  $X(\rho)$ . We say then that  $\hat{\mathfrak{U}}(\rho)$  is a measure covering of  $X(\rho)$ .

*Admissible refinements of measure coverings.* Let  $\hat{\mathfrak{U}}(\rho)$  and  $\hat{\mathfrak{U}}^*(\rho)$  be two measure coverings of  $X(\rho)$ . We say that  $\hat{\mathfrak{U}}^*(\rho)$  is an admissible refinement of  $\hat{\mathfrak{U}}(\rho)$  if the following conditions hold:

- 1)  $U_\iota^* \subset \subset U_\iota$  for each  $\iota$ .
- 2) If  $U_{\iota_0 \dots \iota_\lambda}^* = U_{\iota_0}^* \cap \dots \cap U_{\iota_\lambda}^*$  we put  $(U_{\iota_0 \dots \iota_\lambda}^*)_v = \Phi_v^{-1}(U_{\iota_0 \dots \iota_\lambda}^* \times E^n(\rho))$  for each  $v \in \{ \iota_0 \dots \iota_\lambda \}$ . It is now required that  $(U_{\iota_0 \dots \iota_\lambda}^*)_v \subset (U_{\iota_0 \dots \iota_\lambda})_\mu$  for all  $v, \mu \in \{ \iota_0 \dots \iota_\lambda \}$ .
- 3)  $\hat{U}_{\iota_0 \dots \iota_\lambda}^* = \hat{U}_{\iota_0}^* \cap \dots \cap \hat{U}_{\iota_\lambda}^* \subset (U_{\iota_0 \dots \iota_\lambda})_\mu$  for each  $\mu \in \{ \iota_0 \dots \iota_\lambda \}$ .

#### EXISTENCE OF ADMISSIBLE REFINEMENTS OF MEASURE COVERINGS

*Existence Theorem.* For every fixed integer  $s$  we can find, for some  $\rho > 0$ , a sequence  $\mathfrak{U}_s \ll \mathfrak{U}_{s-1} \ll \dots \ll \mathfrak{U}_1 \ll \mathfrak{U}_0$  of finer measure coverings of  $X(\rho)$  each of which is an admissible refinement of the following.

*Proof.* We first construct a measure covering of  $X(\rho)$  for some  $\rho < \min \rho_\iota$ . Let  $\mathfrak{U}_0 = \{ \mathfrak{U}_\iota \}_{\iota=1}^{\iota^*}$  be a Stein covering of  $X_0$  such that  $U_\iota \subset \subset W_\iota$  for  $\iota \in \{ 1, \dots, \iota^* \}$ . Choose a fixed  $\rho_0 < \min \rho_\iota$ . Now the open sets  $\Phi_\iota^{-1}(U_\iota \times E^n(\rho_0))$  cover  $X_0$  and hence they also cover  $X(\rho)$  for some sufficiently small  $\rho$ . Hence  $\mathfrak{U}_0$  defines a measure covering of  $X(\rho)$ . It is also clear that  $\mathfrak{U}_0$  defines a measure covering of  $X(\rho')$  for each  $\rho' \leq \rho$ . Let us now construct  $\mathfrak{U}_1$ . We let  $\mathfrak{U}^* = \{ U_\iota^* \}_{\iota=1}^{\iota^*}$  be a Stein covering such that  $U_\iota^* \subset \subset U_\iota$  always holds. Now we can find  $\rho_1 \leq \rho$  such that  $\{ \hat{U}_\iota^*(\rho_1) = \Phi_\iota^{-1}(U_\iota^* \times E^n(\rho_1)) \}_{\iota=1}^{\iota^*}$  cover  $X(\rho_1)$ . Hence  $\hat{\mathfrak{U}}^*(\rho_1)$  and  $\hat{\mathfrak{U}}(\rho_1)$  are measure coverings of  $X(\rho_1)$ . But we do not yet know if  $\hat{\mathfrak{U}}^*(\rho_1) \ll \hat{\mathfrak{U}}(\rho_1)$ . We claim that if  $\rho_2 \leq \rho_1$  is sufficiently small then  $\hat{\mathfrak{U}}^*(\rho_2) \ll \hat{\mathfrak{U}}(\rho_2)$ . For suppose this is false. Say that 2) fails for  $\hat{\mathfrak{U}}^*(\rho_2)$  and  $\hat{\mathfrak{U}}(\rho_2)$  when  $0 < \rho_2 \leq \rho_1$ . Hence  $\Phi_v^{-1}(U_{\iota_0 \dots \iota_\lambda}^* \times E^n(\rho_2)) - \Phi_\mu^{-1}(U_{\iota_0 \dots \iota_\lambda} \times E^n(\rho_2))$  are non empty for suitable indices while  $\rho_2 \rightarrow 0$ . Choose a point  $x_t$  from each of these sets. Because  $x_t \in X(\rho_1)$  which is relatively compact we may assume that  $x_t \rightarrow x_0$ . Obviously we get  $x_0 \in \overline{U_{\iota_0 \dots \iota_\lambda}^*} - U_{\iota_0 \dots \iota_\lambda}$ , a contradic-

tion because  $\overline{U_{i_0 \dots i_\lambda}^*} \subset \overline{U_{i_0}^*} \cap \dots \cap \overline{U_{i_\lambda}^*} \subset U_{i_0 \dots i_\lambda}$ . In the same way we can prove that condition 3) is satisfied if  $\rho_2$  is sufficiently small and the theorem is clear.

### GENERAL THEORY

Let  $G$  be an analytic manifold. We put  $\hat{G} = G \times E^n(\rho_1)$  where  $\rho_1$  is an  $n$ -tuple of positive numbers. Let  $\pi: \hat{G} \rightarrow E^n(\rho_1)$  and  $\mathfrak{P}: \hat{G} \rightarrow G$  be the projection maps.  $\hat{G}^* \subset \hat{G}$  denotes an open subset and  $G^* = \hat{G}^* \cap G \times \{0\}$ .

The set  $G^*$  can be identified with an open subset of  $G$ . We denote by  $\alpha: G^* \times E^n(\rho_1) \rightarrow \hat{G}^*$  a biholomorphic fiber preserving map, i.e.  $\pi \circ \alpha = \pi^*$  where  $\pi^*: G^* \times E^n(\rho_1) \rightarrow E^n(\rho_1)$  is the natural projection. Let  $\rho \leq \rho_2 = \gamma \rho_1 < \rho_1$  where  $0 < \gamma < 1$  is a fixed number. We put  $\hat{G}(\rho) = G \times E^n(\rho)$ . If  $f$  is a holomorphic function on  $\hat{G}(\rho)$  we write  $f = \sum a_v (t/\rho)^v$  with  $a_v \in I(G)$ . We define the norm  $\|f\|_\rho$  of  $f$  by  $\|f\|_\rho = \sup_v \{ \sup_v |a_v(G)| \}$ .

If  $f \in I(\hat{G}(\rho))$  we see that  $f \circ \alpha$  is a well defined function on  $G^* \times E^n(\rho)$  because  $\alpha$  is fiber preserving. We define  $\|f \circ \alpha\|_\rho$  using  $G^*$  instead of  $G$  as above. We have the proposition:

*Proposition 1.* There exists a constant  $K$  such that  $\|f \circ \alpha\|_\rho \leq K \|f\|_\rho$  where  $K = K(\rho_2)$  is independent of  $\rho \leq \rho_2$ .

*Proof.* We write  $f = \sum_{|v|=0}^{\infty} a_v (t/\rho)^v$  with  $a_v \in I(G)$ . Now we get  $f \circ \alpha = \sum (a_v \circ \mathfrak{P} \circ \alpha) (t/\rho)^v$  because  $\alpha$  is fiber preserving. Since  $\mathfrak{P}(\hat{G}^*) \subset G$  we get  $|a_v \circ \mathfrak{P}(\hat{G}^*)| \leq |a_v(G)| \leq \|f\|_\rho$ . Now  $a_v \circ \mathfrak{P} \circ \alpha$  admits a Taylor series:  $a_v \circ \mathfrak{P} \circ \alpha = \sum C_{v\lambda} (t/\rho)^\lambda$  with  $C_{v\lambda} \in I(G^*)$ . Since  $|\sum C_{v\lambda} (t/\rho)^\lambda| \leq \|f\|_\rho$  in  $G^* \times E^n(\rho_1)$  and  $\rho \leq \rho_2 = \gamma \rho_1$  Cauchy's inequalities give us  $|C_{v\lambda}(G^*)| \leq \|f\|_\rho \gamma^{|\lambda|}$ . Let us put  $b_\mu = \sum_{v+\lambda=\mu} C_{v\lambda}$ . We get  $|b_\mu(G^*)| \leq \|f\|_\rho \sum \gamma^{|\lambda|} = \|f\|_\rho (1-\gamma)^{-n} = K \|f\|_\rho$ . Now we can write  $f \circ \alpha = \sum_v a_v \circ \mathfrak{P} \circ \alpha (t/\rho)^v = \sum_{\lambda,v} C_{v\lambda} (t/\rho)^\lambda (t/\rho)^v = \sum_\mu b_\mu (t/\rho)^\mu$ . By definition we have  $\|f \circ \alpha\|_\rho = \sup_\mu |b_\mu(G^*)| \leq K \|f\|_\rho$ .

Let us now consider  $\mathbf{h} = (h_{v\mu})$  which is a  $q \times q$  matrix with  $h_{v\mu} \in I(\hat{G})$ . The  $h_{v\mu}$  are also assumed to be bounded on  $\hat{G}$ .