

§3. Applications to theorems of the Schne-Landau type

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **15 (1969)**

Heft 1: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

exactly as (2.2) (a) implies (2.7). Since now $\gamma > \sigma_r \geq \rho$, the above condition in its turn implies

$$\overline{\lim}_{n \rightarrow \infty} \max_{l_n \leq l_m < l_n + \varepsilon l_n} (b_n + b_{n+1} + \dots + b_m) = o_R(1), \quad \varepsilon \rightarrow 0.$$

By Theorem A with hypothesis (1.2) (a) and $a = b = 0$, it follows that $\sum a_n l_n^{-s}$ is convergent for any σ such that $\sigma \geq \gamma > \sigma_r$ and therefore $\sigma_0 \leq \sigma_r$. But, in any case, $\sigma_0 \geq \sigma_k \geq \sigma_r$ for $0 \leq k < r$ and so we have the conclusion (2.5).

In the preceding argument we have supposed that $\sigma_r < \infty$ since $\sigma_r = \infty$ implies trivially $\sigma_k = \infty$.

§ 3. APPLICATIONS TO THEOREMS OF THE SCHNEE-LANDAU TYPE

Theorem II given next is the simplest of the theorems of the type mentioned above and it is a direct combination of Theorems I, B. Theorems V, VI are generalizations, respectively of Ananda-Rau's and Ganapathy Iyer's extensions of the Schnee-Landau theorem ([2], Theorem 9; [7], Theorem 10), as given by Chandrasekharan and Minakshisundaram ([6], pp. 88-9, Corollaries 3.73, 3.74). Theorems III, IV are apparently new counterparts of Theorems V, VI, the newness consisting in the replacement of the two-sided Tauberian conditions of the latter pair of theorems by analogous one-sided conditions suitably supplemented.

THEOREM II. *Suppose that (i) the Dirichlet series,*

$$\sum_1^\infty \frac{a_n}{l_n^s}, \quad s = \sigma + i\tau,$$

is summable (R, l_n, q) for some $q \geq 0$ when $\sigma > \rho$, (ii) the sum-function $f(s)$ thus defined is regular for $\sigma > \eta$ when $\eta < \rho$, and satisfies the condition

$$f(s) = O(|\tau|^r), \quad r > 0, \quad \text{uniformly for } \sigma \geq \eta + \varepsilon > \eta,$$

(iii) *the coefficients a_n of the Dirichlet series satisfy ONE of the two alternatives (a), (b) of (2.2), but with $\theta(x) \equiv x^{1-(\rho-\eta)/r}$. Then the Dirichlet series is summable (R, l_n, k), $0 \leq k < r$, for*

$$\sigma \geq \frac{(r-k)\rho + k\eta}{r}.$$

Proof. By Theorem B, the Dirichlet series is summable (R, l_n, r'), $r' > r$, for $\sigma > \eta$ and hence $\sigma_r \leq \eta < \rho$. Therefore it is evident from the proof of

Theorem I (A) ending with (2.10) that the Dirichlet series is summable (R, l_n, k) , $0 \leq k < r'$, for

$$\sigma \geq \frac{(r' - k)\rho + k\eta}{r'},$$

whence the desired conclusion follows when we let $r' \rightarrow r$.

THEOREM III. *In Theorem II, let ρ be replaced by $\alpha+1$ in hypotheses (i) and (ii); also let hypothesis (iii) be replaced by*

$$a_n = O_R [l_n^\alpha (l_n - l_{n-1})], \quad l_n - l_{n-1} = O \left(l_n^{\frac{r-\alpha+\eta}{r+1}} \right). \quad (3.1)$$

Then the conclusion is that $\sum a_n l_n^{-s}$, $s = \sigma + it$, is summable (R, l_n, k) , $0 \leq k < r$, for

$$\sigma > \frac{(r - k)(\alpha + 1) + (k + 1)\eta}{r + 1}. \quad (3.2)$$

Proof. As in the proof of Theorem II, the series $\sum a_n l_n^{-s}$ is summable (R, l_n, r') , $r' > r$, for $\sigma > \eta$ where now $\eta < \alpha + 1$, so that $\sigma_{r'} \leq \eta < \alpha + 1$. We begin by choosing γ and correspondingly $\theta(x)$ as follows:

$$\eta < \gamma < \alpha + 1, \quad \theta(x) \equiv x^{(r'-\alpha+\gamma)/(r'+1)}. \quad (3.3)$$

Then, since $r' > r$ and $\gamma > \eta$, we have

$$\frac{r' - \alpha + \gamma}{r' + 1} > \frac{r - \alpha + \gamma}{r + 1} > \frac{r - \alpha + \eta}{r + 1}.$$

And so (3.1) gives us, as $n \rightarrow \infty$,

$$a_n = O_R \left[l_n^\alpha l_n^{\frac{r-\alpha+\eta}{r+1}} \right] = o_R \left[l_n^\alpha l_n^{\frac{r'-\alpha+\gamma}{r'+1}} \right] = o_R [l_n^\alpha \theta(l_n)]. \quad (3.4)$$

Also, if $l_n \leq l_m < l_n + \varepsilon \theta(l_n)$, (3.1) again gives us as $n \rightarrow \infty$,

$$a_{n+1} + a_{n+2} + \dots + a_m = \begin{cases} O_R [l_m^\alpha (l_m - l_n)] & \text{if } \alpha \geq 0, \\ O_R [l_n^\alpha (l_m - l_n)] & \text{if } \alpha < 0, \end{cases}$$

so that, whether $\alpha \geq 0$ or $\alpha < 0$,

$$a_{n+1} + a_{n+2} + \dots + a_m = O_R [l_n^\alpha \varepsilon \theta(l_n)]. \quad (3.5)$$

In (3.4) and (3.5),

$$l_n^\alpha \theta(l_n) = l_n^{\rho'} \quad \text{where} \quad \rho' = \alpha + \frac{r' - \alpha + \gamma}{r' + 1} (> \gamma).$$

Hence, combining (3.4) and (3.5), we get

$$\overline{\lim}_{n \rightarrow \infty} \max_{l_m \leq l_n < l_n + \varepsilon \theta(l_n)} \frac{a_n + a_{n+1} + \dots + a_m}{l_n^{\rho'}} = o_R(1), \quad \varepsilon \rightarrow 0. \quad (3.6)$$

(3.6) and the fact, following from Theorem B, that $\sum a_n l_n^{-s}$ is summable (R, l_n , r'), enables us to use (2.10) in the proof of Theorem I (A) with r , ρ replaced by r' , ρ' respectively, so as to infer that $\sum a_n l_n^{-s}$ is summable (R, l_n , k), $0 \leq k < r'$, for

$$\sigma \geq \frac{(r' - k) \rho' + k\gamma}{r'} = \frac{(r' - k)(\alpha + 1) + (k + 1)\gamma}{r' + 1}.$$

This yields (3.2) as required when we let $r' \rightarrow r$ and recall that $\gamma (> \eta)$ can be taken arbitrarily close to η .

THEOREM IV. *In Theorem III, (3.1) alone can be changed to*

$$\left. \begin{aligned} \sum_{v=1}^n (a_v + |a_v|) l_v^p (l_v - l_{v-1})^{1-p} &= O(l_n^{p(\alpha+1)+1})^1, \quad l_n - l_{n-1} = \\ &= O\left[l_n^{\frac{r-\alpha-p-1+\eta}{r+1-p-1}}\right], \quad p > 1, \quad \alpha + 1 + p^{-1} \geq 0, \end{aligned} \right\} \quad (3.7)$$

with the conclusion changed in consequence to the assertion that $\sum a_n l_n^{-s}$ is summable (R, l_n , k), $0 \leq k < r$, for

$$\sigma > \frac{(r - k)(\alpha + 1) + (k + 1 - p^{-1})\eta}{r + 1 - p^{-1}}. \quad (3.8)$$

Proof. We observe that Theorem III may be viewed as the limiting case $p = \infty$ of Theorem IV.

The proof itself is similar to that of Theorem III with the difference that the choice of γ and $\theta(x)$ in (3.3) is now altered as below:

$$\eta < \gamma < \alpha + 1, \quad \theta(x) \equiv x^{(r' - \alpha - p^{-1} + \gamma)/(r' + 1 - p^{-1})}$$

¹⁾ We suppose that $l_0 = 0$.

And furthermore the step corresponding to (3.6) is obtained as follows. Writing $1 - 1/p = 1/p'$, we get, for $l_n \leq l_m < l_n + \varepsilon\theta(l_n)$,

$$\begin{aligned}
 a_{n+1} + a_{n+2} + \dots + a_m &\leq a_{n+1} + |a_{n+1}| + \dots + a_m + |a_m| \\
 &= \sum_{v=1}^{m-n} (a_{v+n} + |a_{v+n}|) l_{v+n} (l_{v+n} - l_{v+n-1})^{(1-p)/p} \times \\
 &\quad \times \frac{(l_{v+n} - l_{v+n-1})^{1/p'}}{l_{v+n}} \\
 &\leq \left[\sum_{v=1}^{m-n} (a_{v+n} + |a_{v+n}|)^p l_{v+n}^p (l_{v+n} - l_{v+n-1})^{1-p} \right]^{1/p} \times \\
 &\quad \times \left[\sum_{v=1}^{m-n} \frac{l_{v+n} - l_{v+n-1}}{l_{v+n}^{p'}} \right]^{1/p'} \\
 &= O \left[l_m^{\alpha+1+1/p} \frac{(l_m - l_n)^{1/p'}}{l_{n+1}} \right] (n \rightarrow \infty) \\
 &= O \left[l_n^{\alpha+1+1/p} \frac{\{\varepsilon\theta(l_n)\}^{1/p'}}{l_n} \right] \tag{3.9}
 \end{aligned}$$

where we have used the hypothesis (3.7) in the passage to the step preceding (3.9). Taking $m = n+1$ in the step preceding (3.9), we get also

$$\begin{aligned}
 a_{n+1} &= O_R \left[l_n^{\alpha+1+1/p} \frac{(l_{n+1} - l_n)^{1/p'}}{l_{n+1}} \right] (n \rightarrow \infty) \\
 &= O_R \left[l_{n+1}^{\alpha+1/p} l_{n+1}^{(r-\alpha-p^{-1}+\eta)/(r+1-p^{-1})p'} \right] \\
 &= o_R \left[l_{n+1}^{\alpha+1/p} \{\theta(l_{n+1})\}^{1/p'} \right]. \tag{3.10}
 \end{aligned}$$

From (3.9) and (3.10) with $n+1$ changed to n , we obtain, instead of (3.6) in the proof of Theorem III,

$$\overline{\lim}_{n \rightarrow \infty} \max_{l_n \leq l_m < l_n + \varepsilon\theta(l_n)} \frac{a_n + a_{n+1} + \dots + a_m}{l_n^{\rho'}} = o_R(1), \quad \varepsilon \rightarrow 0,$$

where

$$\rho' = \alpha + \frac{1}{p} + \frac{(r' - \alpha - p^{-1} + \gamma)}{(r' + 1 - p^{-1})p'}.$$

After this the proof is completed exactly like that of Theorem III subsequent to (3.6).

It may be observed that the assumption $\alpha+1+p^{-1} \geq 0$ involves no loss of generality since $\alpha+1+p^{-1} < 0$ makes successively $a_n + |a_n| \equiv 0$, $a_n \equiv 0$ and so $\sigma_r = -\infty$ for all $r \geq 0$.

THEOREM V. *In Theorem II, let hypothesis (i) be omitted on account of its being implicit (with $q = 0$, $\rho = \alpha+1$) in hypothesis (iii) modified as under. Let hypothesis (ii) be retained with ρ changed to $\alpha+1$, and hypothesis (iii) replaced by*

$$a_n = O [l_n^\alpha (l_n - l_{n-1})]. \quad (3.11)$$

Then the conclusion is that $\sum a_n l_n^{-s}$ is summable (R, l_n, k), $0 \leq k < r$, for σ satisfying (3.2).

THEOREM VI. *If, in Theorem V, (3.11) alone is changed to*

$$\sum_{v=1}^n |a_v|^p l_v^p (l_v - l_{v-1})^{1-p} = O [l_n^{p(\alpha+1)+1}], \quad p > 1, \quad \alpha + 1 + p^{-1} \geq 0,$$

the conclusion will become the assertion that $\sum a_n l_n^{-s}$ is summable (R, l_n, k), $0 \leq k < r$, for σ satisfying (3.8).

The proofs of Theorems V, VI are omitted, being obvious simplifications of those of Theorems III, IV, involving the use of Theorem I (A) with hypothesis (2.2) (b) instead of (2.2.) (a) as formerly. Theorems V and VI, as pointed out by Chandrasekharan and Minakshisundaram, yield Ananda Rau's and Ganapathy Iyer's extensions of the Schnee-Landau theorem when $\alpha \rightarrow +0$.

§ 4. FURTHER APPLICATIONS

Theorem I (A) is a base which, combined with Theorem B, produces Theorem II, and in this sense Theorem I (A) may be said to correspond to Theorem II. There are results corresponding to each of Theorems III-VI in the same sense. For instance, Deduction 1 below corresponds to Theorem III and shows how other deductions corresponding to Theorems IV-VI may be formulated. Deductions 2,3 are further examples of results based on Theorem I.

DEDUCTION 1. (A) *In Theorem I (A), suppose that $\sigma_r < \alpha+1$ and that (2.2) (a) is replaced by*

$$a_n = O_R [l_n^\alpha (l_n - l_{n-1})], \quad l_n - l_{n-1} = O(l_n^{(r-\alpha+\sigma_r)/(r+1)}). \quad (4.1)$$