

§4 Basis and fundamental sets

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **22 (1976)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **23.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

3.7 The following construction would help in developing the case where $|X|$ is even along the lines of 3.2-3.5, which I won't do. Let X be of odd order $|X| = 2g + 1$, and define $X' = X \amalg \{X\}$, thus $|X'| = 2g + 2$. We have a natural linear map

$$2^X \rightarrow 2^{X'}$$

and this is compatible with p, e, Q, q_0 . Composing this with the passage to the quotient, I have a linear isomorphism

$$2^X \rightarrow P_2(X'),$$

and by compatibility with p, p' , isomorphisms

$$\begin{aligned} 2_+^X &\rightarrow P_2^+(X') \\ 2_-^X &\rightarrow P_2^-(X'). \end{aligned}$$

The first is compatible with e, e' , and with the canonical quadratic forms if g is odd. The second is compatible with Q, Q' if g is even.

§ 4 BASIS AND FUNDAMENTAL SETS

4.1 *Normal basis.* Let (J, e) be a symplectic pair. A *normal basis* for (J, e) is a basis $(x_i)_{i \in I}$ for J with the property that $e(x_i, x_j) = 1$ for $i \neq j$, the set of ordered normal basis (i.e. for $I = \{1, \dots, 2g\}$ if $2g = \dim J$) will be denoted $ONB(J, e)$. The symplectic group $Sp(J, e)$ clearly acts on $ONB(J, e)$ and it does it simply transitively, because if two ordered normal bases for (J, e) are given, the unique linear automorphism transforming one into the other is obviously symplectic.

I have not yet shown that the set $ONB(J, e)$ is non-empty, this we will see as a consequence of the following construction, that relates symplectic basis (0.1) with normal basis. The set $SB(J, e)$ of symplectic basis is a torsor over $Sp(J, e)$, thus if $ONB(J, e)$ is non-empty, both torsors should be isomorphic and indeed there would be as many isomorphisms as elements in the group $Sp(J, e)$. What I proceed to exhibit now is a definite isomorphism

$$\alpha: SB(J, e) \rightarrow ONB(J, e)$$

with inverse β . If

$$x \in SB(J, e), x = (x_1, \dots, x_g, x'_1, \dots, x'_g)$$

let's put $y = \alpha(x)$, then by definition

$$\begin{aligned} y_{2k-1} &= x_1 + \dots + x_k + x'_1 + \dots + x'_{k-1} \\ y_{2k} &= x_1 + \dots + x_{k-1} + x'_1 + \dots + x'_k \quad k = 1, \dots, g. \end{aligned}$$

As for the inverse, if $y \in ONB(J, e)$, and $x = \beta(y)$, then one gets from the definition of α

$$\begin{aligned} x_k &= y_1 + \dots + y_{2k-2} + y_{2k-1} \\ x'_k &= y_1 + \dots + y_{2k-2} + y_{2k} \quad k = 1, \dots, g. \end{aligned}$$

It is clear from this definition that α is compatible with the actions of $Sp(J, e)$ on both sets.

4.2 Azygetic sets. Let (S, Q) be a symplectic torsor over a symplectic pair (J, e) . A subset $A \subset S$ is *azygetic* if for any three different elements $s_1, s_2, s_3 \in A$ one has $Q(s_1) + Q(s_2) + Q(s_3) + Q(s_1 + s_2 + s_3) = 1$, or equivalently if $e(s_1, +s_2, s_1 + s_3) = 1$. A is *homogeneous* if Q is constant on it, i.e. if either $A \subset S^+$ or $A \subset S^-$. And the subset A is *linearly independent* if for some (or equivalently, for any) $s \in A$, the subset $s + (A - \{s\}) \subset J$ is linearly independent, or equivalently if $A + A$ spans a subspace of J of dimension $|A| - 1$.

Let A be an azygetic subset, $s \in A$, and let $B = s + (A - \{s\})$, I will show that the only possible linear relation on B is $\sum_{x \in B} x = 0$. Indeed, if $\sum \lambda_x x = 0$ is such a relation, for any $y \in B$, one has

$$\begin{aligned} 0 &= e(y, \sum_x \lambda_x x) = \sum_x \lambda_x e(y, x) = \sum_{\substack{x \in B \\ x \neq y}} \lambda_x \\ \sum_{x \neq y} \lambda_x &= 0 \end{aligned}$$

Adding these equations for any $y, y' \in B$, one concludes that $\lambda_y = \lambda_{y'}$, which was to be shown. As a consequence of this, it follows that any azygetic subset of odd order is linearly independent, and that an azygetic subset has at most $2g + 2$ elements. It is easy to verify that if A is an azygetic subset of odd order and if $s = \sum_{t \in A} t$, $A \cup \{s\}$ is still azygetic.

4.3 Basis for symplectic torsors. A *basis* for a symplectic torsor (S, Q) over (J, e) is a maximal homogeneous, linearly independent, azygetic subset of S . A basis has exactly $2g + 1$ elements, where g is the genus of (S, Q) . This comes from the fact that any symplectic torsor is isomorphic to one of the form (S_X, Q_X) constructed in § 3 because of the uniqueness result in 1.4, that for S_X , $X \subset S_X$ is clearly a basis with $2g + 1$ elements, and that a linearly independent subset can have at most $2g + 1$ elements.

The set of ordered basis for (S, Q) will be denoted by $OB(S, Q)$, the group $Sp(S, Q)$ acts on it.

The following construction is fundamental. Let $X \subset S$ be a basis, we have then a map

$$F_X: 2^X \rightarrow E(S)$$

(cf. 1.5.a) for the definition of $E(S)$), defined by

$$F_X(A) = \sum_{s \in A} s$$

It is clear that F_X is a group homomorphism, that sends subsets of X of even (resp. odd) order into J (resp. S), thereby inducing a linear homomorphism

$$\sigma_X: 2_+^X \rightarrow J$$

and a map compatible with the respective group actions

$$f_X: 2_-^X \rightarrow S.$$

To proceed further, let's choose a total order on X , $X = \{s_0, \dots, s_{2g}\}$. Then, the $X_i = \{s_0, s_i\}$ (resp. $x_i = s_0 + s_i$) for $i = 1, \dots, 2g$ constitute an ordered normal basis for 2_+^X (resp. J), and as $\sigma_X(X_i) = x_i$ we have that σ_X is a symplectic isomorphism. It follows that f_x is a bijection, and indeed f_x defines an isomorphism of symplectic torsors between (S_X, Q_X) and (S, Q) . To see this, we have to prove that if $A, A' \subset X$ are such that $|A| \equiv |A'| \pmod{4}$, then

$$Q\left(\sum_{s \in A} s\right) = Q\left(\sum_{s \in A'} s\right).$$

We know that Q is constant on X , and the condition on X of being azygetic means that for any three different $s_1, s_2, s_3 \in X$, $Q(s_1 + s_2 + s_3)$ is different from the value of Q on X . From this remark, the fact to be proved follows easily by induction and using the defining property (1.1.1) of symplectic torsors. For example, if $|A| = 5$, and we order $A = \{s_1, \dots, s_5\}$, we have

$$Q(\Sigma s_1) + Q(s_1) = Q(s_1 + s_2 + s_3) + Q(s_1 + s_4 + s_5)$$

because $e(s_2 + s_3, s_4 + s_5) = 0$, thus

$$Q(s_1) = Q(\Sigma s_i).$$

Summing up: starting from a basis $X \subset S$, one gets an isomorphism of symplectic pairs

$$\sigma_X: (J_X, e_X) \xrightarrow{\sim} (J, e)$$

underlying an isomorphism of symplectic torsors

$$f_X: (S_X, Q_X) \simeq (S, Q).$$

As a consequence of this, we have that a basis is necessarily contained in S^+ for $g \equiv 0, 1 \pmod{4}$, in S^- for $g \equiv 2, 3 \pmod{4}$ (cf. 3.2.1).

4.4 PROPOSITION. *The set $OB(S, Q)$ of ordered basis for a symplectic torsor (S, Q) is a torsor over the group $Sp(S, Q)$. Moreover, the map*

$$OB(S, Q) \rightarrow ONB(J, e)$$

defined by

$$(s_i)_{0 \leq i \leq 2g} \mapsto (s_0 + s_i)_{1 \leq i \leq 2g}$$

is an isomorphism of torsors over $Sp(S, Q) \simeq Sp(J, e)$.

4.4.1 Proof. The map defined above is clearly compatible with the actions of $Sp(S, Q)$, $Sp(J, e)$ and the isomorphism between these groups described in 1.4. To prove the proposition, it is enough to show that this map is bijective. As $OB(S, Q)$ is non-empty and $ONB(J, e)$ is a torsor, this map is onto. It is injective too, because starting from the $x_i = s_0 + s_i$ I can recover the s_i in the following way. If $s = \sum_{0 \leq i \leq 2g} s_i$, by the identification $S \simeq Q(J, e)$ in 1.5, s corresponds to the unique quadratic form q_s on J whose value on each of the x_i is 1 as it can be easily seen, thus s can be defined in terms of the x_i ; but then

$$s_i = s + \sum_{j \neq i} x_j \quad (0 \leq i \leq 2g, 1 \leq j \leq 2g).$$

4.5 Fundamental sets. A *fundamental set* for a symplectic torsor (S, Q) is a maximal azygetic subset $F \subset S$. Any basis X for S defines a fundamental set, it suffices to put $F_X = X \cup \{s_X\}$, where $s_X = \sum_{s \in X} s$. Also, if F is a fundamental set and if $x \in J$, $x + F$ is a fundamental set too, as it is easily seen. In fact, for any fundamental set F , there exists a basis X and an $x \in J$ such that $F = x + F_X$. Let $F = \{t_0, \dots, t_{2g+1}\}$ be an ordering of F , it is clear that if

$$x_i = t_0 + t_i \quad (1 \leq i \leq 2g+1),$$

the x_i for $1 \leq i \leq 2g$ constitute a normal basis for J , thus there exists a unique ordered basis $X = \{s_0, \dots, s_{2g}\}$ for S such that $x_i = s_0 + s_i$ (4.4). Then, if $x = s_0 + t_0$, we have $t_i = x + s_X$, because $\sum t_i = 0$ and $s_X = \sum s_i$.

Observe that a fundamental set arising from a basis is homogeneous iff g is even. Indeed, it is homogeneous iff $2g + 1 \equiv 1 \pmod{4}$, i.e. iff g is even.

It follows from the last part of prop. 3.2.1 that, in this case, the number of odd characteristics in the fundamental sets is congruent to $g \pmod{4}$. We will see that this is a general fact.

4.5.1 PROPOSITION. *Let $O(F)$ be the number of odd characteristics in a fundamental set F . Then $O(F) \equiv g \pmod{4}$. Conversely, for any $l \equiv g \pmod{4}$, and $l \leq 2g + 2$, there are fundamental sets F with $O(F) = l$.*

4.5.2 Proof. We may safely restrict ourselves to the case where the symplectic torsor is S_X with its standard basis X , and $F = \{A\} + (X \cup \{X\})$ where $A \subset X$ is of even order $|A| = 2k$ (cf. 4.3). Then, in F there are $2k$ characteristics corresponding to subsets of X with $2k - 1$ elements, $2(g - k) + 1$ characteristics with $2k + 1$ elements, and 1 characteristic with $2(g - k) + 1$ elements, namely the ones obtained adding A to respectively the characteristics of the form $\{s\}$ ($s \in A$), $\{s\}$ ($s \notin A$), X . When g is even the second and third types have the same parity; when g is odd the first and third types have the same parity. From these remarks, it is easy to see that the number of elements of the same parity in F and $X \cup \{X\}$ are congruent mod 4, and that with this only restriction, this number can be arbitrary for F by conveniently choosing A . The proposition follows from this and from what was said just before its statement.

4.5.3 In Coble [1], additional material on fundamental sets may be found.

REFERENCES

- [1] COBLE, A. An application of finite geometry to the characteristic theory of the odd and even theta functions. *Trans. A.M.S.* 7 (1906), pp. 241-276.
- [2] FAY, J. Theta functions on Riemann surfaces. *Lecture Notes in Mathematics*, No. 352, Springer Verlag (1973).
- [3] IGUSA, J. *Theta Functions*. Springer Verlag (1972).
- [4] MUMFORD, D. On the equations defining Abelian varieties I. *Invent. Math.* 1 (1966), pp. 287-354.
- [5] ——— Theta characteristics of an algebraic curve. *Ann. Scient. ENS* 4 (1971), pp. 181-192.
- [6] WEBER, H. *Lehrbuch der algebra*, Band 2, Braunschweig (1896).

(Reçu le 20 février 1976)

Neantro Saavedra Rivano

Dept. de Matemática
 Universidad Simón Bolívar
 Apartado 5354
 Caracas, Venezuela