

## §2. The theorem of Kahane

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **23 (1977)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **05.06.2024**

### Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

### Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

2. The function  $t_q = \sum_{p=1}^n \lambda_p \varphi_q(x_p)$  maps the cube  $\mathcal{I}^n$  into the circle  $|t| = 1$ .

3. The transformation  $\Psi$  given by the equalities  $t_q = \sum_{p=1}^n \lambda_p \varphi_q(x_p)$  ( $q = 1, \dots, 2n+1$ ) is one-to-one on  $\mathcal{I}^n$ .

4. For any function  $f$  continuous on  $\mathcal{I}^n$  there exists a function  $g(z)$  continuous on the disk  $|z| \leq 1$ , holomorphic inside that disk, and such that  $f = \sum_{q=1}^{2n+1} g \left( \sum_{p=1}^n \lambda_p \varphi_q(x_p) \right)$ .

The transformation  $\Psi$  gives an embedding of the cube  $\mathcal{I}^n$  into the torus  $|t| = 1$  ( $q = 1, \dots, 2n+1$ ) such that any function continuous on the cube  $\tilde{\mathcal{I}}^n = \Psi(\mathcal{I}^n)$  is represented in the form  $f(t_1, \dots, t_{2n+1}) = \sum_{q=1}^{2n+1} g(t_q)$ , where  $g$  is a function holomorphic in the unit disk. This means in particular that any function continuous on  $\tilde{\mathcal{I}}^n$  has an analytic extension to the polydisk  $|t_q| \leq 1$  ( $q = 1, \dots, 2n+1$ ).

## § 2. *The theorem of Kahane*

Let  $M$  be a complete metric space. We recall that a set is called a set of second category if it is the intersection of a countable family of open sets which are everywhere dense in  $M$ . By the theorem of Baire in a complete metric space no set of second category is empty. The massivity of such sets is characterized by the fact that the intersection of a countable family of sets of second category is again a set of second category and consequently is not empty.

We will say that a statement is true for quasi every element of  $M$  if it is true for a set of elements of second category.

Let us consider an example. Let  $\Phi$  be the space with uniform norm consisting of all functions continuous and non-decreasing on the segment  $\mathcal{I}^1$  ( $0 \leq t \leq 1$ ). It can be shown easily that quasi every element of  $\Phi$  is a strictly increasing function.

In fact, any strictly increasing function belongs to any set defined as  $\varphi(r') < \varphi(r'')$ , where  $r' < r''$  are fixed rational numbers. Any set defined by an inequality of that type is open and everywhere dense in  $\Phi$ , and the set of all such sets is countable.

Let  $\mathcal{I}^n$  be the cube  $\{0 \leq x_i \leq 1, i = 1, \dots, n\}$ ;  $C(\mathcal{I}^n)$ -the space of all functions continuous on  $\mathcal{I}^n$  with the uniform norm;  $\Phi$ -the space of functions continuous and non-decreasing on the segment  $\mathcal{I}^1$  (with the uniform norm);  $\Phi^k = \Phi \times \dots \times \Phi$  the  $k$ -th power of the space  $\Phi$ .

**THEOREM 3.2.1.** *Let  $\lambda_p (p=1, \dots, n)$  be a collection of rationally independent constants. Then for quasi every collection  $\{\varphi_1, \dots, \varphi_{2n+1}\} \in \Phi^{2n+1}$  it is true that any function  $f \in C(\mathcal{I}^n)$  can be represented on  $\mathcal{I}^n$  in the form*

$$f(x) = \sum_{q=1}^{2n+1} g \left( \sum_{p=1}^n \lambda_p \varphi_q(x_p) \right),$$

where  $g$  is a continuous function.

### § 3. The main lemma

We fix a function  $f \in C(\mathcal{I}^n)$ , positive numbers  $\lambda_p (p=1, \dots, n)$  and a positive  $\varepsilon$ . We will denote by  $\Omega_f$  the set of all collections  $\{\varphi_1, \dots, \varphi_{2n+1}\} \in \Phi^{2n+1}$  for each of which there exists a continuous function  $h$  such that  $\|h\| \leq \|f\|$  and  $\|f(x) - h \left( \sum_{p=1}^n \lambda_p \varphi_q(x_p) \right)\| < (1-\varepsilon) \|f\|$ . The latter inequality is strict and consequently the set  $\Omega_f$  is open.

The idea of the construction is contained in the following statement.

**LEMMA 3.3.1.** *If  $\|f\| \neq 0$ , the numbers  $\{\lambda_p\}$  are rationally independent, and  $0 < \varepsilon < \frac{1}{2n+2}$ , than the corresponding set  $\Omega_f$  is everywhere dense in  $\Phi^{2n+1}$ .*

*Proof.* Let us fix an open set  $\Omega \subset \Phi^{2n+1}$  and prove that  $\Omega \cap \Omega_f$  is not empty. This will imply that  $\Omega_f$  is everywhere dense in  $\Phi^{2n+1}$ .

We choose a number  $\delta > 0$  and denote by  $\mathcal{I}_q(j)$  the segment defined by the inequality

$$\begin{aligned} q \cdot \delta + (2n+1)j \cdot \delta &\leq t \leq q \cdot \delta + (2n+1)j\delta + 2n\delta \\ (q=1, \dots, 2n+1, j \text{ is an integer}) . \end{aligned}$$

The value  $\delta$  will be determined below. Now we notice, firstly, that for any  $q$  the segments  $\mathcal{I}_q(j) (j=0, \pm 1, \pm 2)$  are pairwise disjoint and every two consecutive segments are separated by an interval of length  $\delta$  and, secondly, that, every point of the real axis belongs to at least  $2n$  of the sets  $\sum_j \mathcal{I}_q(j), (q=1, \dots, 2n+1)$ .