

§2. (,)-entropy of the set of linear superpositions

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **23 (1977)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **26.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

COROLLARY 5.1.3. For any continuous functions p_i and continuously differentiable functions $q_{1,i}, q_{2,i}, \dots, q_{k,i}$, $k < n$ ($i = 1, 2, \dots, N$) and every region G_n there exists a continuous function that is not equal in G_n to any superposition of the form (V).

§ 2. (ε, δ) -entropy of the set of linear superpositions

We denote by $S(\delta, z)$ the disc of radius δ with centre at z . Let $p(z) = p(x, y)$ and $q(z) = q(x, y)$ be functions defined in a closed region G of the x, y -plane and having the properties:

- a) $p(x, y), \frac{\partial q(x, y)}{\partial x}, \frac{\partial q(x, y)}{\partial y}$ are continuous in G and have modulus of continuity $\omega(\delta)$,
- b) the inequalities $0 < \gamma \leq |\operatorname{grad}[q(r)]| \leq \frac{1}{\gamma}$ and $|p(z)| \leq \frac{1}{\gamma}$, where γ is some constant, are satisfied everywhere in G .

LEMMA 5.2.1. Let $S(\delta, z) \subset G$ and let $\mu_q(t)$ be the function equal to $2 \sqrt{\delta^2 - (t - q(z))^2} |\operatorname{grad}[q(z)]|^{-2}$ on

$$q(z) - \delta |\operatorname{grad}[q(z)]| \leq t \leq q(z) + \delta |\operatorname{grad}[q(z)]|$$

and equal to zero elsewhere. Then

$$\int_{-\infty}^{\infty} |\mu_q(t) - h_1(e(q, t) \cap S(\delta, z))| dt \leq c_1(\gamma) \omega(\delta) \delta^2,$$

where $c_1(\gamma)$ is a constant depending only on γ .

Proof. Let $[a, b] \subset e(q, t) \cap S(\delta, z)$ be the segment of the level curve $e(q, t)$, endpoints a and b , lying on the boundary of $S(\delta, z)$; $[z, a]$ and $[z, b]$ the vectors with origin at z and endpoints at a and b , respectively;

$$\alpha_1 = \gamma([\overrightarrow{z, a}], \operatorname{grad}[q(z)]), \quad \alpha_2 = \gamma([\overrightarrow{z, b}], \operatorname{grad}[q(z)]).$$

We have

$$\begin{aligned} |t - q(z)| &= |q(a) - q(z)| = \left| \int_{s \in [z, a]} \frac{\partial q}{\partial s} ds \right| \\ &= \delta \cos \alpha_1 |\operatorname{grad}[q(z)]| (1 + O(1) \omega(\delta)) \end{aligned}$$

Hence

$$\delta \sin \alpha_1 = \sqrt{\delta^2 - (t - q(z) + 0(\gamma) \delta \omega(\delta))^2} |\operatorname{grad} [q(z)]|^{-2}$$

and similarly

$$\delta \sin \alpha_2 = \sqrt{\delta^2 - (t - q(z) + 0(\gamma) \delta \omega(\delta))^2} |\operatorname{grad} [q(z)]|^{-2}$$

By b) the size of the angle swept out by the tangent vector to the level curve $e(q, t)$ on moving along $[a, b]$ does not exceed $C_2(\gamma) \omega(\delta)$. Therefore

$$h_1([a, b]) = \delta (\sin \alpha_1 + \sin \alpha_2) (1 + 0(\gamma) \omega(\delta)) \\ = 2 \sqrt{\delta^2 - (t - q(z) + 0(\gamma) \delta \omega(\delta))^2} |\operatorname{grad} [q(z)]|^{-2} + 0(\gamma) \delta \omega(\delta).$$

If $\alpha_1 \geq C_3(\gamma) \omega(\delta)$ (C_3 is a sufficiently large constant), then $[a, b] = e(q, t) \cap S(\delta, z)$. Consequently, for

$$|t - q(z)| \leq \theta = \delta \cos [C_3 \omega(\delta)] |\operatorname{grad} [q(z)]| \times (1 + 0(1) \omega(\delta))$$

we have $h_1(e(q, t) \cap S(\delta, z)) = h_1([a, b])$. Since for every t (by b))

$$h_1(e(q, t) \cap S(\delta, z)) \leq C_4(\gamma) \delta (1 + \omega(\delta)),$$

we have

$$\int_{-\infty}^{\infty} |h_1(e(q, t) \cap S(\delta, z)) - \mu_q(t)| dt = \\ = \int_{q(z) - \theta}^{q(z) + \theta} |h_1(e(q, t) \cap S(\delta, z)) - \mu_q(t)| dt + 0(\gamma) \delta^2 \omega(\delta).$$

We now estimate

$$\int_{q(z) - \theta}^{q(z) + \theta} |h_1(e(q, t) \cap S(\delta, z)) - \mu_q(t)| dt = \\ = \int_{q(z) - \theta}^{q(z) + \theta} |h_1([a, b]) - \mu_q(t)| dt \leq \\ \leq 2 \int_{q(z) - \theta}^{q(z) + \theta} (\sqrt{\delta^2 - (t - q(z) + 0(\gamma) \delta \omega(\delta))^2} |\operatorname{grad} [q(z)]|^{-2} \\ - \sqrt{\delta^2 - (t - q(z))^2} |\operatorname{grad} [q(z)]|^{-2}) dt + 0(\gamma) \delta^2 \omega(\delta) \\ = 0(\gamma) \delta^2 \omega(\delta) \int_{-1}^1 \frac{d\tau}{\sqrt{1 - \tau^2}} + 0(\gamma) \delta^2 \omega(\delta) = 0(\gamma) \delta^2 \omega(\delta).$$

Here we have the mean value theorem. This proves the lemma.

LEMMA 5.2.2. Let $p(z), q(z)$ satisfy conditions a) and b); $S(\delta, z) \subset G$; let $f(t)$ be an arbitrary continuous function, uniformly bounded in modulus by the constant m . Then

$$\begin{aligned} & \int \int_{(u, v) \in S(\delta, z)} p(u, v) f(q(u, v)) dudv \\ &= p(z) \left| \operatorname{grad} [q(z)] \right|^{-1} \int_{-\infty}^{\infty} f(t) \mu_q(t) dt + \lambda(z) m \delta^2 \omega(\delta), \end{aligned}$$

where $|\lambda(z)| \leq C_5(\gamma)$.

Proof. Using a) and b) and Lemma 5.2.1 we have

$$\begin{aligned} & \int \int_{S(\delta, z)} p(u, v) f(q(u, v)) dudv \\ &= p(z) \int \int_{(u, v) \in S(\delta, z)} f(q(u, v)) dudv + O(1) m \delta^2 \omega(\delta) \\ &= p(z) \int_{-\infty}^{\infty} \left\{ f(t) \int_{s \in e(q, t) \cap S(\delta, z)} \left| \operatorname{grad} [q(s)] \right|^{-2} ds \right\} dt + O(1) m \delta^2 \omega(\delta) \\ &= p(z) \left| \operatorname{grad} [q(z)] \right|^{-1} \int_{-\infty}^{\infty} \left\{ f(t) \int_{s \in e(q, t) \cap S(\delta, z)} ds \right\} dt + O(\gamma) m \delta^2 \omega(\delta) \\ &= p(z) \left| \operatorname{grad} [q(z)] \right|^{-2} \int_{-\infty}^{\infty} f(t) h_1(e(q, t) \cap S(\delta, z)) dt + O(\gamma) m \delta^2 \omega(\delta) \\ &= p(z) \left| \operatorname{grad} [q(z)] \right|^{-1} \int_{-\infty}^{\infty} f(t) \mu_q(t) dt + O(\gamma) m \delta^2 \omega(\delta). \end{aligned}$$

This proves the lemma.

LEMMA 5.2.3. Suppose that a number $\alpha > 0$ and functions $p(z), q(z), f(t)$ satisfying the conditions of Lemma 5.2.2. are given. If for every integer k such that

$$\min_{z \in G} q(z) \leq t_k = k \delta \frac{\alpha}{m} \leq \max_{z \in G} q(z)$$

and any integer l such that

$$\min_{z \in G} \left| \operatorname{grad} [q(z)] \right| \leq t_l' = l \frac{\alpha}{m} \leq \max_{z \in G} \left| \operatorname{grad} [q(z)] \right|,$$

the inequality

$$\left| \int_{t_k - t_l' \delta}^{t_k + t_l' \delta} f(t) \sqrt{\delta^2 - \left(\frac{t - t_k}{t_l'} \right)^2} dt \right| \leq \alpha \delta^2$$

is satisfied, then for every disc $S(\delta, z) \subset G$

$$\left| \int_{(u,v) \in S(\delta,z)} p(u,v) f(q(u,v)) du dv \right| \leq c_6(\gamma) (\alpha \delta^2 + m \delta^2 \omega(\delta)).$$

Proof. Suppose that a disc $S(\delta, z) \subset G$ is given. By the condition of the lemma there are integers k and l such that $|q(z) - t_k| \leq \delta \alpha / m$ and $\|\operatorname{grad}[q(z)]\| - t_l' \leq \alpha / m$. From Lemma 5.2.2 we obtain

$$\begin{aligned} \left| \int_{(u,v) \in S(\delta,z)} p(u,v) f(q(u,v)) du dv \right| &\leq \frac{|p(z)|}{\|\operatorname{grad}[q(z)]\|} \left| \int_{-\infty}^{\infty} f(t) \mu_q(t) dt \right| \\ &+ c_5(\gamma) m \delta^2 \omega(\delta) \leq \frac{2}{\gamma^2} \left| \int_{t_k - \delta | \operatorname{grad}[q(z)] |}^{t_k + \delta | \operatorname{grad}[q(z)] |} f(t) \sqrt{\delta^2 - \frac{(t - q(z))^2}{\|\operatorname{grad}[q(z)]\|^2}} dt \right| \\ &- \int_{t_k - \delta | \operatorname{grad}[q(z)] |}^{t_k + \delta | \operatorname{grad}[q(z)] |} f(t) \sqrt{\delta^2 - \left(\frac{t - t_k}{t_l'} \right)^2} dt \Big| + \frac{2}{\gamma^2} \alpha \delta^2 + c_5(\gamma) m \delta^2 \omega(\delta) \leq \end{aligned}$$

(by the mean value theorem)

$$\begin{aligned} &\leq \frac{2}{\gamma^2} \alpha \delta^2 + c_5(\gamma) m \delta^2 \omega(\delta) + \frac{2}{\gamma^2} \left(\int_{-1}^1 \frac{\delta m d\tau}{\sqrt{1 - \tau^2}} \right) \delta \frac{\alpha}{m} \\ &+ \frac{2}{\gamma^2} \left(\int_{-1}^1 \frac{\delta^2 m d\tau}{\sqrt{1 - \tau^2}} \right) \frac{\alpha}{m} \leq c_6(\gamma) (\alpha \delta^2 + m \delta^2 \omega(\delta)). \end{aligned}$$

This proves the lemma.

We denote by $F_m = F_m(D; p_1, p_2, \dots, p_N; q_1, q_2, \dots, q_N)$ the set of superpositions of the form

$$f(x, y) = \sum_{i=1}^N p_i(x, y) f_i(q_i(x, y)), \text{ where } \{p_i(x, y)\}$$

and $\{q_i(x, y)\}$ are fixed functions, defined in the closed region D of the x, y plane and satisfying conditions a) and b) with a constant γ not depending on i and $\{f_i(t)\}$ are arbitrary continuous functions, defined on $\{[a_i, b_i]\} = \{[\min_{z \in D} q_i(z); \max_{z \in D} q_i(z)]\}$ and uniformly bounded in modulus by the constant m .

THEOREM 5.2.1. *There exist constants A and B such that if $\varepsilon > Am\omega(\delta)$ then for the (ε, δ) -entropy of the set of functions F_m , $H_{\varepsilon, \delta}(F_m) \leq \frac{B}{\delta} \left(\frac{m}{\varepsilon} \right)^2$, where A and B depend only on γ, N and D .*

Proof. We put

$$R(f(z), \delta) = \max_{S(\delta, z) \subset D} \left| \frac{1}{\pi \delta^2} \iint_{(u, v) \in S(\delta, z)} f(u, v) dudv \right|.$$

We denote by $\mathcal{H}_{\varepsilon, \delta}(F_m)$ the ε -entropy of the space F_m , taking as the distance between the functions $f_1(z), f_2(z) \in F_m$ the number $R(f_1(z) - f_2(z), \delta)$. The inequality $H_{2\varepsilon, \delta}(F_m) \leq \mathcal{H}_{\varepsilon, \delta}(F_m)$ holds owing to the fact that if two functions $f_1(z)$ and $f_2(z)$ are (ε, δ) -distinguishable, then they are ε -distinguishable also in the sense of the metric $R(f_1(z) - f_2(z), \delta)$. We now estimate the value of $\mathcal{H}_{\varepsilon, \delta}(F_m)$. Let k and l be integers such that

$$\min_{z \in D} q_i(z) \leq t_k = k\delta \frac{\alpha}{m} \leq \max_{z \in D} q_i(z)$$

and

$$\min_{z \in D} |\operatorname{grad} [q_i(z)]| \leq t_l' = l \frac{\alpha}{m} \leq \max_{z \in D} |\operatorname{grad} [q_i(z)]|.$$

To compute the function

$$f_\delta(z) = \frac{1}{\pi \delta^2} \iint_{(u, v) \in S(\delta, z)} f(u, v) dudv,$$

where $f(x, y) \in F_m$, $S(\delta, z) \subset D$ to within ε , it is sufficient by Lemma 5.2.3 to give the values of

$$v_i(t_k, t_l') = \frac{1}{\pi \delta^2} \int_{t_k - t_l' \delta}^{t_k + t_l' \delta} f_i(t) \sqrt{\delta^2 - \left(\frac{t - t_k}{t_l'} \right)^2} dt$$

to within $\alpha = \pi \varepsilon / (2 NC_B(\gamma))$ and to assume that δ is small enough so that

$$\varepsilon > \frac{2NC_B(\gamma)m\omega(\delta)}{\pi} = A(\gamma, N)m\omega(\delta).$$

Since $|v_i(t_k, t_l')| \leq C_1 m$, to write the numbers $v_i(t_k, t_l')$ (i, k, l fixed) $\log_2(C_1 m/\alpha)$ binary digits are sufficient. Since

$$|v_i(t_{k+1}, t_l') - v_i(t_k, t_l')| \leq c_8 \frac{1}{\delta^2} \left(\int_{-1}^1 \frac{\delta m d\tau}{\sqrt{1-\tau^2}} \right) \delta \frac{\alpha}{m} = c_9(\gamma) \alpha$$

(here we again use the mean value theorem), to store the numbers $v_i(t_{k+1}, t_l') - v_i(t_k, t_l')$ to within α , $\log_2 C_9$ binary digits are sufficient. Therefore to write the numbers $v_i(t_k, t_l')$ (i, l fixed; k any admissible number)

$C_{10}(\gamma) \left[\log_2 \frac{m}{\alpha} + (b_i - a_i) \frac{m}{\delta \alpha} \right] = \mathcal{H}_{i,l}$ binary digits are sufficient. Consequently the total number of digits sufficient to store all the numbers $v_i(t_k, t_l')$ to within α , that is, to store the functions $f_\delta(z)$ to within ε , is

$$\mathcal{H} = \sum_{i,l} \mathcal{H}_{i,l} \leq N c_{10}(\gamma) \left[\log_2 \frac{m}{\alpha} + (b_i - a_i) \frac{m}{\delta \alpha} \right] \frac{1}{\gamma} \frac{m}{\alpha} \leq \frac{B(\gamma, N, D)}{\delta} \left(\frac{m}{\varepsilon} \right)^2.$$

This proves the theorem.

§ 3. Functional “dimension” of the space of linear superpositions

Suppose that continuous functions $p_i(x, y)$ and continuously differentiable functions $q_i(x, y)$ ($i = 1, 2, \dots, N$) are fixed. Let G be a closed region of the x, y plane. We denote by $F = F(G, \{p_i\}, \{q_i\})$ the set of superpositions of the form $f(x, y) = \sum_{i=1}^N p_i(x, y) f_i(q_i(x, y))$, where $(x, y) \in G$ and $\{f_i(t)\}$ are arbitrary continuous functions of one variable. We are interested in the functional dimension of the set F .

THEOREM 5.3.1. *In every region D of the x, y plane there exists a closed subregion $G \subset D$ such that*

$$r(F(G, \{p_i\}, \{q_i\})) \leq 1.$$

Proof. By Theorem 4.5.1, in D there exists a closed subregion $G^* \subset D$ such that the set of superpositions $F(G^*, \{p_i\}, \{q_i\})$ is closed (in the uniform metric) in $C(G^*)$, and the functions $\{q_i(x, y)\}$ satisfy the condition: for any i , either $\text{grad}[q_i(x, y)] \neq 0$ on G^* or $q_i(x, y) \equiv \text{const}$ on G^* . We show that $r(F(G^*, \{p_i\}, \{q_i\})) \leq 1$. By Banach's open mapping theorem, there exists a constant K such that for any superposition $\sum_{i=1}^N p_i(x, y) f_i(q_i(x, y)) = f(x, y) \in F(G^*, \{p_i\}, \{q_i\})$ there are con-