Zeitschrift:	L'Enseignement Mathématique		
Herausgeber:	Commission Internationale de l'Enseignement Mathématique		
Band:	23 (1977)		
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE		
Artikel:	STABILITY OF PROJECTIVE VARIETIES		
Autor:	Mumford, David		
Kapitel:	§1. Stable points of représentation, examples and Chow forms		
DOI:	https://doi.org/10.5169/seals-48919		

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

## **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

# Download PDF: 02.07.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

§ 1. STABLE POINTS OF REPRESENTATION, EXAMPLES AND CHOW FORMS

For more details on the notations, definitions and properties which follow see Mumford [14], which we will call G.I.T. or Seshadri [20].

Fix k an algebraically closed field,

G a reductive algebraic group over k (i.e.  $G = [\text{semi-simple group} \times \mathbf{G}_m^n]/\text{finite central subgroup}),$ 

V an *n*-dimensional representation of G,  $x \in V$ .

There are three possibilities for x whose equivalent formulations are summarized in table 1.1 below.

1.1.

x unstable	x semi-stable	x stable	
$(a_u)  0 \in \overline{O^G(x)}$	$(a_{ss}) \\ 0 \notin \overline{O^G(x)}$	<ul> <li>(a<sub>s</sub>)</li> <li>i) O<sup>G</sup>(x) is closed</li> <li>in V</li> <li>ii) stab (x) is finite</li> </ul>	
$(b_u)$ $\forall$ non-constant G-invariant homogeneous polyno- mials $f$ f(x) = 0	$(b_{ss}) \exists$ a non-constant <i>G</i> -invariant homogeneous polyno- mial <i>f</i> s.t. $f(x) \neq 0$	(b <sub>s</sub> ) i) $\forall y \in V - O^G(x)$ , $\exists$ a <i>G</i> -invariant polynomial <i>f</i> s.t. $f(x) \neq f(y)$ ii) tr deg <sub>k</sub> k (V) <sup>G</sup> = dimV-dimG	
( $c_u$ ) $\exists$ a 1-PS $\lambda$ of G s.t. the weights of x with respect to $\lambda$ are all positive	( $c_{ss}$ ) $\forall$ 1-PS's $\lambda$ of G the weights of x with respect to $\lambda$ are not all positive	(c <sub>s</sub> ) for all non-trivial 1-PS's $\lambda$ of G, x has both positive and negative weights with respect to $\lambda$	

1.2. REMARKS. i) Recall that a 1-PS (one parameter subgroup)  $\lambda$  of G is just a homomorphism  $\lambda: \mathbf{G}_m \to G$ . Such  $\lambda$  can always be diagonalized in a suitable basis:

$$\lambda(t) = \begin{bmatrix} t^{r_1} & 0 \\ & \cdot & \\ & & \cdot \\ 0 & & t^{r_n} \end{bmatrix}$$

If in this basis  $x = (x_1, ..., x_n)$ , the set of weights of x with respect to  $\lambda$  is the set of  $r_i$  for which  $x_i \neq 0$ .

ii) Unstable is *not* the opposite of stable, but of semi-stable. We will use non-stable as the opposite of stable.

iii) The important part of stability is the condition:  $O^G(x)$  closed in V. In virtually all the cases that will interest us the finiteness of stab (x) will be automatic (but cf. the remark following 1.15).

iv) A point x is stable if it merely has negative weights with respect to every non-trivial 1-PS  $\lambda$ , for then it also has positive weights with respect to  $\lambda$ , namely, its negative weights with respect to  $\lambda^{-1}$ .

v) The proofs of  $c_u \Rightarrow a_u \Rightarrow b_u$  and of  $b_s \Rightarrow a_s \Rightarrow c_s$  are obvious: for example, if  $\lambda$  is a 1-PS for which all weights of x are positive, then  $\lambda(t) x \to 0$  at  $t \to 0$ ; i.e.  $c_u \Rightarrow a_u$ .

vi) The proofs of  $a_s \Rightarrow b_s$  and  $b_u \Rightarrow a_u$  are achieved by reduction to the special case called geometric reducivity of G. A group G is called geometric reductive if

a) whenever  $V_0$  is an invariant codimension-1 subspace of a vector space V in which G is represented, there exists an n for which the codimension-1 invariant subspace  $V^0 \cdot \text{Symm}^{n-1}V \subset \text{Symm}^n V$  has an invariant 1-dimensional complement.

But notice that this is the same as saying that

b) whenever  $x \neq 0$  is a G-invariant point, then there exists a G-invariant polynomial f such that  $f(x) \neq 0$  and f(0) = 0. (Just consider x as a functional on the dual  $\hat{V}$  and apply a) to its kernel there).

And b) is a special case of  $a_s \Rightarrow b_s$ . When char k = 0 we can take the polynomial f to be linear, for by complete reducibility the invariant subspace generated by x is invariantly complemented. A simple example shows this does not happen in char p. Take p = 2, G = SL(2), V = the space of

symmetric bilinear functions on  $k^2$ , and x a non-degenerate skew-symmetric form ( $x \in V$  because p = 2 !). Then x is SL (2)-invariant and there are no G-invariant non-zero linear functionals on V. A quadratic f which does work is the determinant.

vii) The remaining implications  $c_s \Rightarrow a_s$  and  $a_u \Rightarrow c_u$  are essentially consequences of the surjectivity of the natural map

$$\begin{cases} 1\text{PS's }\lambda \text{ of }G\\ \lambda: \mathbf{G}_m \to G \end{cases} \longrightarrow G(k[[t]]) \qquad G(k((t))) \qquad G(k[[t]]) \end{cases}$$

where  $\lambda$  is considered as a k((t))-valued point of G by composition with the canonical map

Spec 
$$k((t)) \rightarrow$$
 Spec  $k[t, t^{-1}] = \mathbf{G}_m$ 

1.3. Let  $V_{ss}$  (resp.  $V_s$ ) denote the Zariski-open cones of semi-stable (resp. stable) points.  $V-V_{ss}$  is the Zariski-closed cone of unstable points. The conditions b of 1.1 tell us that if we try to map  $\mathbf{P}(V)$  to a projective space by invariant polynomials, we can only hope to achieve a well-defined map on  $\mathbf{P}(V)_{ss}$  and an embedding on  $\mathbf{P}(V)_s$ . From the point of view of quotients this can be expressed by:

**PROPOSITION 1.3.** Let  $X = \operatorname{Proj} k [V]^{G}$ . Then there is a diagram

$$\mathbf{P}(V) \supset \mathbf{P}(V)_{ss} \supset \mathbf{P}(V)_{s}$$

$$\pi \left| \begin{array}{c} \pi \\ X \end{array} \right| \\ X \supset X_{s} \end{array}$$

such that i) if  $x, y \in \mathbf{P}(V)_s$ ,  $\pi_s(x) = \pi_s(y) \Leftrightarrow \exists g \in G \text{ s.t. } x = g y$ ii) if  $x, y \in \mathbf{P}(V)_{ss}$ ,  $\pi(x) = \pi(y) \Leftrightarrow \overline{O^G(x)} \cap \overline{O^G(y)} \cap \mathbf{P}(V)_{ss} \neq \emptyset$ .

We now want to look at some examples to illustrate the application of these ideas.

1.4. "BAD" ACTIONS. Using results of T. Kimura and M. Sato [11]<sup>1</sup>), we can give a list of all representations of simple algebraic groups in charac-

ないないのであった。

<sup>&</sup>lt;sup>1</sup>) Plus help given by J. Tits.

teristic 0 in which all vectors are unstable. The point is that there are very few such representations.

G	V
$SL(W)$ $\begin{cases} SL(W) \\ dim W odd \\ Sp(W) \end{cases}$	$W^{i}, \widehat{W}^{i}, 1 \leq k < \dim W$ $\Lambda^{2} W, \Lambda^{2} \widehat{W}$ $\Lambda^{2} W \oplus \widehat{W}, \Lambda^{2} \widehat{W} \oplus W$ $W$
Spin (10)	W or $W \oplus W$ where W is a 16-dimensional half-spin representation

1.5. DISCRIMINANT. If G is semi-simple and char k = 0 then any irreducible representation V has the form  $V = \Gamma(G/B, L)$  for a suitable line bundle L on G/B (B is a Borel subgroup of G). To a point x in V associate the divisor  $H_x$  on G/B which is the zero set of the corresponding section. Except in the extremely unusual case that the set of singular  $H_x$  is of co-dimension > 1, there is an irreducible invariant polynomial  $\delta$ , the discriminant, such that

1)  $\delta(x) = 0 \Leftrightarrow H_x$  is singular

2)  $V - (\delta = 0)$  consists of semi-stable points.

An interesting case is

LEMMA 1.6. Let G = SL(n),  $V = \Lambda^{l}(k^{n})$ . If  $W \subset k^{n}$  is a subspace of codimension l then let  $\Phi_{W}$  denote the natural map  $\Lambda^{2} W \otimes \Lambda^{l-2}(k^{n}) \rightarrow \Lambda^{l}(k^{n})$ . If 2 < l < n - 2 or n is even l = 2 or n - 2, then there is a G-invariant  $\delta$  such that  $\delta(x) = 0 \Leftrightarrow x \in \text{Im}(\Phi_{W})$  for some W.

When l = 2 and n = 2m + 1 we have seen that there are no invariants; corresponding to these cases the Grassmanian of lines in  $\mathbf{P}^{2m}$  in its Plücker embedding in projective space has the unusual property that the singular hyperplane sections are of codimension  $\ge 2$  in the set of all such sections.

Question: if not every point of V is unstable, then is the set of singular hyperplane sections  $H_x$  of codimension 1?

For l = 2 and *n* even or l = 3,  $n \le 8$ , one can check that *x* is unstable  $\Leftrightarrow \delta(x) = 0$ , hence  $\delta$  generates the ring of invariants. It would be nice to have a necessary and sufficient condition for a 3-form to be unstable for higher *n* as well.

1.7. 0-CYCLES. For 
$$G = SL(W)$$
, dim  $W = 2$ ,  
 $V_n = \operatorname{Symm}^n(\widehat{W})$   
 $= \operatorname{vector space of homogeneous polynomials}$   
of degree *n* on *W*,  
 $\mathbf{P}(V_n) = \operatorname{space of 0-cycles of n unordered points of$ 

 $\mathbf{P}(V_n)$  = space of 0-cycles of *n* unordered points on the projective line  $\mathbf{P}(W)$ , the roots of an *f* determining the cycle.

f

If  $f = \sum_{i=0}^{n} a_i x^{n-i} y^i$  and  $\lambda$  is the one-parameter subgroup given by  $t \mapsto \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix}$  in these coordinates, then  $\lambda(t) f = \sum_{i=0}^{n} a_i t^{n-2i} x^{n-i} y^i$ . For f to be stable, the weights (n-2i) associated to the non-zero coefficients of f must lie on both sides of 0: i.e. if  $j \ge n/2$ , neither  $x^j$  nor  $y^j$  divide f.



In fact, the stability of f is equivalent to the same condition with respect to all linear forms  $l: l^j \not> f$  if  $j \ge n/2$ .

Thus  $\mathbf{P}(V_n)_s = \{0 \text{-cycles with no points of multiplicity} \ge n/2\}$  $\mathbf{P}(V_n)_{ss} = \{0 \text{-cycles with no points of multiplicity} > n/2\}.$ 

たいたいですいとうでんでもというないである

1.8. REMARK. In the example above we can also prove that semistability is a purely topological character. I claim that if n is odd and f is unstable then the action of G near  $\overline{f} \in \mathbf{P}(V_n)$  is bad: on all open neighbourhoods of the orbit of  $\overline{f}$ , G acts non-properly and the orbit space is non-Haussdorf. Let's see this for n = 7. Consider the following deformations of a 7-point cycle. (Subscripts indicate multiplicities)



At each intermediate stage the two cycles are projectively equivalent, but the unstable limiting cycle in the right is clearly not equivalent to the limit on the left. In fact, any pair of cycles with the multiplicities indicated on the line t = 0 arise in this way as simultaneous limits of projectively equivalent 0-cycles. Moreover, there are cycles of the same type as the left hand limit in any neighbourhood of the orbit of the right limit—just bring a multiplicity one point in towards the triple point; so the orbit space cannot be Hausdorff near the right limit.

1.9. CURVES. Here G = SL(W), dim W = 3,  $V_n = \text{Symm}^n(W)$ , as before, and a point  $f \in V_n$  defines a plane curve of degree *n*. There is a very simple way to decide the stability of *f*. Represent *f* as below by a triangle of coefficients, *T*.



— 46 —

We can coordinatize this triangle by 3 coordinates  $i_x$ ,  $i_y$ ,  $i_z$  (the exponents

of x, y and z respectively) related by  $i_x + i_y + i_z = n$ . The condition that a line L with equation  $ai_x + bi_y + ci_z = 0$ ,  $(a, b, c) \neq (0, 0, 0)$ , should pass through the centre of this triangle is just a + b + c = 0; if L also passes through a point with integral coordinates then a, b and c can be chosen integral. It is now easy to check that the weights of the 1-PS

のためのです。

$$t \longmapsto \begin{pmatrix} t^a & 0 \\ & t^b & \\ 0 & & t^c \end{pmatrix}$$

at f are just the values of the form defining L at the non-zero coefficients of f. In suitable coordinates every 1-PS is of this form so:

f is unstable  $\Leftrightarrow$  in some coordinates, all non-zero coefficients of f lie to one side of some L

f is stable  $\Leftrightarrow$  for all choices of coordinates and all L, f has non-zero

(resp. semi-stable) coordinates on both sides of L (resp. f has non-zero coordinates on both sides of L or has non-zero coefficients on L).

Roughly speaking, a stable f can only have certain restricted singularities. We summarize what happens for small n, showing the "worst" triangle T for f with given singularities, and the associated L when f is not stable.

1.10. n = 2: We can achieve the diagram below for a non-singular quadric f by choosing coordinates so that  $(1, 0, 0) \in f$  and z = 0 is the tangent line there, so f is never stable. We cannot make the xz coefficient of f zero without making f singular so f is always semi-stable; indeed, we know f always has non-zero discriminant. A singular quadric always has a diagram like that on the right: make (1, 0, 0) the double point. Henceforth, we leave the checking of the diagrams to the reader.



1.11. n = 3: It is well known that in this case the ring of invariants is generated by two invariants, A of degree 4 and B of degree 6. If we set  $\Delta = 27A^3 + 4B^2$ , then up to a constant the classical *j*-invariant is just  $A^3/\Delta$ . The possibilities are:



We remark that in this case, we have

$$\mathcal{M}_1 \cong \mathbf{A}^1$$
$$\stackrel{\bigcirc}{\mathcal{M}_1} = \mathbf{P}^2$$

and that the *j*-invariant is a true modulus. Note that from a moduli point of view all three semi-stable types are equivalent.

1.12. n = 4: There are already quite a few diagram types here. Their enumeration can be summarized by saying that f is unstable if and only if f has a triple point or consists of a cubic and an inflectional tangent line; f is stable if and only if f has only ordinary double points or ordinary cusps (i.e. singularities with local equation  $y^2 = x^3 + \text{higher terms}$ ). The remaining f's with a tacnode (a double point with local equation  $y^2 = x^4 + \text{higher}$ terms) are strictly semi-stable.

1.13. REMARK. The fact that for  $n \ge 4$  curves with sufficiently tame cusps are semi-stable (or even stable!) is a definite problem because

- i) such curves do not appear in the good compactification  $\overline{\mathcal{M}}_g$  of the moduli space of non-singular curves of genus g. But
- ii) if we wish to obtain a compactification of  $\mathcal{M}_g$  as the quotient space of some subset of  $\mathbf{P}(V_n)$  by G, the natural candidate is  $\mathbf{P}(V_n)_{ss}$ ; so these curves must be let in.

For example, when n = 4, we have

M 3,	non-hyperelliptic	$\cong [\mathbf{P}(V_4) - (\delta = 0)]/G$
$\cap$		$\cap$
$\mathcal{M}_3$		$\mathbf{P}(V_4)_s/G$
$\cap$	α	$\cap$
$\overline{\mathcal{M}}_3$	<	$=$ $=$ $=$ $=$ $=$ $=$ $=$ $\mathbf{P}(V_4)_{ss}/G$
	В	

 $\overline{\mathcal{M}}_3$  is the moduli space for "stable" curves of genus 3: (see introduction). Recall from Proposition 1.3 that  $\mathbf{P}(V_4)_{ss}/G$  is just the projectivization of the full rings of invariants of  $\mathbf{P}(V_4)$ . The rational maps  $\alpha$  and  $\beta$  induced by the top isomorphism enable us to make a topological comparison of these two compactifications. Let's see geometrically how cuspidal curves in  $\mathbf{P}(V_4)_{ss}$  prevent  $\alpha$  and  $\beta$  from being continuous.

First  $\alpha$ : the diagram below shows on the left a deformation on  $\mathcal{M}_3$  with limit in  $\overline{\mathcal{M}}_3$ , and on the right the same deformation followed to its limit in  $\mathbf{P}(V_4)_{ss}/G$ .

4



In the limit on the right, the value of the *j*-invariant of the shrinking elliptic curve has been lost! So  $\alpha$  blows up a point representing a curve C with a cusp to the set of points representing joins of an arbitrary elliptic curve with

the desingularization C of C.  $\alpha$  also blows up the point representing a double conic to the family of all hyperelliptic curves.

As for  $\beta$ , look at the double pinching below:



Here it is the manner in which the tangent spaces of the two branches have been glued at the tacnodal point which has been lost in the limiting curve on the left: this glueing corresponds on the left to the relative rate at which the two pinches are made. Thus  $\beta$  has blown up the point corresponding to the double join of two elliptic curves to a family of tacnodal quartics.

「ないたい」という

1.14. SURFACES. Here G = SL(W), dim W = 4 and  $V_n = \text{Symm}^n(W)$  as before. The technique for determining stability here is essentially that given for curves in 1.9 except that one has a tetrahedron T of coefficients and 1-PS's determine central planes, L: and, of course, the computations required to apply the technique are much more complicated (cf. the case n = 4 below). For small n, the situation is summarized below.

<i>n</i>	Type of singularities		STABILITY
n = 2	non-singular singular		semi-stable, not stable unstable
n = 3	non-singular or with ordinary double points of type A1	}	stable
	ordinary double points of type A2		semi-stable, not stable
	triple points, double curve, higher double points	}	unstable
n = 4 (due to Jayant Shah [26])	singularities at most rational double points, or ordinary double curves poss- ibly with pinch points, but no double <i>line</i> , and if reducible then no component a plane, no multiple components		stable
	A triple point whose tangent cone has only ordinary double points; or a double line not as below; or an irrational double point not as below; or a plane plus a cubic meeting in a plane cubic curve with only ordinary double points; or a non- singular quadric counted twice		semi-stable — but not stable
	a) quadruple point, or triple point whose tangent cone has cusp,		
	b) $x = y = 0$ is double line and $\hat{f} \in (x^2, xyz^2, xy^2, y^3)$	}	unstable
	c) a higher double point of form: $f \in (x^2, xy^2, xyz^2, xz^3, y^3z, y^4)$		

# 1.15. Adjoint stability.

**PROPOSITION 1.15.** Let G be any semi-simple group with Lie algebra g. Then  $X \in g$  is unstable  $\Leftrightarrow$  ad X is nilpotent.

*Proof*: ( $\Rightarrow$ ) From the formula ad (Ad g(x)) = Ad  $g \circ$  ad  $x \circ$  Ad $g^{-1}$  it is immediate that the characteristic polynomial det (tI – ad x) is G-invariant, hence that is coefficients are invariant functions. If x is unstable, these all vanish so adx is nilpotent.

( $\Leftarrow$ ) If adx is nilpotent then the { exp  $t(x) | t \in k$  } is a unipotent subgroup of G which must be contained in the unipotent radical  $R_u(B)$  of some Borel subgroup B of G. Fix a maximal torus  $T \subset B$ , so  $B = R_u \cdot T$ . Then by the structure theorem of semi-simple groups we can write  $g = t + \left(\sum_{\alpha>0} g_{\alpha}\right)$  $+ \left(\sum_{\alpha<0} g_{\alpha}\right)$  where t = Lie(T) and  $\left(\sum_{\alpha>0} g_{\alpha}\right) = \text{Lie}(R_u(B))$ . Let  $\chi_{\alpha}$  be the character of T, which is associated to  $\alpha = (\alpha_i)$  (i.e. if  $w \in T$ ,  $y \in g_{\alpha}$ then Ad  $(w)(y) = \chi_{\alpha}(w) y$ ), and let *l* be a linear functional on the group of characters of T defining the given ordering: i.e.,

 $l(\chi_{\alpha}) = \sum_{i} c_{i} \alpha_{i} > 0$  if  $\alpha > 0$  and  $l(\chi_{\alpha}) < 0$  if  $\alpha < 0$ .

We can always choose l so that all the  $c_i$  are integers. If we define a 1-PS  $\lambda$ :  $\mathbf{G}_m \to T$  by  $\lambda(t) = (..., t^{c_i}, ...)$ , then the weights of X with respect to  $\lambda$  are some subset of  $\{l(\alpha) \mid \alpha > 0\}$ , hence are positive. Thus X is unstable.

REMARK. There are no stable points. One can show that the regular semi-simple elements of g have closed orbits of maximal dimension but their stabilizers will be their centralizers, i.e. maximal tori of G, and hence far from finite.

1.16. CHOW FORM. The Chow form is the answer to the problem of describing by an explicit set of numbers a general subvariety  $V^r \subset \mathbf{P}^n$ . In two cases, the problem has a very easy answer: a hypersurface has its equation F and a linear space  $L^r$  has its Plücker coordinates. The Chow form is just a clever combination of these two special cases. Suppose  $V^r$  has degree d. There are two ways to proceed

i) If  $u = (u_i) \in \mathbf{P}^n$  write  $H_u$  for the hyperplance  $\sum u_i X_i = 0$ . One shows that there is an irreducible polynomial  $\Phi_V$  such that

 $\left[V \cap H_u^{(0)} \cap \dots \cap H_u^{(r)} \neq \emptyset\right] \Leftrightarrow \left[\Phi_V(u_i^{(0)}, \dots, u_i^{(r)}) = 0\right]$ 

Moreover  $\Phi_V$  is multihomogeneous of degree d in each of the sets of variables  $(u_0^{(j)}, ..., u_n^{(j)}), \Phi_V$  is unique up to a scalar, and  $\Phi_V$  determines V.

ii) If G = Grassmanian of  $L^{n-r-1's}$  in  $\mathbb{P}^n$  and  $\mathcal{O}_G(1)$  is the ample line bundle on G defined by its Plücker embedding, then the set of  $L \in G$  such that  $L \cap V \neq \emptyset$  is the divisor  $D_V$  of zeroes of some section if  $\mathcal{O}_G(d)$  and Vand  $D_V$  determine each other. (Unfortunately,  $D_V$  is almost always a singular divisor.)

These methods give the same result via the identification:

$$\bigoplus_{d=0}^{\infty} \Gamma(G, \mathcal{O}_G(d)) = \begin{cases} \text{Homogeneous} \\ \text{coordinate} \\ \text{ring of } G \end{cases}$$
$$= \begin{cases} \text{Subring } W \text{ of } \mathbf{C} [..., U_i^{(j)}, ...] \text{ generated} \\ \text{by the Plücker coordinates} \\ P_{i_0, ..., i_r} = \det_{(r+1, r+1)} (U_{i_l}^{(j)}), i_0 < i_1 < ... < i_r \end{cases}$$

Letting  $W_d$  be the  $d^{th}$  graded piece of W, the identification furnishes an irreducible representation r+1

$$\operatorname{Symm}^{d}\left(\Lambda^{r+1}\left(\mathbf{C}^{n+1}\right)\right) \to W_{d} \subset W_{d} \subset \operatorname{Symm}^{d}\left(\mathbf{C}^{n+1}\right)$$

Thus, although we will usually consider the Chow form as a point of the SL(n+1) representation  $\otimes^{r+1}$  Symm<sup>d</sup> ( $\mathbb{C}^{n+1}$ ) this form lies in the irreducible piece  $W_d$  and can be thought of as defining a divisor on the Grassmanian. For more details on Chow forms, see Samuel [17, Ch. 1 § 9].

1.17. ASYMPTOTIC STABILITY. We will say that a variety  $V^r \subset \mathbf{P}^n$  is *Chow stable* or simply *stable* if its Chow form is stable for the natural SL(n+1)-action. If L is an ample line bundle on V, we say that (V, L) is *asymptotically stable* if

$$\exists n_0 \text{ s.t. } \forall n \ge n_0, \Phi_{\Gamma(L^n)}(V) \subset \mathbf{P}^{h^0(L^n)-1} \text{ is stable.}$$

Attention: a stable variety need not be asymptotically stable (nor, of course, vice versa). Indeed, one of the main goals of this exposition is to show that the asymptotically stable curves are exactly the "stable" curves of Deligne and Mumford, and that by using asymptotic stability we can construct  $\overline{\mathcal{M}}_g$  as a "quotient" moduli space for these curves.