

# 1. Introduction

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **23 (1977)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

# THE GELFAND-NAIMARK THEOREMS FOR C\*-ALGEBRAS

by Robert S. DORAN and Josef WICHMANN

## 1. INTRODUCTION

Many of the Banach spaces which attract attention are at the same time algebras under some multiplication. In spite of this fact their study from this richer point of view was taken up only after the publication in 1932 of Banach's book [6]. One of the early fundamental results in the general theory of Banach algebras was a generalization of the classical theorem of Frobenius that any finite dimensional division algebra over the complex field is isomorphic to the field of complex numbers. S. Mazur [35] announced in 1938 that every complex normed division algebra is isomorphic to the field of complex numbers. Since the first published proof was given by I. M. Gelfand [22] this result is often called the Mazur-Gelfand theorem [43], [55]. As an immediate consequence one obtains the following beautiful characterization of the complex field among normed algebras: any normed algebra satisfying the norm condition  $\|xy\| = \|x\| \cdot \|y\|$  for all elements  $x$  and  $y$  is isometrically isomorphic to the field of complex numbers.

Many important Banach algebras carry a natural involution. In the case of an algebra of functions the involution is the operation of taking the complex-conjugate and in the case of an algebra of operators on a Hilbert space it is the operation of taking the adjoint operator. Motivated by these observations the Soviet mathematicians Israel M. Gelfand and Mark A. Naimark [23] proved, under some additional assumptions, the following two theorems:

**THEOREM I.** *Let  $A$  be a commutative Banach algebra with involution satisfying  $\|x^*x\| = \|x^*\| \cdot \|x\|$  for all  $x$  in  $A$ . Then  $A$  is isometrically \*-isomorphic to  $C_0(X)$ , the algebra of all continuous complex-valued functions which vanish at infinity on some locally compact Hausdorff space  $X$ .*

**THEOREM II.** *Let  $A$  be a Banach algebra with involution satisfying  $\|x^*x\| = \|x^*\| \cdot \|x\|$  for all  $x$  in  $A$ . Then  $A$  is isometrically \*-iso-*

morphic to a norm-closed \*-subalgebra of bounded linear operators on some Hilbert space.

The purpose of this paper is to present a thorough discussion of these two representation theorems. We shall trace, as carefully as we have been able, the interesting and rather tangled history which led to their present form. Then proofs of the theorems will be given. Finally, we shall survey some recent developments inspired by the theorems.

## 2. DEFINITIONS AND MOTIVATION

A *\*-algebra* is a complex associative linear algebra  $A$  with a mapping  $x \rightarrow x^*$  of  $A$  into itself such that for all  $x, y \in A$  and complex  $\lambda$ : (a)  $x^{**} = x$ ; (b)  $(\lambda x)^* = \bar{\lambda}x^*$ ; (c)  $(x + y)^* = x^* + y^*$ ; and (d)  $(xy)^* = y^*x^*$ . The map  $x \rightarrow x^*$  is called an *involution*; because of (a) it is clearly bijective. A subalgebra  $B$  of  $A$  is called a *\*-subalgebra* if  $x \in B$  implies  $x^* \in B$ .

An algebra which is also a Banach space satisfying  $\| xy \| \leq \| x \| \cdot \| y \|$  for all  $x$  and  $y$  is called a *Banach algebra*. A Banach algebra which is also a *\*-algebra* is called a *Banach \*-algebra*. The involution in a Banach *\*-algebra* is said to be *continuous* if there is a constant  $M$  such that  $\| x^* \| \leq M \| x \|$  for all  $x$ ; the involution is *isometric* if  $\| x^* \| = \| x \|$  for all  $x$ .

A norm on a *\*-algebra* is said to satisfy the *B\*-condition* if  $\| x^*x \| = \| x^* \| \cdot \| x \|$  for all  $x$ ; a *B\*-algebra* is a Banach *\*-algebra* whose norm satisfies the *B\*-condition*. A *B\*-algebra* with isometric involution clearly satisfies the condition  $\| x^*x \| = \| x \|^2$ . On the other hand, if  $A$  is a Banach *\*-algebra* satisfying  $\| x \|^2 \leq \| x^*x \|$  (in particular if equality holds), then  $A$  is easily seen to be a *B\*-algebra* with isometric involution.

The Banach space  $C(X)$  of continuous complex-valued functions on a compact Hausdorff space is a commutative *B\*-algebra* under point-wise multiplication  $(fg)(t) = f(t)g(t)$ , involution  $f^*(t) = \overline{f(t)}$ , and sup-norm. Similarly, the algebra  $C_0(X)$  of continuous complex-valued functions which vanish at infinity on a locally compact Hausdorff space is a commutative *B\*-algebra*.

Examples of noncommutative *B\*-algebras* are provided by the algebra  $B(H)$  of bounded linear operators on a Hilbert space  $H$ . Multiplication in  $B(H)$  is operator composition, the involution  $T \rightarrow T^*$  is the usual adjoint operation, and the norm is the operator norm  $\| T \| = \sup \{ \| T\xi \| : \|\xi\| \leq 1, \xi \in H \}$ . A norm-closed *\*-subalgebra* of  $B(H)$  is called a *C\*-algebra*; clearly, every *C\*-algebra* is a *B\*-algebra*.