

5. The register theorem

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **23 (1977)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

The Whitehead product theory for ex-spaces has been worked out by Eggar [4]. His definition is such that if A, B, Y are as in §2 and $\alpha \in \pi_G(\Sigma A, Y)$, $\beta \in \pi_G(\Sigma B, Y)$ then

$$(4.1) \quad [P_{\#}\alpha, P_{\#}\beta] = P_{\#}[\alpha, \beta]$$

in $\pi_X(\Sigma(P_{\#}A \wedge P_{\#}B), P_{\#}Y)$. Since we shall only be concerned with elements in the image of $P_{\#}$ we can introduce (4.1) as a piece of notation, without going into the details of Eggar's theory.

5. THE REGISTER THEOREM

In this section we suppose that X is a finite simply-connected CW-complex, although the results obtained can no doubt be generalized. We define the *register* $\text{reg}(X)$ of X to be the number of positive integers r such that, for some abelian group A , the cohomology group $H^r(X; A)$ is non-trivial. If X is a sphere, for example, then $\text{reg}(X) = 1$.

Let $p: M \rightarrow X$ be a fibration with fibre N . If a cross-section $s: X \rightarrow M$ exists then $sp: M \rightarrow M$ is a fibre-preserving map which is constant on the fibre. Conversely if $k: M \rightarrow M$ is a fibre-preserving map which is nullhomotopic on the fibre then M admits a cross-section as shown by Noakes [11]. We use similar arguments to prove

THEOREM (5.1). *Let $k: M \rightarrow M$ be a fibre-preserving map such that $l: N \rightarrow N$ is nullhomotopic, where $l = k|N$, and let $s, t: X \rightarrow M$ be cross-sections. Then $k^r s$ and $k^r t$ are vertically homotopic, where $r = \text{reg}(X)$.*

The n -section ($n = 0, 1, \dots$) of the complex X is denoted by X^n . Since X is connected we have a vertical homotopy of s into t over X^0 . This starts an induction. Suppose that for some $n \geq 1$ and some $q = q(n) \geq 1$ we have a vertical homotopy of $k^q s$ into $k^q t$ over X^{n-1} , so that the separation class

$$d = d(k^q s, k^q t) \in H^n(X; \pi_n(N))$$

is defined. If the cohomology group vanishes then $d = 0$ and $k^q s \simeq k^q t$ over X^n . But in any case the induced endomorphism l_* of $\pi_n(N)$ is trivial, by hypothesis, and so d lies in the kernel of the coefficient endomorphism $l_{\#}$ determined by l_* . Therefore

$$d(k^{q+1} s, k^{q+1} t) = l_{\#} d = 0,$$

and so $k^{q+1}s \simeq k^{q+1}t$ over X^n . Hence, by induction, we obtain (5.1). Of course the value of r can often be improved in particular cases.

COROLLARY (5.2). *Let E be a fibre bundle over X with locally compact fibre F , which admits a cross-section. Choose a cross-section and so regard E as an ex-space. Let $f: E \rightarrow E$ be an ex-map such that $g: F \rightarrow F$ is null-homotopic, where $g = f|_F$. Then $f^{r+1} \simeq c$, the trivial ex-map, where $r = \text{reg}(X)$.*

To see this, take $M = M_X(E, E)$, in (5.1), and define $k: M \rightarrow M$ by post-composition with f . We take s, t to be the cross-sections $f', e': X \rightarrow M$ determined by f, e , and obtain (5.2).

Now let $\alpha, \beta \in \pi_X(E, E)$ be elements such that

- (i) $\alpha^2 = \beta^2$ and $\alpha\beta = \beta\alpha$,
- (ii) $\Phi_*\alpha = \Phi_*\beta$,

where $\Phi_*: \pi_X(E, E) \rightarrow \pi(F, F)$ is given by restriction. Suppose that $E = \Sigma E'$, for some ex-space E' , and that $\alpha = \Sigma_*\alpha'$, $\beta = \Sigma_*\beta'$, for some $\alpha' \in \pi_{X'}(E', E')$. Take f in (5.2) to be a representative of $\alpha - \beta$. Then f^{r+1} is a representative of $(\alpha - \beta)^{r+1} = 2^r(\alpha - \beta)\alpha^r$, and so (5.2) shows that

$$(5.3) \quad 2^r\alpha = 2^r\beta.$$

Applications will be given in §8 below.

6. THE EXACT SEQUENCE

Let X be a CW-complex with basepoint x_0 a 0-cell. Let $p: M \rightarrow X$ be a fibration with fibre $N = p^{-1}(x_0)$, and let Γ denote the function-space of cross-sections. By evaluating at x_0 we obtain a fibration $q: \Gamma \rightarrow N$. It may be noted that, under fairly general conditions, this fibration admits a cross-section if and only if the original fibration is trivial, in the sense of fibre homotopy type.

Now choose a basepoint $y_0 \in N$ so that $q^{-1}(y_0) = \Gamma_0$, the space of pointed cross-sections. Choose such a cross-section s as basepoint in $\Gamma_0 \subset \Gamma$, and consider the homotopy exact sequence of the fibration as follows:

$$\dots \rightarrow \pi_{r+1}(N) \xrightarrow{\Delta} \pi_r(\Gamma_0) \xrightarrow{u_*} \pi_r(\Gamma) \xrightarrow{q_*} \pi_r(N) \rightarrow \dots$$

Note that Γ_0 is a deformation retract of $\tilde{\Gamma}_0$, the space of pointed maps $t: X \rightarrow M$ such that $pt \simeq 1$.