

# 5. SCHWARZIAN DERIVATIVE AND UNIVALENCE

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **24 (1978)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **24.05.2024**

## **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

$> \sigma_1 - \varepsilon$ . If  $w$  is defined by  $w(z) = z + e^{i\theta}/z$  and  $f = w \circ h^{-1}$ , then  $f$  is univalent in  $A$  and

$$\|S_f\|_A = \|S_w - S_h\|_D \geq |S_w(0) - S_h(0)| = |6e^{i\theta} + S_h(0)|.$$

By choosing  $\varphi$  suitably we obtain  $\|S_f\|_A > 6 + \sigma_1 - \varepsilon$ .

## 5. SCHWARZIAN DERIVATIVE AND UNIVALENCE

5.1 *Constant  $\sigma_3$ .* Let  $A$  again be a simply connected domain with more than one boundary point. As a kind of opposite to the constant  $\sigma_2$  we define

$$\sigma_3 = \sup \{a \mid \|S_f\| \leq a \text{ implies } f \text{ univalent in } A\}.$$

Note that the number  $a = 0$  is always in the above set. In this definition, sup can be replaced by max, as can be shown by a standard normal family argument.

Nehari [12] proved that in a disc, the condition  $\|S_f\| \leq 2$  implies the univalence of  $f$ , and Hille [5] showed that the bound 2 is best possible. In other words,  $\sigma_3 = 2$  for a disc.

A closer study of  $\sigma_3$  leads to the universal Teichmüller space and reveals an intrinsic significance of quasiconformal mappings in the theory of univalent functions. The gist is the following result.

**THEOREM 5.1.** *The constant  $\sigma_3$  is positive if and only if  $A$  is bounded by a quasicircle.*

*Proof:* The sufficiency of the condition was established by Ahlfors [1] who actually proved more: If  $A$  is bounded by a  $K$ -quasicircle, there is an  $\varepsilon > 0$  depending only on  $K$ , such that whenever  $\|S_f\|_A < \varepsilon$ , then  $f$  is univalent and can be continued to a quasiconformal mapping of the plane. In the proof, the extension of the given meromorphic  $f$  is explicitly constructed by means of a continuously differentiable quasiconformal reflection  $\varphi$  in  $\partial A$  with bounded  $|d\varphi|/|dz|$  (cf. 3.3).

The necessity was proved by Gehring [2]. His proof was in two steps. It was first shown, by aid of an example, that if  $A$  is not  $b$ -locally connected for any  $b$ , then  $\sigma_3 = 0$ . After this, the desired conclusion was drawn from the result we stated above as Lemma 3.2.

5.2 *Universal Teichmüller space.* Henceforth, we assume that the domain  $A$  is bounded by a quasicircle. Let  $Q(A)$  be the Banach space

consisting of all holomorphic functions  $\varphi$  of  $A$  with finite norm. We introduce the subsets

$$U(A) = \{\varphi = S_f \mid f \text{ univalent in } A\},$$

$$T(A) = \{S_f \in U(A) \mid f \text{ can be extended to a quasiconformal mapping of the plane}\}.$$

Both sets are well defined. The set  $T(A)$  is called the *universal Teichmüller space* of  $A$ .

**THEOREM 5.2.** *The sets  $T(A)$  and  $U(A)$  are connected by the relation  $T(A) = \text{interior of } U(A)$ .*

*Proof:* We first show that  $T(A)$  is open. Choose  $S_f \in T(A)$ ,  $S_h \in Q(A)$ , and set  $g = h \circ f^{-1}$ . Then  $g$  is meromorphic in the domain  $f(A)$ . Since  $\partial A$  is a quasicircle,  $\partial f(A)$  is also a quasicircle. By the theorem of Ahlfors cited in the proof of Theorem 5.1, there is an  $\varepsilon > 0$  such that if

$$(5.1) \quad \|S_g\|_{f(A)} < \varepsilon,$$

then  $S_g \in T(f(A))$ . Now, choose  $h$  so that  $\|S_f - S_h\|_A < \varepsilon$ . Then (5.1) holds, and it follows that  $S_h = S_{g \circ f} \in T(A)$ .

After this it suffices to prove that  $\text{int } U(A) \subset T(A)$ . Choose  $S_f \in \text{int } U(A)$  and then an  $\varepsilon > 0$ , so that the ball  $B = \{\varphi \in Q(A) \mid \|\varphi - S_f\| < \varepsilon\}$  is contained in  $U(A)$ . Let  $g$  be an arbitrary meromorphic function in  $f(A)$  for which  $\|S_g\|_{f(A)} < \varepsilon$ . If  $h = g \circ f$ , then  $\|S_f - S_h\|_A = \|S_g\|_{f(A)} < \varepsilon$ . Thus  $S_h \in U(A)$ . But then also  $g = h \circ f^{-1}$  is univalent, and we have proved that  $\sigma_3$  is positive for the domain  $f(A)$ . By Theorem 5.1, the boundary  $\partial f(A)$  is a quasicircle. Hence, by the remark in 3.3,  $S_f \in T(A)$ .

**COROLLARY 5.1.** *If  $f$  is univalent in  $A$  and  $\|S_f\|_A < \sigma_3$ , then  $f$  can be extended to a quasiconformal mapping of the plane.*

*Proof:* This follows immediately from Theorem 5.2, in view of our previous remark that the closed ball  $\{\varphi \in Q(A) \mid \|\varphi\|_A \leq \sigma_3\}$  is contained in  $U(A)$ .

By this Corollary, we have for  $A$ ,

$$\sigma_3 = \sup \{a \mid \|S_f\|_A < a \text{ implies that } f \text{ is univalent and can be extended to a quasiconformal mapping of the plane}\}.$$

5.3 *New characterization for  $\sigma_3$ .* Theorem 5.2 was proved by Gehring [2] in the case where  $A$  is a half-plane. As is seen from the above proof, the generalization for an arbitrary  $A$  is immediate. In fact, the sets  $Q(A)$ ,  $U(A)$  and  $T(A)$  corresponding to different domains  $A$  are isomorphic:

LEMMA 5.1. *Let  $h$  be a conformal mapping of the upper half-plane  $H$  onto  $A$ . Then the mapping  $h^*: Q(A) \rightarrow Q(H)$ , defined by  $h^*(S_f) = S_{f \circ h}$ , is a bijective isometry. It maps  $U(A)$  and  $T(A)$  onto  $U(H)$  and  $T(H)$ , respectively.*

*Proof:* Clearly  $h^*$  is well defined and a bijection of  $Q(A)$ ,  $U(A)$  and  $T(A)$  onto  $Q(H)$ ,  $U(H)$  and  $T(H)$ , respectively. That  $h^*$  is an isometry follows from formula (4.3).

The function  $h^*$  maps the origin of  $Q(A)$  onto the point  $S_h \in T(H)$ , which has the distance  $\sigma_1$  from the origin of  $Q(H)$ . If  $B = \{\varphi \in Q(A) \mid \|\varphi\|_A \leq \sigma_3\}$ , then

$$h^*(B) = \{\psi \in Q(H) \mid \|\psi - S_h\| \leq \sigma_3\}.$$

From this and the definition of  $\sigma_3$  we infer that  $\sigma_3$  is equal to the distance from the point  $S_h$  to the boundary of  $U(H)$ . The following characterization seems to be more useful:

LEMMA 5.2. *The constant  $\sigma_3$  of  $A$  is equal to the distance of the point  $S_h$  to the boundary of  $T(H)$ .*

*Proof:* Let  $d$  denote the distance function in  $Q$ . Since  $T(H) \subset U(H)$  we conclude from what we just said above that  $\sigma_3 \geq d(\{S_h\}, U(H) - T(H))$ . On the other hand, it follows from Theorem 5.2 that  $\text{int } B \subset T(A)$  and hence  $\text{int } h^*(B) \subset T(H)$ . Therefore,  $\sigma_3 \leq d(\{S_h\}, U(H) - T(H))$ .

A standard normal family argument shows that  $U(A)$  is a closed subset of  $Q(A)$ . Therefore, the closure of  $T(A)$  is contained in  $U(A)$ . Gehring [3] showed recently that this inclusion is proper, thus disproving a famous conjecture of Bers.

However, it is true that on every sphere  $\|\varphi\| = r$  of  $Q(H)$ ,  $2 \leq r \leq 6$ , there are points of  $U(H) - T(H)$  which belong to the closure of  $T(H)$  ([9]).

5.4 *Estimates for  $\sigma_3$ .* Lemma 5.2 can be used to deriving estimates for  $\sigma_3$  in terms of  $\sigma_1$  ([9]). Suppose first that  $0 \leq \sigma_1 < 2$ . Then  $S_h$  lies in the ball  $\{\varphi \in Q(H) \mid \|\varphi\| < 2\}$  which is a subset of  $T(H)$ . Since  $\|S_h\| = \sigma_1$ ,

we conclude that  $d(\{S_h\}, U(H) - T(H)) \geq 2 - \sigma_1$ . Consequently, by Lemma 5.2,

$$(5.2) \quad \sigma_3 \geq 2 - \sigma_1.$$

In order to prove that this inequality is sharp, we consider the point  $S_w$ , where  $w$  is the restriction to  $H$  of a branch of the logarithm. Since the boundary of  $w(H)$  is not a quasicircle,  $S_w \in U(H) - T(H)$ . From  $S_w(z) = z^{-2}/2$  it follows that  $\|S_w\|_H = 2$ . Let  $h$  be determined by the condition  $S_h = r S_w$ ,  $0 < r < 1$ , and set  $A = h(H)$ . From  $\|S_h\|_H < 2$  it follows that  $S_h \in T(H)$ , and so  $\partial A$  is a quasicircle. Now

$$\sigma_3 = d(\{S_h\}, U(H) - T(H)) = \|S_w - S_h\| = 2(1-r) = 2 - \sigma_1,$$

showing that (5.2) is sharp.

Suppose that  $2 \leq \sigma_1 < 6$ . We then conclude from the remark at the end of 5.3 that, even though  $\sigma_3 > 0$  for each  $A$ , we have  $\inf \sigma_3 = 0$  for every  $\sigma_1$ .

Similarly, Lemma 5.2 can be used to deriving the upper estimate

$$\sigma_3 \leq \min(2, 6 - \sigma_1).$$

(For the details we refer to [9].)

#### REFERENCES

- [1] AHLFORS, L. V. Quasiconformal reflections. *Acta Math.* 109 (1963), pp. 291-301.
- [2] GEHRING, F. W. Univalent functions and the Schwarzian derivative. *Comment. Math. Helv.* 52 (1977), pp. 561-572.
- [3] ——— Spirals and the universal Teichmüller space. *To appear in Acta Math.*
- [4] GEHRING, F. W. and J. VÄISÄLÄ. Hausdorff dimension and quasiconformal mappings. *J. London Math. Soc.* (1973).
- [5] HILLE, E. Remarks on a paper by Zeev Nehari. *Bull. Amer. Math. Soc.* 55 (1949), pp. 552-553.
- [6] KRAUS, W. Über den Zusammenhang einiger Charakteristiken eines einfach zusammenhängenden Bereichs mit der Kreisabbildung. *Mitt. math. Semin. Giessen* 21 (1932).
- [7] KÜHNAU, R. Wertannahmeprobleme bei quasikonformen Abbildungen mit ortsabhängiger Dilatationsbeschränkung. *Math. Nachr.* 40 (1969).
- [8] LEHTO, O. Schlicht functions with a quasiconformal extension. *Ann. Acad. Sci. Fenn. A I* 500 (1971).
- [9] ——— Domain constants associated with Schwarzian derivative. *Comment. Math. Helv.* 52 (1977), pp. 603-610.
- [10] ——— Univalent functions and Teichmüller theory. *Proc. of the First Finnish-Polish Summer School in Complex Analysis at Podlesice, University of Lodz* (1977), pp. 11-33.