

III. A VARIANT

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **27 (1981)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **23.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

III. A VARIANT

Let us agree to call a scheme S *accessible* if there exists an absolutely finitely generated field K for which the set $S(K)$ of K -valued points of S is non-empty. Thus for example, if K is an absolutely finitely generated field, then for *any* subring $R \subset K$, $\text{Spec}(R)$ is accessible (by the K -valued point $R \hookrightarrow K$); also any subring R' of the power-series ring $K[[X_1, \dots, \dots]]$ over K in any number of variables has $\text{Spec}(R')$ accessible

$$(\text{by } R' \hookrightarrow K \xrightarrow{X \rightarrow 0} K).$$

On the other hand, the spectrum of a field F is accessible if and only if F is absolutely finitely generated.

THEOREM 2. *Let S be a connected, locally noetherian scheme which is accessible. Let X/S be a proper and smooth S -scheme with geometrically connected fibres. Then the group $\text{Ker}(X/S)$ is finite.*

Proof. We begin by reducing to the case when S is a finitely generated field. In view of the accessibility of S , this reduction results from the following simple lemma applied with $T = \text{Spec}(K)$.

LEMMA 4. *Let X/S be proper and smooth with geometrically connected fibres over a connected locally noetherian scheme S . Given a connected locally noetherian S -scheme T , denote by X_T/T the inverse image of X/S on T , i.e. form the cartesian diagram*

$$\begin{array}{ccc} & X_T = X \times_S T & \\ \swarrow & \downarrow & \\ X & & T \\ \downarrow & \swarrow & \\ S & & \end{array}$$

The natural map (cf. 1.5)

$$\text{Ker } (X_T/T) \rightarrow \text{Ker } (X/S)$$

is surjective.

Proof. Let t be a geometric point of T , s the image geometric point of S , and x a geometric point on the fibre X_s . The homotopy exact sequences (SGA I, Exp X, 1.4) for X/S and X_T/T sit in a commutative diagram

$$\begin{array}{ccccccc} \pi_1(X_s, x) & \longrightarrow & \pi_1(X_T, x) & \longrightarrow & \pi_1(T, t) & \longrightarrow & 0 \\ \parallel & & \downarrow & & \downarrow & & \\ \pi_1(X_s, x) & \longrightarrow & \pi_1(X, x) & \longrightarrow & \pi_1(S, s) & \longrightarrow & 0 \end{array}$$

Passing to the abelianizations yields the commutative diagram with exact rows

$$\begin{array}{ccccccc} \pi_1(X_s)^{ab} & \longrightarrow & \pi_1(X_T)^{ab} & \longrightarrow & \pi_1(T)^{ab} & \longrightarrow & 0 \\ \parallel & & \downarrow & & \downarrow & & \\ \pi_1(X_s)^{ab} & \longrightarrow & \pi_1(X)^{ab} & \longrightarrow & \pi_1(S)^{ab} & \longrightarrow & 0 \end{array}$$

whence we find

$$\begin{array}{l} \pi_1(X_s)^{ab} \begin{cases} \nearrow \text{Ker } (X_T/T) = \text{image of } \pi_1(X_s)^{ab} \text{ in } \pi_1(X_T)^{ab} . \\ \searrow \text{Ker } (X/S) = \text{image of } \pi_1(X_s)^{ab} \text{ in } \pi_1(X)^{ab} . \end{cases} \end{array} \quad \text{QED}$$

Thus we are reduced to proving the finiteness of $\text{Ker } (X/K)$ when K is an absolutely finitely generated field, and X/K is proper, smooth, and geometrically connected. We have already proven this finiteness theorem when X/K is an abelian variety (cf. Remark (1) above). We will reduce to this case by making use of the theory of the Picard and Albanese varieties.

At the expense of replacing K by a finite extension, we may assume that X has a K -rational point x_0 . The Picard scheme $\text{Pic}_{X/K}$ is then a commutative group-scheme locally of finite type over K , which represents the functor on $\{\text{Schemes}/K\}$

$$W \rightarrow \left\{ \begin{array}{l} \text{the group of } W\text{-isomorphism classes of pairs } (\mathcal{L}, \varepsilon) \text{ consisting} \\ \text{of an invertible sheaf } \mathcal{L} \text{ on } X \times_K W \text{ together with a} \\ \text{trivialization } \varepsilon \text{ of the restriction } \mathcal{L} \text{ to } \{x_0\} \times_K W \end{array} \right.$$

The subgroup-scheme $Pic_{X/K}^\tau$ of $Pic_{X/K}$ classifies those $(\mathcal{L}, \varepsilon)$ whose underlying \mathcal{L} becomes τ -equivariant to zero when restricted to every geometric fibre of $X \times W/W$ (i.e. for each geometric point w of W , some multiple of $\mathcal{L} \mid X \times w$ is algebraically equivalent to zero). The identity component $Pic_{X/K}^0$ of $Pic_{X/K}$ classifies those $(\mathcal{L}, \varepsilon)$ whose \mathcal{L} becomes algebraically equivalent to zero on each geometric fibre $X \times W/W$. The Picard variety $Pic_{X/K}^{0, \text{red}}$ is an abelian variety over K , and it sits in an *f.p.f.* short exact sequence of commutative group schemes

$$(3.1) \quad 0 \rightarrow Pic_{X/K}^{0, \text{red}} \rightarrow Pic_{X/K}^\tau \rightarrow C \rightarrow 0$$

in which the cokernel C is a finite flat group-scheme over K . This cokernel C should be thought of as the “scheme theoretic” torsion in the Neron-Severi group.

We denote by $Alb_{X/K}$ the Albanese variety of X/K , defined to be the dual abelian variety to the Picard variety $Pic_{X/K}^{0, \text{red}}$. We now recall the expression of $\pi_1(X \otimes \bar{K})^{ab}$ in terms of the Tate module of the Albanese, and a finite “error term” involving the Cartier dual C^\vee of C .

LEMMA 5. *Let K be a field, and X/K a proper, smooth and geometrically connected K -scheme which admits a K -rational point. Then there is a canonical short exact sequence of $\text{Gal}(\bar{K}/K)$ -modules*

$$(3.2) \quad 0 \rightarrow C^\vee(\bar{K}) \rightarrow \pi_1(X \otimes \bar{K})^{ab} \rightarrow T(Alb_{X/K}(\bar{K})) \rightarrow 0.$$

Proof. By Kummer and Artin-Schreier theory, we have for each integer $N \geq 1$ a canonical isomorphism

$$\begin{aligned} & \text{Hom}(\pi_1(X \otimes \bar{K})^{ab}, \mathbf{Z}/N\mathbf{Z}) \\ &= H_{et}^1(X \otimes \bar{K}, \mathbf{Z}/N\mathbf{Z}) \simeq \text{Hom}(\mu_N, (Pic_{X/K}^\tau \otimes \bar{K})). \end{aligned}$$

in which the last Hom is in the sense of \bar{K} -group-schemes. Applying the functor $X \mapsto \text{Hom}(\mu_N, X)$ to the short exact sequence

$$0 \rightarrow Pic^{0, \text{red}} \rightarrow Pic^\tau \rightarrow C \rightarrow 0$$

gives a short exact sequence

$$(3.3) \quad \begin{aligned} 0 &\rightarrow \text{Hom}(\mu_N, (Pic^{0, \text{red}} \otimes \bar{K})) \\ &\rightarrow \text{Hom}(\mu_N, (Pic^\tau \otimes \bar{K})) \rightarrow \text{Hom}(\mu_N, C \otimes \bar{K}) \rightarrow 0 \end{aligned}$$

(the final zero because over an algebraically closed field, the group $\text{Ext}^1(\mu_N, A)$ vanishes for any abelian variety A , cf. the remark at the end of this section). We now “decode” its two end terms, using Cartier-Nishi duality for the first, and Cartier duality for the last.

The first is

$$\begin{aligned}
 \text{Hom}(\mu_N, (Pic^{0, \text{red}}) \otimes \overline{K}) &= \text{Hom}(\mu_N, (Pic^{0, \text{red}})_N \otimes \overline{K}) \\
 &\quad \Downarrow \text{Cartier-Nishi duality} \\
 &\text{Hom}(\text{Alb}_{X/N})_N \otimes \overline{K}, \mathbf{Z}/N\mathbf{Z}) \\
 &\quad \Downarrow \text{evaluation on } \overline{K}\text{-points} \\
 &\text{Hom}((\text{Alb}_{X/K}(\overline{K}))_N, \mathbf{Z}/N\mathbf{Z}) \\
 &\quad \Downarrow \\
 &\text{Hom}(T(\text{Alb}_{X/K}(\overline{K})), \mathbf{Z}/N\mathbf{Z}).
 \end{aligned}$$

The last is

$$\begin{aligned}
 \text{Hom}(\mu_N, C \otimes \overline{K}) &\xrightarrow{\text{Cartier duality}} \text{Hom}(C^\vee \otimes \overline{K}, \mathbf{Z}/N\mathbf{Z}) \\
 &\quad \downarrow \int \text{evaluation} \\
 &\text{Hom}(C^\vee(\overline{K}), \mathbf{Z}/N\mathbf{Z})
 \end{aligned}$$

“Substituting” into the exact sequence (3.2), we find a canonical short exact sequence

$$\begin{aligned}
 (3.4) \quad &0 \rightarrow \text{Hom}(T(\text{Alb}_{X/K}(\overline{K})), \mathbf{Z}/N\mathbf{Z}) \\
 &\rightarrow \text{Hom}(\pi_1(X \otimes \overline{K})^{ab}, \mathbf{Z}/N\mathbf{Z}) \rightarrow \text{Hom}(C^\vee(\overline{K}), \mathbf{Z}/N\mathbf{Z}) \rightarrow 0
 \end{aligned}$$

Passing to the *direct* limit as N grows multiplicatively, we obtain a canonical short exact sequence

$$\begin{aligned}
 (3.5) \quad &0 \rightarrow \text{Hom}(T(\text{Alb}_X(\overline{K})), \mathbf{Q}/\mathbf{Z}) \\
 &\rightarrow \text{Hom}(\pi_1(X \otimes \overline{K})^{ab}, \mathbf{Q}/\mathbf{Z}) \rightarrow \text{Hom}(C^\vee(\overline{K}), \mathbf{Q}/\mathbf{Z}) \rightarrow 0.
 \end{aligned}$$

Taking its Pontryagin dual, we find the required exact sequence (3.2). QED

To complete the reduction of Theorem 2 to the case of abelian varieties, we simply notice that the exact sequence of lemma 5 yields, upon passage to coinvariants, an exact sequence

$$(3.6) \quad (C^\vee(\bar{K}))_{\text{Gal}(\bar{K}/K)} \rightarrow \text{Ker}(X/K) \rightarrow \text{Ker}(\text{Alb}_{X/K}/K) \rightarrow 0$$

whose first term, being a quotient of the finite group $C^\vee(\bar{K})$, is finite. QED

Remark. In the course of the proof of Lemma 5, we appealed to the “well-known” vanishing of $\text{Ext}^1(\mu_N, A)$ over an algebraically closed field, for an abelian variety A and any integer $N > 1$. Here is a simple proof. It is enough to prove this vanishing when N is either prime to the characteristic p of K , or, in case $p > 0$, when $N = p$.

Suppose first N prime to p . Because the ground-field is algebraically closed, we have $\mu_N \simeq \mathbf{Z}/N\mathbf{Z}$, so it is equivalent to prove the vanishing of $\text{Ext}^1(\mathbf{Z}/N\mathbf{Z}, A)$. We will prove that *this* group vanishes for every integer $N > 1$. Consider such an extension:

$$0 \rightarrow A \rightarrow E \rightarrow \mathbf{Z}/N\mathbf{Z} \rightarrow 0$$

Pass to \bar{K} -valued points

$$0 \rightarrow A(\bar{K}) \rightarrow E(\bar{K}) \rightarrow \mathbf{Z}/N\mathbf{Z} \rightarrow 0$$

and consider the endomorphism “multiplication by N ”. Because the group $A(\bar{K})$ is N -divisible, the snake lemma gives an exact sequence

$$0 \rightarrow A(\bar{K})_N \rightarrow E(\bar{K})_N \rightarrow \mathbf{Z}/N\mathbf{Z} \rightarrow 0$$

But a point in $E(\bar{K})_N$ which maps onto “1” $\in \mathbf{Z}/N\mathbf{Z}$ is precisely a splitting of our extension.

Next consider the case $N = p = \text{char}(K)$. We give a proof due to Barry Mazur. Using the *f.p.p.f.* exact sequence

$$0 \rightarrow A_p \rightarrow A \rightarrow A \rightarrow 0.$$

to compute $\text{Ext}(\mu_p, -)$, we obtain a short exact sequence

$$0 \rightarrow \text{Hom}(\mu_p, A) \rightarrow \text{Ext}^1(\mu_p, A_p) \rightarrow \text{Ext}^1(\mu_p, A) \rightarrow 0$$

To prove that $\text{Ext}^1(\mu_p, A) = 0$, we will show that the groups $\text{Hom}(\mu_p, A)$ and $\text{Ext}^1(\mu_p, A_p)$ are both finite, of the same order. Trivially, we have $\text{Hom}(\mu_p, A) = \text{Hom}(\mu_p, A_p)$. Because we are over an algebraically closed field, and A_p is killed by p , its toroidal biconnected-etale decomposition looks like

$$A_p \simeq (\mu_p)^a \times (\text{biconnected}) \times (\mathbf{Z}/p\mathbf{Z})^b; \quad [\text{in fact } a = b].$$

Only the μ_p 's in A_p can “interact” with μ_p . Thus we are reduced to showing that $\text{Hom}(\mu_p, (\mu_p)^a)$ and $\text{Ext}^1(\mu_p, (\mu_p)^a)$ are both finite of the same cardinality p^a .

By Cartier duality, it is equivalent to show that both $\text{Hom}(\mathbb{Z}/p\mathbb{Z}, \mathbb{Z}/p\mathbb{Z})$ and $\text{Ext}^1(\mathbb{Z}/p\mathbb{Z}, \mathbb{Z}/p\mathbb{Z})$ have order p , and this is obvious (resolve the “first” $\mathbb{Z}/p\mathbb{Z}$ by

$$0 \rightarrow \mathbb{Z} \xrightarrow{p} \mathbb{Z} \rightarrow \mathbb{Z}/p\mathbb{Z} \rightarrow 0).$$

For another proof in this case, cf. Oort, [10], 85.

IV. ABSOLUTE FINITENESS THEOREMS

THEOREM 3. *Let \mathcal{O} be the ring of integers in a finite extension K of \mathbb{Q} . Let X be a smooth \mathcal{O} -scheme of finite type whose geometric generic fibre $X \otimes_{\mathcal{O}} \overline{K}$ is connected, and which maps surjectively to $\text{Spec}(\mathcal{O})$ (i.e. for every prime \mathfrak{p} of \mathcal{O} , the fibre over \mathfrak{p} , $X \otimes_{\mathcal{O}} (\mathcal{O}/\mathfrak{p})$, is non empty). Then the group $\pi_1(X)^{ab}$ is finite.*

Proof. This follows immediately from Theorem 1 and global classfield theory, according to which $\pi_1(\text{Spec}(\mathcal{O}))^{ab}$, the galois group of the maximal unramified abelian extension of K , is finite. QED

THEOREM 4. *Let \mathcal{O} be the ring of integers in a finite extension K of \mathbb{Q} , $\mathfrak{p}_1, \dots, \mathfrak{p}_n$ a finite set of primes of \mathcal{O} , $N = p_1 \dots p_n$ the product of their residue characteristics, and $\mathcal{O}[1/\mathfrak{p}_1 \dots \mathfrak{p}_n]$ the ring of “integers outside $\mathfrak{p}_1, \dots, \mathfrak{p}_n$ ” in K . Let X be a smooth $\mathcal{O}[1/\mathfrak{p}_1 \dots \mathfrak{p}_n]$ -scheme of finite type, whose geometric generic fibre $X \otimes_{\mathcal{O}} \overline{K}$ is connected, and which maps surjectively to $\text{Spec}(\mathcal{O}[1/\mathfrak{p}_1 \dots \mathfrak{p}_n])$ (i.e. for every prime $\mathfrak{p} \notin \{\mathfrak{p}_1, \dots, \mathfrak{p}_n\}$, the fibre*

$$X \otimes_{\mathcal{O}} (\mathcal{O}/\mathfrak{p})$$

is non-empty). Then the group $\pi_1(X)^{ab}$ is the product of a finite group and a pro- N group.

Proof. Again an immediate consequence of Theorem 1 and global classfield theory, according to which $\pi_1(\text{Spec}(\mathcal{O}[1/\mathfrak{p}_1 \dots \mathfrak{p}_n]))^{ab}$, the galois group of the maximal abelian, unramified outside $\{\mathfrak{p}_1, \dots, \mathfrak{p}_n\}$ -extension of K is finite times pro- N . QED