

V. Application to ℓ -adic representations

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **27 (1981)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Ein Dienst der *ETH-Bibliothek*

ETH Zürich, Rämistrasse 101, 8092 Zürich, Schweiz, www.library.ethz.ch

<http://www.e-periodica.ch>

THEOREM 5. *Let S be a normal, connected noetherian scheme, whose function field K is absolutely finitely generated. Let $f: X \rightarrow S$ be a smooth surjective morphism of finite type whose geometric generic fibre is connected, and*

which admits a cross-section $X \xrightarrow{\epsilon} S$. Then there are only finitely many connected finite etale X -schemes Y/X which are galois over X with abelian galois group of order prime to $\text{char}(K)$ and which are completely decomposed over the marked section. If in addition we suppose X/S proper, we can drop the proviso "of order prime to $\text{char}(K)$ ".

Proof. This is just the concatenation of Theorems 1 and 2 with the physical interpretation (1.3) of the group $\text{Ker}(X/S)$ in the presence of a section. QED

V. APPLICATION TO l -ADIC REPRESENTATIONS

Let l be a prime number, $\overline{\mathbf{Q}}_l$ an algebraic closure of \mathbf{Q}_l . By an l -adic representation ρ of a topological group π , we mean a finite-dimensional continuous representation

$$\rho: \pi \rightarrow GL(n, \overline{\mathbf{Q}}_l)$$

whose image lies in $GL(n, E_\lambda)$ for some finite extension E_λ of \mathbf{Q}_l .

THEOREM 6. (cf. Grothendieck, via [2], 1.3). *Let K be an absolutely finitely generated field, X/K a smooth, geometrically connected K -scheme of finite type, \bar{x} a geometric point of $X \otimes \overline{K}$, x the image geometric point of \bar{x} in X . Let l be a prime number, and ρ an l -adic representation of $\pi_1(X, x)$;*

$$\rho: \pi_1(X, x) \rightarrow GL(n, \overline{\mathbf{Q}}_l).$$

Let G be the Zariski closure of the image $\rho(\pi_1(X \otimes \overline{K}, \bar{x}))$ of the geometric fundamental group $\pi_1(X \otimes \overline{K}, \bar{x})$ in $GL(n, \overline{\mathbf{Q}}_l)$ and G^0 its identity component. Suppose that either l is different from the characteristic p of K , or that X/K is proper. Then:

- (1) *the radical of G^0 is unipotent, or equivalently:*
- (2) *if the restriction of ρ to the geometric fundamental group $\pi_1(X \otimes \overline{K}, \bar{x})$ is completely reducible, then the algebraic group G^0 is semi-simple.*

Proof. By Theorem 1, for $l \neq p$, or by Theorem 2 if $l = p$ and X/K is proper, we know that the l -part of $\text{Ker}(X/K)$ is finite i.e. (cf. Lemma 1) the image of $\pi_1(X \otimes \bar{K}, \bar{x})$ in $\pi_1(X)^{ab}$ is the product of a finite group and a group of order prime to l . Given this fact, the proof proceeds exactly as in (Deligne [2], 1.3).

QED

Remarks. (1) This theorem is the group-theoretic version of Grothendieck's local monodromy theorem (cf. Serre-Tate ([15], Appendix) for a precise statement, as well as the proof) with X/K "replacing" the spectrum of the fraction field E of a henselian discrete valuation ring with residue field K , and with $\pi_1(X \otimes \bar{K})$ "replacing" the inertia subgroup I of $\text{Gal}(\bar{E}/E)$. The "extra" feature of the "local" case is that the quotient of I by a normal pro- p subgroup is abelian. Therefore any l -adic representation ρ of I , with $l \neq p$, becomes *abelian* when restricted to a suitable open subgroup of I , and hence the associated algebraic group G^0 is automatically abelian. In particular, the radical of G^0 is G^0 itself.

(2) If X/K is itself an abelian variety A/K , then $\pi_1(A \otimes \bar{K}, \bar{x})$ is abelian. Therefore if l is any prime, and ρ any l -adic representation of $\pi_1(A \otimes \bar{K}, \bar{x})$, the associated algebraic groups G and G^0 will be abelian; hence if ρ is the restriction to $\pi_1(A \otimes \bar{K}, \bar{x})$ of an l -adic representation of $\pi_1(A, x)$, then G^0 is unipotent, i.e. the restriction of ρ to an open subgroup of $\pi_1(A \otimes \bar{K}, \bar{x})$ is *unipotent* (compare Oort [11], 2).

(3) Can one give an example of X/K proper smooth and geometrically connected over an absolutely finitely generated field K of characteristic $p > 0$ whose fundamental group $\pi_1(X, x)$ admits an n -dimensional p -adic representation with $n \geq 2$ (resp. $n \geq 3$) for which the associated algebraic group G^0 is $SL(n)$ (resp. $SO(n)$)? Can we find an abelian scheme A over such an X , all of whose fibres have the same p -rank $n \geq 2$, for which the associated p -adic representation of $\pi_1(X, x)$ has $G^0 = SL(n)$? (cf. Oort [11] for the case of p -rank zero).