

# 5. FURTHER NOTES

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **28 (1982)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

since by (3.15) the Lie algebra cohomology space in Theorem 3.16 is a formal harmonic space. It is even true as a matter of fact that under reasonable conditions the formal harmonic space associated to a polarization coincides with a Lie algebra cohomology space; see Penney's Theorem 2 in [68]. The latter cohomology spaces have the form  $H^j(p \cap \text{Ker } \Lambda, \pi_{-\Lambda}^\infty)$  where  $\pi^\infty$  is the space of  $C^\infty$  vectors in a representation  $\pi$  and  $\Lambda$  is considered also as a linear functional on  $g^C$ . By very clever means these spaces are shown to vanish for all  $j$  except  $j =$  the negativity index  $q(p, \Lambda)$  of the polarization (see 4.1). Moreover  $H^{q(p, \Lambda)}(p \cap \text{Ker } \Lambda, \pi_{-\Lambda}^\infty)$  is one-dimensional; see Rosenberg's Theorem 2.4 in [74]; also see Penney [69]. With these remarks in mind an application of Theorem 4.3 gives

**THEOREM 4.5** (J. Rosenberg–R. Penney 1979). *Let  $G$  be a connected, simply connected nilpotent Lie group and let  $p$  be a relatively ideal complex polarization at  $\Lambda \in g^*$ ,  $g = \text{Lie algebra of } G$ . Let  $\pi^j(\Lambda, p, G)$  be the  $j$ -th harmonically induced representation defined in (4.2). Then  $\pi^j(\Lambda, p, G)$  vanishes for  $j \neq$  the negativity index  $q(p, \Lambda)$  (see (4.1)). Moreover  $\pi^{q(p, \Lambda)}(\Lambda, p, G)$  is irreducible and unitarily equivalent to the Kirillov representation  $\pi_\Lambda$ .*

Theorem 4.5 is clearly analogous to Theorem 3.11 and thus it represents the confirmation of a version of the Kostant-Langlands conjecture for nilpotent Lie groups. One may add that as a matter of fact the distinguished integer  $q_\Lambda$  in (3.9) is indeed the negativity index of a complex polarization—namely the polarization is a Borel subalgebra at a regular point.

## 5. FURTHER NOTES

1. We have pointed out earlier that in addition to Schmid's thesis work, early efforts towards proving the Kostant-Langlands conjecture were made by Narasimhan and Okamoto. The latter authors considered the special case when  $G/K$  admits a  $G$  invariant complex structure<sup>1)</sup>. They constructed unitary representations  $\pi_\Lambda^{0,j}$  of  $G$  on  $L_2$ -cohomology spaces associated to holomorphic vector bundles  $E_\Lambda$  over  $G/K$  induced by an irreducible unitary representation of  $K$  with highest weight  $\Lambda$ ; compare remarks following (3.10). The  $\pi_\Lambda^{0,j}$  are shown to be subject to an important *alternating sum formula* which, roughly stated, says that

$$(5.1) \quad \sum_{j=0}^n (-1)^j \text{ character of } \pi_\Lambda^{0,j} = (-1)^{q_\Lambda} \text{ character of } \pi_\Lambda^*$$

---

<sup>1)</sup> Here  $G, K$  are as in section 3.

where  $\pi_\Lambda^*$  is the contragredient to the discrete class  $\pi_\Lambda$  given in Theorem 3.5,  $q_\Lambda$  is the number of non-compact positive roots  $\alpha$  such that  $(\Lambda + \delta, \alpha) > 0$  (compare (3.9)), and  $n = \frac{1}{2} \dim_R G/K$ . A precise statement of (5.1) is given in Theorem 1 of

[60]. Once one has an alternating sum formula the class  $\pi_\Lambda^*$  can be realized by  $\pi_\Lambda^{0, q_\Lambda}$  if a *vanishing theorem*  $H_{\delta, 2}^{0, j}(G/K, E_\Lambda) = 0$  for  $j \neq q_\Lambda$  is proved. The methods of Narasimhan and Okamoto of establishing an alternating sum formula and a vanishing theorem served as a prototype for the work of later authors; see for example [59], [65], [99], [64], [98]. The vanishing theorems in [60] are improved by Parthasarathy in [64].

2. The ambitious program of decomposing a complex flag manifold under the action of a real group and of using the real group orbits as the setting for the geometric realization of unitary representations of semisimple (even reductive) Lie groups is carried out in the profound work of J. Wolf in [97], [99], [96]. A family of unitary representations which support the Plancherel measure are realized on *partially holomorphic* cohomology spaces. This family clearly contains many non-discrete classes. The realizations are similar to realizations by the Kostant-Kirillov method where one uses polarizations of semisimple orbits. However some interesting differences occur in the case when the reductive group has non-commutative Cartan subgroups; see [86], [98].

3. After (3.9) we remarked on the similarity in appearance of the Weyl and Harish-Chandra character formulas of Theorems 2.22 and 3.5. There is however a vast difference of roles which these formulas play. For example Weyl's formula determines the character *on all of the group* (since a compact group is covered by conjugates of a maximal torus) whereas conjugates of  $H$  in section 3 certainly do not cover  $G$ . It seems to be an extremely difficult problem (and perhaps an impossible one to solve) to obtain the character formula explicitly on an *arbitrary* Cartan subgroup  $H$  of a non-compact semisimple group.

4. A new proof of Harish-Chandra's regularity theorem for invariant eigendistributions (cf. remarks following (3.4) due to Atiyah and Schmid is now available; cf. [4], [7], [81]. [6] contains a new and largely self-contained account of the principal theory of the discrete series including existence theory, exhaustion, geometric realization, character formulae, and character behavior. These new methods rely on the Atiyah-Singer  $L_2$ -index theorem [3], [31], [32], [81], [83], [84].

5. It is possible to formulate Frobenius reciprocity for unitary representations on a Hilbert space  $\mathcal{H}(D)$  of  $L_2$ -solutions of an invariant elliptic differential operator  $D$  on homogeneous bundles over a homogeneous space  $G/H$  whose isotropy subgroup  $H$  is compact modulo the center of  $G$ . Here  $G$  is a connected unimodular Lie group (not necessarily semisimple) subject to some mild structural constraints. In [33] Connes and Moscovici show that  $\mathcal{H}(D)$  decomposes as a finite direct sum of irreducible unitary representations all of which are square-integrable modulo the center of  $G$  and occur with finite multiplicity. They derive for  $\mathcal{H}(D)$  a reciprocity analogous to that expressed for the  $L_2$ -cohomology spaces in Theorem 3.15 and Theorem 4.3.

## REFERENCES

- [1] AHIEZER, D. Cohomology of compact homogeneous spaces. *Mat. Sb.* 84 (126) (1971), 290-300 = *Math. USSR Sb.* 13 (1971), 285-296.
- [2] ANDREOTTI, A. and E. VESSENTINI. Carleman estimates for the Laplace-Beltrami operator on complex manifolds. *Publ. I.H.E.S.* No. 25 (1965), 313-362.
- [3] ATIYAH, M. Elliptic operators, discrete groups and von Neumann algebras. *Astérisque* 32-33 (1976), 43-72.
- [4] —— *Characters of semisimple Lie groups*. Mimeographed lecture notes, Univ. of Oxford.
- [5] —— The Harish-Chandra character. *Lecture No. 7 from Proceedings of the SRC/LMS Research Symposium on Representations of Lie Groups, London Math. Soc. Series 34*, Cambridge Univ. Press, 176-181.
- [6] ATIYAH, M. and W. SCHMID. A geometric construction of the discrete series for semi-simple Lie groups. *Inventiones Math.* 42 (1977), 1-62.
- [7] —— A new proof of the regularity theorem for invariant eigendistributions on semisimple Lie groups. *To appear*.
- [8] AUSLANDER, L. and B. KOSTANT. Polarization and unitary representations of solvable Lie groups. *Inventiones Math.* 14 (1971), 255-354.
- [9] BERNAT, P., N. CONZE, M. DUFLO, M. LÉVY-NAHAS, M. RAIS, P. RENOUARD et M. VERGNE. *Représentations des groupes de Lie résolubles*. Monographies de la Société Mathématique de France, Dunod, éditeur. (1972) Paris.
- [10] BLATTNER, R. On induced representations. *Amer. J. of Math.* 83 (1961), 79-98.
- [11] BOREL, A. et A. WEIL. Représentations linéaires et espaces homogènes Kähleriens des groupes de Lie compacts. *Séminaire Bourbaki*, May 1954 (exposé by J.-P. Serre).
- [12] BOTT, R. Homogeneous vector bundles. *Annals of Math.* (2) 66 (1957), 203-248.
- [13] —— Induced representations. *Seminars on Analytic Functions*, Vol. 2, Institute for Advanced Study, Princeton, N.J., 1957.
- [14] BRUHAT, F. Travaux de Harish-Chandra. *Séminaire Bourbaki*, exposé 143 (1957), 1-9.
- [15] CARMONA, J. Représentations du groupe de Heisenberg dans les espaces de  $(0, q)$ -formes. *Math. Ann.* 205 (1973), 89-112.
- [16] CARTAN, E. Les tenseurs irréductibles et les groupes simples et semisimples. *Bull. Sci. Math.* 49 (1925), 130-152.