

5.2. Spherical functions of type

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **28 (1982)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

5.2. SPHERICAL FUNCTIONS OF TYPE δ

Let G be a unimodular lcsc. group with compact subgroup K . Let

$$K^* := \{(k, k) \in G \times K \mid k \in K\}.$$

Let $\delta \in \hat{K}$ and let τ be a K -unitary representation of G . Then $\tau \otimes \check{\delta}$ ($\check{\delta}$ the contragredient representation to δ) is a K^* -unitary representation of $G \times K$ on $\mathcal{H}(\tau) \otimes \mathcal{H}(\delta)$.

LEMMA 5.1. *The multiplicity of δ in $\tau|_K$ is equal to the multiplicity of the representation 1 of K^* in $\tau \otimes \check{\delta}|_{K^*}$. τ is irreducible iff $\tau \otimes \check{\delta}$ is irreducible. τ is unitary iff $\tau \otimes \check{\delta}$ is unitary.*

This can be proved immediately. By using the results summarized in §5.1 we conclude that $(G \times K, K^*)$ is a Gelfand pair if there exists a continuous involutive homomorphism α on G such that for each $(g, k) \in G \times K$ we have $\alpha(g) = k_1 g^{-1} k_2$, $\alpha(k) = k_1 k^{-1} k_2$ for certain $k_1, k_2 \in K$. Furthermore, if $(G \times K, K^*)$ is a Gelfand pair and if the irreducible representation τ of G is unitary or K -finite then τ is K -multiplicity free. In particular, this applies to $SU(1, 1)$:

PROPOSITION 5.2. *If $G = SU(1, 1)$ then $(G \times K, K^*)$ is a Gelfand pair.*

Proof. For $g \in SU(1, 1)$ define $\alpha(g) := {}^t(g^{-1})$. Then α is a continuous involutive automorphism on G and $\alpha(a_t) = a_{-t}$ on A , $\alpha(u_\theta) = u_{-\theta}$ on K . Since $G = KAK$, α has the required properties. \square

Let $(G \times K, K^*)$ be a Gelfand pair. Identify $G \times \{e\}$ with G . A spherical function on $G \times K$ is completely determined by its restriction to G . By using the results mentioned in §5.1 we obtain the following properties. First, a continuous function ϕ on G is the restriction to G of a spherical function on $G \times K$ iff $\phi \neq 0$ and

$$\phi(x)\phi(y) = \int_K \phi(xkyk^{-1})dk, \quad x, y \in G.$$

Next, let

$$\begin{aligned} & I_c(G) \text{ (or } I_c^\infty(G)) \\ & := \{f \in C_c(G) \text{ (or } C_c^\infty(G)) \mid f(kgk^{-1}) = f(g), \\ & \quad g \in G, k \in K\}. \end{aligned}$$

These are commutative topological algebras under convolution and their characters are precisely of the form (5.1), where ϕ is a spherical function on $G \times K$. If ϕ is a spherical function on $G \times K$ then there is a $\delta \in \hat{K}$ such that for all $x \in G$ the function $k \rightarrow \phi(xk)$ on K belongs to δ . Then δ is called a *spherical function of type δ* on G (with respect to K), cf. GODEMENT [19]. It is funny that spherical functions of type δ are on the one hand generalizations of ordinary spherical functions for (G, K) , on the other hand restrictions to G of ordinary spherical functions for $(G \times K, K^*)$.

For convenience, we take a one-dimensional $\delta \in \hat{K}$. Then a spherical function ϕ on $G \times K$ is of type δ iff

$$\phi(xk) = \phi(kx) = \delta(k)\phi(x), \quad x \in G, k \in K.$$

Let

$$\begin{aligned} & I_{c, \delta}(G) \text{ (or } I_{c, \delta}^\infty(G)) \\ & := \{f \in C_c(G) \text{ (or } C_c^\infty(G)) \mid f(xk) = f(kx) \\ & \quad = \delta(k)f(x), x \in G, k \in K\}. \end{aligned}$$

These are closed subalgebras of $I_c(G)$ (or $I_c^\infty(G)$) and their characters are precisely of the form (5.1), where ϕ is a spherical function of type δ . Finally, if τ is a K -unitary representation of G and if $\mathcal{H}(\tau)$ contains a unit vector v satisfying $\tau(k)v = \delta(k)v$, unique up to a constant factor, then $x \rightarrow (\tau(x)v, v)$ is a spherical function of type δ .

5.3. THE GENERALIZED ABEL TRANSFORM

Let G be a connected noncompact real semisimple Lie group with finite center. Use the notation of §2.2. For given Haar measures dk, da, dn on K, A, N , respectively, normalize the Haar measure on G such that

$$(5.2) \quad \int_G f(g)dg = \int_{K \times A \times N} f(kan)e^{2\rho(\log a)} dk da dn, \quad f \in C_c(G)$$

(cf. HELGASON [25, Ch. X, Prop. 1.11]). Note the property

$$(5.3) \quad \int_N f(n)dn = e^{2\rho(\log a)} \int_N f(ana^{-1})dn, \quad f \in C_c(N), a \in A$$

(cf. [25, Ch. X, proof of Prop. 1.11]).