

6.1. A CRITERIUM FOR UNITARIZABILITY

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **28 (1982)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

theorem for noncompact semisimple Lie groups due to Casselman (cf. WALLACH [47, Cor. 7.5]). Casselman's theorem improves HARISH-CHANDRA's [22, Theorem 4] *subquotient theorem*.

5.6.3. The generalized Abel transform $f \rightarrow F_f^\delta$ can be defined for general K -type δ . It was introduced by HARISH-CHANDRA [24, p. 595] in the spherical case, TAKAHASHI [40, §2] in the case $G = SO_0(n, 1)$ and WARNER [49, 6.2.2] in the general case. The injectivity of this transform holds generally, cf. WARNER [49]. The image of $I_{c, \delta}^\infty(G)$ under this transform is known in the spherical case (cf. GANGOLLI [16]) and if G has real rank 1 and δ is one-dimensional (cf. WALLACH [46]), but seems to be unknown in the general case (cf. WARNER [49, p. 36]).

5.6.4. In [39] TAKAHASHI also reduces the proof of Theorem 5.4 to Proposition 5.5. However, he proves Prop. 5.5 by considering eigenfunctions of the Casimir operator, since he did not know, then, how to invert the transform $f \rightarrow F_f^n$. In [42] he independently obtained a proof of Prop. 5.5 similar to ours. Earlier, in [40, §4.1] he used a similar method in the spherical case of $G = SO_0(n, 1)$. NAIMARK [34, Ch. 3, §9] proved the subquotient theorem for $SL(2, \mathbf{C})$ by methods somewhat related to ours.

5.6.5. Part of Lemma 5.8 is contained in WHITNEY [50]. See SCHWARZ [37] for a theorem on C^∞ -functions which are invariant under a more general Weyl group.

5.6.6. Theorem 5.10 more generally holds with Gegenbauer polynomials of integer or half integer order as kernels, cf. DEANS [6], [7], KOORNWINDER [27, §5.9]. Deans' proof uses the inversion formula for the Radon transform. The author's proof uses Weyl fractional integral transforms and generalized fractional integral transforms studied by SPRINKHUIZEN [38]. MATSUSHITA [30, §2.3] considers the transformation $f \rightarrow F_f^n$ for general real n in the context of the universal covering group of $SL(2, \mathbf{R})$ and he derives the inversion formula with a proof due to T. Shintani, which uses Mellin transforms.

6. UNITARIZABILITY OF IRREDUCIBLE SUBREPRESENTATIONS OF THE PRINCIPAL SERIES

6.1. A CRITERIUM FOR UNITARIZABILITY

Remember that a representation of an lcsc. group G on a Hilbert space is strongly continuous if and only if it is weakly continuous (cf. WARNER [48, Prop. 4.2.2.1]). Thus, if τ is a (strongly continuous) Hilbert representation of G then $\tilde{\tau}$ defined by

$$(6.1) \quad \tilde{\tau}(g) := \tau(g^{-1})^*, g \in G,$$

is again a (strongly continuous) Hilbert representation of G on $\mathcal{H}(\tau)$. The representation $\tilde{\tau}$ is called the *conjugate contragredient* to τ . The representation τ is unitary iff $\tilde{\tau} = \tau$.

THEOREM 6.1. *Let G be an lcsc. group with compact abelian subgroup K . Let τ be a K -multiplicity free representation of G . Let $\{\phi_\delta\}$ be a K -basis for $\mathcal{H}(\tau)$. Let $c_\delta (\delta \in \mathcal{M}(\tau))$ be positive real numbers. Then the following statements are equivalent to each other :*

(a) τ is Naimark equivalent to some unitary representation.

(b) $\tau \xrightarrow{A} \tilde{\tau}$ with $A\phi_\delta = c_\delta \phi_\delta (\delta \in \mathcal{M}(\tau))$.

(c) $\overline{\tau_{\gamma, \delta}(g^{-1})} = \frac{c_\delta}{c_\gamma} \tau_{\delta, \gamma}(g), \gamma, \delta \in \mathcal{M}(\tau), g \in G$.

If, moreover, τ is irreducible then (a), (b) and (c) are equivalent to :

(d) For some $\delta \in \mathcal{M}(\tau)$ we have

$$\overline{\tau_{\gamma, \delta}(g^{-1})} = \frac{c_\delta}{c_\gamma} \tau_{\delta, \gamma}(g), g \in G, \text{ for all } \gamma \in \mathcal{M}(\tau).$$

If (b) holds then $\tau(g)(g \in G)$ is unitary with respect to a new inner product $\langle \cdot, \cdot \rangle$ on $\mathcal{D}(A)$ defined by

$$(6.2) \quad \langle \phi_\gamma, \phi_\delta \rangle := \begin{cases} 0 & \text{if } \gamma \neq \delta, \\ c_\delta & \text{if } \gamma = \delta. \end{cases}$$

Proof. First observe that $\tilde{\tau}(k)\phi_\delta = \delta(k)\phi_\delta (k \in K)$, so $\{\phi_\delta\}$ is a K -basis with respect to $\tilde{\tau}$ as well. We have

$$(6.3) \quad \tilde{\tau}_{\gamma, \delta}(g) = \overline{\tau_{\delta, \gamma}(g^{-1})}.$$

(a) \Rightarrow (b) : Let $\tau \xrightarrow{B} \sigma$ with σ unitary. Then $\sigma = \tilde{\sigma}$ and $\tilde{\sigma} \xrightarrow{B^*} \tilde{\tau}$. Let $\{\psi_\delta\}$ be a K -basis for $\mathcal{H}(\sigma)$. Let $B\phi_\delta = b_\delta \psi_\delta (\delta \in \mathcal{M}(\tau))$. Then, by Theorem 4.5:

$$\tilde{\tau}_{\gamma, \delta} = \overline{\frac{b_\gamma}{b_\delta}} \sigma_{\gamma, \delta} = \left| \frac{b_\gamma}{b_\delta} \right|^2 \tau_{\gamma, \delta},$$

so (b) holds.

(b) \Rightarrow (a) : Assume (b). Then A is self-adjoint and positive definite. Define a new inner product $\langle \cdot, \cdot \rangle$ on $\mathcal{D}(A)$ by $\langle v, w \rangle := (Av, w)$. Then, for $v, w \in \mathcal{D}(A)$, $g \in G$, we have :

$$\begin{aligned}\langle \tau(g)v, \tau(g)w \rangle &= (A\tau(g)v, \tau(g)w) = (\tilde{\tau}(g^{-1})A\tau(g)v, w) \\ &= (A\tau(g^{-1})\tau(g)v, w) = (Av, w) = \langle v, w \rangle,\end{aligned}$$

i.e. $\langle \tau(g)v, \tau(g)w \rangle = \langle v, w \rangle$. Thus τ is a unitary representation on $\mathcal{D}(A)$ with respect to the new inner product. (Weak continuity of τ is easily proved.) Let σ be the extension of this representation to a unitary representation in the Hilbert space completion $\overset{B}{\mathcal{H}}(\sigma)$ of $\mathcal{D}(A)$ with respect to $\langle \cdot, \cdot \rangle$. Then $\tau \simeq \sigma$, where B is the closure of the identity operator on $\mathcal{D}(A)$ (cf. Lemma 4.4). Note that we have also proved the last part of the theorem.

The equivalence of (c) or (d) with (b) follows from Theorem 4.5. \square

6.2. THE CASE $SU(1, 1)$

It follows from (2.30) that

$$(6.4) \quad \overline{c_{\xi, \lambda, n, m}} = (-1)^{m-n} c_{\xi, -\bar{\lambda}, m, n}.$$

Combination of (6.3), (2.29) and (6.4) yields

$$(6.5) \quad \tilde{\pi}_{\xi, \lambda} = \pi_{\xi, -\bar{\lambda}}.$$

In §6.1 we showed that a necessary condition for unitarizability of an irreducible subquotient representation τ of $\pi_{\xi, \lambda}$ is the equivalence of τ and $\tilde{\tau}$. In view of (6.5) and Theorem 4.7 this is only possible if $\bar{\lambda} = \pm \lambda$, that is, if λ is real or imaginary. If λ is imaginary then $\tilde{\pi}_{\xi, \lambda} = \pi_{\xi, \lambda}$, so $\pi_{\xi, \lambda}$ is already unitary. Let us now examine the case that λ is real and nonzero. Then $\tilde{\pi}_{\xi, \lambda} = \pi_{\xi, -\lambda}$. If τ is an irreducible subquotient representation of $\pi_{\xi, \lambda}$ then $\tau \overset{A}{\simeq} \tilde{\tau}$ with (cf. (4.10))

$$(6.6) \quad A\phi_m = c_{\xi, \lambda, m} \phi_m, \quad \phi_m \in \mathcal{H}(\tau),$$

where $c_{\xi, \lambda, m}$ is given by (4.9). Now a sufficient condition for the unitarizability of τ is that the coefficients $c_{\xi, \lambda, m}$ are all positive or all negative for $\phi_m \in \mathcal{H}(\tau)$. Referring to the classification in Theorem 3.4 we will examine these coefficients. (Because of equivalence, it is not necessary to treat the cases where $\lambda < 0$.)