

§3. CONFORMAL ACTIONS ON SURFACES OF LEAST GENUS

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **31 (1985)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **26.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

The only identifications under the action are: vu gets identified to vx and wu gets identified to wx . It follows that $P/T(r, s, t)$ is the 2 sphere and the branched covering $P/\Delta \rightarrow P/T(r, s, t)$ has 3 branch points coming from the vertices u, v, w .

Now notice that Δ is torsion free. This follows from the facts:

(1) the elements of finite order in $T(r, s, t)$ are the conjugates of A, B, C .

(2) elements of finite order in $T(r, s, t)$ map to elements of the same order in G . From this it follows that the orders of the branch points are r, s, t respectively.

Finally we consider the Riemann-Hurwitz formula:

$$\chi(P/\Delta) = |G| \left(\chi(P/T(r, s, t)) - \left(1 - \frac{1}{r}\right) - \left(1 - \frac{1}{s}\right) - \left(1 - \frac{1}{t}\right) \right)$$

i.e.,

$$2 - 2g = |G| \left(\frac{1}{r} + \frac{1}{s} + \frac{1}{t} - 1 \right).$$

Therefore
$$g = 1 + \frac{|G|}{2} \left(1 - \frac{1}{r} - \frac{1}{s} - \frac{1}{t} \right)$$
 Q.e.d.

§ 3. CONFORMAL ACTIONS ON SURFACES OF LEAST GENUS

If (A, B, C) is an (r, s, t) triple generating $PSl_2(p)$ then we have a short exact sequence

$$1 \rightarrow \Delta \rightarrow T(r, s, t) \rightarrow PSl_2(p) \rightarrow 1$$

where Δ is torsion free. Then it follows that $H/T(r, s, t)$ is S^2 and the branched covering $H/\Delta \rightarrow H/T(r, s, t)$ has 3 branch points with orders r, s, t .

Conversely we have:

(3.1). THEOREM. *If S is a Riemann surface of least genus for $PSl_2(p)$ then $S/PSl_2(p)$ is S^2 and $\pi: S \rightarrow S/PSl_2(p)$ has 3 branch points.*

Proof. There exists a short exact sequence $1 \rightarrow \Delta \rightarrow T(2, 3, p) \rightarrow PSl_2(p) \rightarrow 1$ arising from a $(2, 3, p)$ triple and consequently

$$\text{genus}(H/\Delta) = 1 + \frac{|G|}{2} \left(\frac{1}{6} - \frac{1}{p} \right).$$

Let $g = \text{genus}(S)$, $h = \text{genus}(S/PSl_2(p))$ and suppose $\pi: S \rightarrow S/PSl_2(p)$ has b branch points x_1, \dots, x_b of respective orders n_1, \dots, n_b . Then the Riemann-Hurwitz formula tells us

$$(3.2). \quad 2 - 2g = |G| \left(2 - 2h - \sum \left(1 - \frac{1}{n_i} \right) \right).$$

That is $g = 1 + \frac{|G|}{2} \left(2h - 2 + \sum \left(1 - \frac{1}{n_i} \right) \right)$. Since g is the least genus this leads to the inequality

$$(3.3). \quad 2h - 2 + \sum \left(1 - \frac{1}{n_i} \right) \leq \frac{1}{6} - \frac{1}{p}.$$

From this we immediately see that $h = 0, 1$.

Therefore we suppose that $h = 1$. Since all $n_i \geq 2$ this implies that $b = 0$, and hence $PSl_2(p)$ is acting fixed point freely on S with orbit space the torus. But this immediately gives an epimorphism $\mathbf{Z} \oplus \mathbf{Z} \twoheadrightarrow PSl_2(p)$. However, this is a contradiction since G is not abelian. Therefore $h = 0$ and $S/PSl_2(p)$ is a 2-sphere.

To prove that there are 3 branch points put $h = 0$ into (3.3):

$$(3.4) \quad -2 + \sum_{i=1}^b \left(1 - \frac{1}{n_i} \right) \leq \frac{1}{6} - \frac{1}{p}.$$

Since $1 - \frac{1}{n_i} \geq \frac{1}{2}$ for all i this gives $b \leq 4$. If $b = 0$ we have an unbranched covering $S \rightarrow S^2$ with deck transformation group $PSl_2(p)$. But this is clearly a contradiction.

Thus assume $b = 1$. Then we have the regular unbranched covering

$$S - \pi^{-1}(x_1) \rightarrow S^2 - \{x_1\}$$

with deck transformation group $PSl_2(p)$. But again this is impossible since $S^2 - \{x_1\} \cong \mathbf{R}^2$.

Next we put $b = 2$ and consider the regular covering

$$S - \pi^{-1}\{x_1, x_2\} \rightarrow S^2 - \{x_1, x_2\}.$$

Then we have the exact sequence coming from fundamental groups $1 \rightarrow \mathbf{Z} \rightarrow \mathbf{Z} \rightarrow PSl_2(p) \rightarrow 1$, which is again a contradiction.

Finally we suppose $b = 4$. The inequality (3.4) is

$$(3.5). \quad 2 - \frac{1}{n_1} - \frac{1}{n_2} - \frac{1}{n_3} - \frac{1}{n_4} \leq \frac{1}{6} - \frac{1}{p}$$

and is clearly satisfied by $n_1 = n_2 = n_3 = n_4 = 2$. However, this choice of n_i 's gives $g = 1$ by (3.2); in other words $PSl_2(p)$ is toroidal. However, no nonabelian finite simple group G can act on $S^1 \times S^1$ because covering space theory implies there are branch points and hence the orbit space is S^2 . Hence the induced homomorphism $PSl_2(p) \rightarrow \text{Aut}(\mathbb{Z}^2)$ is nontrivial and also has a kernel since $\text{Aut}(\mathbb{Z}^2)$ has no p torsion for $p \geq 7$. This contradicts G simple. Therefore this case is excluded and we have

$$2 - \frac{1}{n_1} - \frac{1}{n_2} - \frac{1}{n_3} - \frac{1}{n_4} \geq 2 - \frac{1}{2} - \frac{1}{2} - \frac{1}{2} - \frac{1}{3} = \frac{1}{6}$$

contradicting (3.5).

Q.e.d.

If we let the orders of the 3 branch points be r, s, t then the Riemann-Hurwitz formula is

$$\chi(S) = |PSl_2(p)| \left(2 - \left(1 - \frac{1}{r} \right) - \left(1 - \frac{1}{s} \right) - \left(1 - \frac{1}{t} \right) \right).$$

But $|PSl_2(p)| = \frac{p(p^2-1)}{2}$ and therefore

$$\text{genus}(S) = 1 + \frac{p(p^2-1)}{4} \left(1 - \frac{1}{r} - \frac{1}{s} - \frac{1}{t} \right).$$

To take advantage of this formula we must know what sort of branching data (r, s, t) can occur. To this end we quote a very general theorem of Tucker [T].

(3.6). THEOREM. Suppose G is a finite group acting effectively on a closed orientable surface S by orientation preserving homeomorphisms. If $g = \text{genus}(S/G)$ and there are b branch points of orders n_1, \dots, n_b then G has a presentation of the form

$$\{x_1, y_1, \dots, x_g, y_g, e_1, \dots, e_b \mid \prod_{i=1}^g [x_i, y_i] e_1 \dots e_b = e_1^{n_1} = \dots = e_b^{n_b} = 1, ETC\}$$

(3.7). COROLLARY. If S is a Riemann surface of least genus for $PSl_2(p)$ then there exist integers $r, s, t, \geq 2$ so that

(a) there is an extension $1 \rightarrow \Delta \rightarrow T(r, s, t) \rightarrow PSl_2(p) \rightarrow 1$;

$$(b) \text{ genus } (PSl_2(p)) = 1 + \frac{p(p^2-1)}{4} \left(1 - \frac{1}{r} - \frac{1}{s} - \frac{1}{t} \right).$$

If A, B, C are the usual generators of $T(r, s, t)$ then it is in fact true that the orders of A, B, C in $PSl_2(p)$ are r, s, t . Putting (2.22) and (3.7) together gives

(3.8). COROLLARY. *The genus of $PSl_2(p)$ is given by*

$$g = \min \left\{ 1 + \frac{p(p^2-1)}{4} \left(1 - \frac{1}{r} - \frac{1}{s} - \frac{1}{t} \right) \right\}$$

where the minimum is taken over all (r, s, t) for which there exist (r, s, t) triples generating $PSl_2(p)$.

The last step in the determination of the genus is to identify those (r, s, t) which are relevant. This is accomplished in the following manner:

(1) first find all (r, s, t) so that

$$1 - \frac{1}{r} - \frac{1}{s} - \frac{1}{t} \leq \frac{1}{6} - \frac{1}{d}, \text{ assuming } p \geq 13.$$

- (2) then eliminate those triples (r, s, t) corresponding to either spherical or Euclidean triangle groups.
- (3) make a comparison of the triples remaining so as to eliminate those with larger genus.

In the following table we give some pertinent data:

TABLE I

| (r, s, t) | $1 - \frac{1}{r} - \frac{1}{s} - \frac{1}{t}$ | type | condition for $1 - \frac{1}{r} - \frac{1}{s} - \frac{1}{t} \leq \frac{1}{6} - \frac{1}{d}$ |
|----------------------------------|---|------------|---|
| $(2, 2, t)$ | $-\frac{1}{t}$ | spherical | $d \geq 6$ |
| $(2, 3, t)$ where $3 < t < 5$ | $\frac{1}{6} - \frac{1}{t}$ | spherical | $t \leq d$ |
| $(2, 3, 6)$ | 0 | euclidean | $t < d$ |
| $(2, 3, t)$ where $t > 7$ | $\frac{1}{6} - \frac{1}{t}$ | hyperbolic | $t \leq d$ |
| $(2, 4, 4)$ | 0 | euclidean | $d > 6$ |
| $(2, 4, 5)$ | $\frac{1}{20}$ | hyperbolic | $d \geq 9$ |
| $(2, 4, 6)$ | $\frac{1}{12}$ | hyperbolic | $d \geq 12$ |
| $(2, 4, 7)$ | $\frac{3}{28}$ | hyperbolic | $d \geq 17$ |
| $(2, 4, 8)$ | $\frac{1}{8}$ | hyperbolic | $d \geq 24$ |
| $(2, 4, 9)$ | $\frac{5}{36}$ | hyperbolic | $d \geq 36$ |
| $(2, 4, 10)$ | $\frac{3}{20}$ | hyperbolic | $d \geq 60$ |
| $(2, 4, 11)$ | $\frac{7}{44}$ | hyperbolic | $d \geq 132$ |

TABLE I (*suite*)

| | (r, s, t) | $1 - \frac{1}{r} - \frac{1}{s} - \frac{1}{t}$ | <i>type</i> | condition for $1 - \frac{1}{r} - \frac{1}{s} - \frac{1}{t} \leq \frac{1}{6}$ |
|-----|---|--|-------------|---|
| 13. | $(2, 4, t)$ | $\frac{1}{4} - \frac{1}{t} \geq \frac{1}{6}$ | hyperbolic | never |
| 14. | $(2, 5, 5)$ | $\frac{1}{10}$ | hyperbolic | $d \geq 15$ |
| 15. | $(2, 5, 6)$ | $\frac{2}{15}$ | hyperbolic | $d \geq 30$ |
| 16. | $(2, 5, 7)$ | $\frac{11}{70}$ | hyperbolic | $d \geq 105$ |
| 17. | $(2, 5, t)$ where $t > 8$ | $\frac{3}{10} - \frac{1}{t} > \frac{1}{6}$ | hyperbolic | never |
| 18. | $(2, s, t)$ | $\frac{1}{2} - \frac{1}{s} - \frac{1}{t} \geq \frac{1}{6}$ | hyperbolic | never |
| 19. | $(3, 3, 3)$ | 0 | euclidean | $d > 7$ |
| 20. | $(3, 3, 4)$ | $\frac{1}{12}$ | hyperbolic | $d \geq 12$ |
| 21. | $(3, 3, 5)$ | $\frac{2}{15}$ | hyperbolic | $d \geq 30$ |
| 22. | $(3, 3, t)$ | $\frac{1}{3} - \frac{1}{t} \geq \frac{1}{6}$ | hyperbolic | never |
| 23. | $(3, s, t)$ where $t > s > 4$ | $\frac{2}{3} - \frac{1}{s} - \frac{1}{t} \geq \frac{1}{6}$ | hyperbolic | never |
| 24. | (r, s, t) where $t > s > r > 4$ | $1 - \frac{1}{r} - \frac{1}{s} - \frac{1}{t} \geq \frac{1}{4}$ | hyperbolic | never |

Examining this table we see that we can eliminate cases 13, 17, 18, 22, 23 and 24 since $\frac{1}{6} - \frac{1}{d}$ will always be less than $1 - 1/r - 1/s - 1/t$. We can also eliminate cases 1, 2, 3, 5 and 19 since these triples are not hyperbolic. Now notice that cases 7, ..., 12 need never be considered since if there are such triples generating $PSl_2(p)$ then there will also be a (3, 3, 4) triple generating $PSl_2(p)$, in which case the genus calculation from the (3, 3, 4) case is at least as small. In a similar fashion we can ignore cases 15, 16 and 21 by comparing them with case 14. Finally, we can use Lemma (2.3) to eliminate case 4. The triples remaining after this will be (2, 3, p), (2, 3, d), (2, 5, 5), (2, 4, 5) and (3, 3, 4). Minimization of the genera for these triples leads directly to the corollary in the introduction.

REFERENCES

- [B] BURNSIDE, W. *Theory of Groups of Finite Order*. Cambridge University Press, 1911.
- [FLM] FRENKEL, I., J. LEPOWSKY and A. MEURMAN. A natural representation of the Fischer-Griess Monster with the modular function J as character (preprint).
- [Gr] GREENBERG, L. Maximal groups and signatures. *Annals of Math. Studies* 79 (1974), 207-226.
- [Gu] GUNNING, R. C. *Lectures on Modular Forms*. Annals of Math. Studies No. 48, Princeton University Press, 1962.
- [GS] GLOVER, H. and D. SJERVE. $PSl_2(p)$ as the automorphism group of a Riemann surface (in preparation).
- [H] HURWITZ, A. Über algebraische Gebilde mit eindeutigen Transformationen in sich. *Math. Ann.* 41 (1892), 403-442.
- [M] MAGNUS, W. *Noneuclidean Tesselations and Their Groups*. Academic Press, 1974.
- [N] NEWMANN, M. *Integral Matrices*. Academic Press, 1972.
- [S] SUZUKI, M. *Group Theory I*. Springer-Verlag 1982.
- [T] TUCKER, T. Finite groups Acting on Surfaces and the genus of a group. *Journal of Combinatorial Theory B* 34 (1983), 82-92.

(Reçu le 17 octobre 1984)

Henry Glover

Ohio State University
Columbus, OH 43210

Denis Sjerve

University of British Columbia
Vancouver, BC V6T1W5