

VIII. Appendix: Extending the Main Theorem to the class E OF NORM-TOPOLOGICAL SPACES

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **31 (1985)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **26.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

Remark 30. By Theorem 28 the isometry type of a definite space with admissible topology is characterized by the sequence $(\langle e_i \rangle)_{i \in \mathbb{N}}$ where $(e_i)_{i \in \mathbb{N}}$ is a maximal orthogonal family in \mathfrak{E} . Conversely, for each $(\alpha_i) \in k^{\mathbb{N}}$ there is a definite space \mathfrak{E} with $L_c(\mathfrak{E}) = L_s(\mathfrak{E})$ admitting a maximal orthogonal family $(e_i)_{i \in \mathbb{N}}$ with $\langle e_i \rangle = \alpha_i$ ($i \in \mathbb{N}$) provided that

- (A) $\xi_i := \varphi \alpha_i \in \Gamma$ satisfies the (type-) condition expressed in (8)
- (B) The form $\langle \cdot, \cdot \rangle$ defined on $\mathfrak{F} := k(e_i)_{i \in \mathbb{N}}$ by $\langle e_i, e_j \rangle = 0$ ($i \neq j$), $\langle e_i \rangle = \alpha_i$ ($i \in \mathbb{N}$) is definite.

These two conditions are implemented by many fields. In order to satisfy (A) one may, e.g. pick fields of generalized formal power series that are complete under a valuation φ with group Γ a prescribed Hahn product [30, p. 31] with sufficiently many factors not 2-divisible, e.g. $\Gamma = \mathbf{Z}^{(\mathbb{N})}$ ordered antilexicographically. Let k be any field with (A) and $t \in \Gamma/2\Gamma$; set $\mathfrak{F}_t = \{\text{span } e_i \mid \varphi \alpha_i + 2\Gamma = t\}$. By (A) $\dim \mathfrak{F}_t < \infty$; furthermore

$$\mathfrak{F} = \bigoplus^\perp \{\mathfrak{F}_t \mid t \in \Gamma/2\Gamma\}.$$

In order to check whether the form $\langle \cdot, \cdot \rangle$ satisfies the triangle inequality on \mathfrak{F} it suffices to verify said inequality on each \mathfrak{F}_t . A. Fässler has given a handy criterium for $\langle \cdot, \cdot \rangle$ to be definite if Hahnproducts Γ are used, as indicated, to construct k with (A), [6, Lemma 15, 16].

VIII. APPENDIX: EXTENDING THE MAIN THEOREM TO THE CLASS \mathcal{E} OF NORM-TOPOLOGICAL SPACES

The arguments applied to the spaces in the class \mathcal{D} can be extended to a larger class \mathcal{E} . First we have (cf. Definition 15):

Definition 31. An infinite dimensional anisotropic quadratic space $(\mathfrak{E}; \langle \cdot, \cdot \rangle)$ over a *-valued field $(k, *, \varphi, \Gamma)$ is called norm-topological if the sets $\mathfrak{U}_\gamma := \{x \in \mathfrak{E} \mid \varphi \langle x \rangle > \gamma\}$ form a 0-neighbourhood basis of a vector space topology on \mathfrak{E} . Let \mathcal{E} be the class of all norm-topological spaces.

Definite spaces are norm-topological, obviously.

A proper subgroup Δ of Γ is *convex* (or isolated) if “ $0 \leqslant x \leqslant y \& y \in \Delta$ ” implies “ $x \in \Delta$ ”. If the subgroup $\Delta \subset \Gamma$ is convex then the factor group Γ/Δ is ordered by setting $\gamma + \Delta \leqslant \delta + \Delta$ iff $\gamma < \delta$ or $\gamma - \delta \in \Delta$; furthermore, $\varphi_\Delta: k \rightarrow \Gamma/\Delta \cup \{\infty\}$ defined by $\varphi_\Delta(\alpha) = \varphi(\alpha) + \Delta$ is a valuation (a “coarser valuation”) which yields the same topology on k as φ .

In order to make the mechanism of types work in the context of norm-topological spaces, i.e., in order to salvage the statement of Corollary 26 in the new context, the concept of type has to be coarsened as follows. For $\gamma \in \Gamma$ we introduce

$$(12) \quad \Delta(\gamma) := \{\delta \in \Gamma \mid \forall n \in \mathbf{N} : n |\delta| \leq |\gamma|\}$$

and

$$(13) \quad \Theta(\gamma) := \bigcap_{\delta \in \Gamma} \Delta(\gamma + 2\delta)$$

If $\gamma \neq 0$ then $\Delta(\gamma)$ is the largest convex subgroup of Γ not containing γ ([21]).

Remark 32. The group defined in (13) for $\gamma = \varphi\langle e \rangle$, $e \in \mathfrak{E}$, represents yet another possibility to introduce a “type” for the vectors in a definite space. The fundamental property expressed in Lemma 25 can be replaced and reproved (along the same lines), cf. [21]:

(14) If U is a convex subgroup in Γ and $(e_i)_{\mathbf{N}}$, $(f_i)_{\mathbf{N}}$ are two maximal orthogonal families in a norm-topological space that satisfies (iii) in Theorem 28 then

$$\text{card } \{i \in I \mid \Theta(\varphi\langle e_i \rangle) \subset U\}_{\neq} = \text{card } \{j \in \mathbf{N} \mid \Theta(\varphi\langle f_j \rangle) \subset U\}_{\neq}.$$

One has the following analogue of Lemma 14:

LEMMA 33. ([21]). *Let $(\mathfrak{E}; \langle \cdot, \cdot \rangle; \varphi, \Gamma, *)$ be a norm-topological space and $\varphi(2) = 0$ (cf. Remark 35 below). Then there is a valuation $\tilde{\varphi} : k \rightarrow \tilde{\Gamma} \cup \{\infty\}$ coarser than φ such that the following holds: Either $(\mathfrak{E}; \langle \cdot, \cdot \rangle; \tilde{\varphi}, \tilde{\Gamma}, *)$ is a definite space, in the sense of Definition 15, or else there are no analytically nilpotent elements $\alpha \in k$ (i.e., for no $\alpha \neq 0$ shall we have $\lim_{\mathbf{N}} \alpha^n = 0$) and then the following weakened versions of the statements in Lemma 14 hold:*

- (i)' $\tilde{\varphi}_{\Lambda}\langle x + y \rangle \geq \min\{\tilde{\varphi}_{\Lambda}\langle x \rangle, \tilde{\varphi}_{\Lambda}\langle y \rangle\}$
- (ii)' $\tilde{\varphi}\langle x \rangle \leq \tilde{\varphi}\langle y \rangle \& \langle x, y \rangle = 0 \Rightarrow \tilde{\varphi}_{\Lambda}\langle x \rangle = \tilde{\varphi}_{\Lambda}\langle x + y \rangle$
- (iii)' $\tilde{\varphi}\langle x, y \rangle \geq \min\{\tilde{\varphi}_{\Lambda}\langle x \rangle, \tilde{\varphi}_{\Lambda}\langle y \rangle\}$
- (iv)' $2\tilde{\varphi}_{\Lambda}\langle x, y \rangle \geq \tilde{\varphi}_{\Lambda}\langle x \rangle + \tilde{\varphi}_{\Lambda}\langle y \rangle$
where $\Lambda = \Theta(\tilde{\varphi}\langle x \rangle)$ and $\Delta = \Theta(\tilde{\varphi}\langle x \rangle) \cap \Theta(\tilde{\varphi}\langle y \rangle)$.

The inequalities in Lemma 33 suffice to salvage all results proved previously on definite spaces; in particular we have the following strengthening of Theorem 28 (cf. Remark 35 below):

THEOREM 34 [21]. *Let \mathfrak{E} be a norm-topological space in the sense of Definition 31 and assume $\varphi(2) = 0$. Then the statements (i), (ii), (iii) in Theorem 28 are equivalent.*

Remark 35. In Definition 15, Lemma 33 and in Theorem 34 we stipulated that $\varphi(2) = 0$ for the valuation φ of the base field. However, it is neither necessary to assume this nor that $\text{char } k$ be different from two. As technicalities increase if 2 is not a unit for φ the general case has been banned from this elementary survey. Refer to [21].

IX. APPENDIX: ORTHOMODULAR SPACES OVER ORDERED FIELDS

A Baer order of a *-field k is a subset $\Pi \subset S := \{\alpha \in k \mid \alpha = \alpha^*\}$ with $1 \in \Pi$, $0 \notin \Pi$, $\Pi + \Pi \subset \Pi$, $\forall \alpha \neq 0 : \alpha \Pi \alpha^* \subset \Pi$, $-\Pi \cup \Pi = S \setminus \{0\}$. ([14]). The map $\alpha \mapsto \alpha^* \alpha = : \|\alpha\|$ has the properties of a norm and defines a topology on k ; if * is continuous then k is a topological *-field [14, Theorem 4.1, p. 231]. The theory of positive definite orthomodular spaces over archimedean ordered fields is settled in [9]: There are but the classical Hilbert spaces over $\mathbf{R}, \mathbf{C}, \mathbf{H}$. If the order is non-archimedean we shall assume that

(15) the subgroup S generated by all $\alpha^* \alpha^{-1}$ is bounded.

There is [14, Sec. 4.5, p. 234] a valuation on k that induces the norm-topology. We remark that the boundness condition on S is always satisfied for the usual orderings on commutative fields, for Prestel's semi-orderings and for all *-ordered fields that are known hitherto.

A family $(e_i)_{i \in I}$ of vectors in a *positive definite space* $(\mathfrak{E}; \langle \cdot, \cdot \rangle)$ over an ordered *-field k is said to satisfy the *type condition* (cf. Definition 21) iff for all $(\alpha_i)_{i \in I} \in k^I$ the following holds: if $(\langle \alpha_i e_i \rangle)_{i \in I}$ is bounded then $(\alpha_i e_i)_{i \in I}$ converges to $0 \in \mathfrak{E}$.

With this version of type condition we have

THEOREM 36. *Let $(\mathfrak{E}; \langle \cdot, \cdot \rangle)$ be a positive definite space over a non-archimedean ordered *-field that satisfies (15). Then the statements (i), (ii), (iii) in Theorem 28 are equivalent.*