

# §11. Application to the b-function (see [SKKO])

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **32 (1986)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

is an isomorphism.

In particular if  $\text{Supp } \mathcal{M} \subset \Lambda_1 \cup \Lambda_2$  and if  $\dim(\Lambda_1 \cap \Lambda_2) \leq n-2$ , then  $\mathcal{M}$  is a direct sum of two holonomic  $\mathcal{E}_X$ -modules supported on  $\Lambda_1$  and  $\Lambda_2$ , respectively.

Here is another type of theorem.

**THEOREM 10.4.3 ([SKKO]).** Let  $\mathcal{M} = \mathcal{E}u = \mathcal{E}/\mathcal{J}$  be a holonomic  $\mathcal{E}$ -module defined on a neighborhood of  $p \in T^*X$ . Assume  $\text{Supp } \mathcal{M} = \Lambda_1 \cup \Lambda_2$  and

- (i)  $\Lambda_1, \Lambda_2$  and  $\Lambda_1 \cap \Lambda_2$  are non-singular and  $\dim \Lambda_1 = \dim \Lambda_2 = n$ ,  $\dim(\Lambda_1 \cap \Lambda_2) = n-1$ .
- (ii)  $T_{p'} \Lambda_1 \cap T_{p'} \Lambda_2 = T_{p'}(\Lambda_1 \cap \Lambda_2)$  for any  $p'$  in a neighborhood of  $p$  in  $\Lambda_1 \cap \Lambda_2$ .
- (iii) The symbol ideal of  $\mathcal{J}$  coincides with the ideal of functions vanishing on  $\Lambda_1 \cup \Lambda_2$ .

Setting  $k = \text{ord}_{\Lambda_1} u - \text{ord}_{\Lambda_2} u - 1/2$ , we have

- (a)  $\mathcal{M}$  has a non-zero quotient supported on  $\Lambda_1 \Leftrightarrow \mathcal{M}$  has a non-zero submodule supported on  $\Lambda_2 \Leftrightarrow k \in \mathbf{Z}$ .
- (b)  $\mathcal{M}_p$  is a simple  $\mathcal{E}_p$ -module  $\Leftrightarrow k \notin \mathbf{Z}$ .

*Sketch of the proof.* By a quantized contact transformation, we can transform  $p, \Lambda_1, \Lambda_2$  and  $\mathcal{J}$  as follows:

$$\begin{aligned} p &= (0, dx_1) \\ \Lambda_1 &= \{(x, \xi); x_1 = \xi_2 = \dots = \xi_n = 0\} \\ \Lambda_2 &= \{(x, \xi); x_1 = x_2 = \xi_3 = \dots = \xi_n = 0\} \\ \mathcal{J} &= \mathcal{E}(x_1 \partial_1 - \lambda) + \mathcal{E}(x_2 \partial_2 - \mu) + \sum_{j>2} \mathcal{E} \partial_j \end{aligned}$$

In this case, we can easily check the theorem.

## § 11. APPLICATION TO THE $b$ -FUNCTION (see [SKKO])

11.1. As one of the most successful application of microlocal analysis, we shall sketch here how to calculate the  $b$ -function of a function under certain conditions.

11.2. Let  $f$  be a holomorphic function on a complex manifold  $X$ . Then, it is proved ([Bj], [Be] [K1]) that there exist (locally) a non zero polynomial  $b(s)$  and  $P(s) \in \mathcal{D}[s] = \mathcal{D} \underset{\text{def}}{\otimes} \mathbf{C}[s]$  such that  $P(s)f(x)^{s+1} = b(s)f(x)^s$

for any  $s \in \mathbf{N}$ . Such a polynomial  $b(s)$  of smallest degree is called the  $b$ -function of  $f(x)$  and is denoted by  $b_f(s)$ . For the relations between the  $b$ -function and the local monodromy see [M1], [K3].

11.3. Set  $\mathcal{J} = \{P(s) \in \mathcal{D}[s]; P(s)f^s = 0 \text{ for } s \in \mathbf{N}\}$  and  $\mathcal{N} = \mathcal{D}[s]/\mathcal{J}$ . We shall denote the canonical generator of  $\mathcal{N}$  by  $f^s$ . Then  $t: \mathcal{N} \ni P(s)f^s \rightarrow P(s+1)f \cdot f^s \in \mathcal{N}$  gives a  $\mathcal{D}$ -endomorphism of  $\mathcal{N}$  and  $t\mathcal{N} = \mathcal{D}[s]f^{s+1}$ . Here  $f^{s+1} = f \cdot f^s \in \mathcal{N}$ . In this terminology  $b_f(s)$  is the minimal polynomial of  $s \in \mathcal{E}nd_{\mathcal{D}}(\mathcal{N}/t\mathcal{N})$ .

For  $\lambda \in \mathbf{C}$ , we set  $\mathcal{M}_\lambda = \mathcal{D}[s]/(\mathcal{J} + \mathcal{D}[s](s-\lambda))$  and denote by  $f^\lambda$  the canonical generator of  $\mathcal{M}_\lambda$ . Then  $f^{\lambda+1} \mapsto f f^\lambda$  defines a  $\mathcal{D}$ -linear homomorphism  $\mathcal{M}_{\lambda+1} \rightarrow \mathcal{M}_\lambda$ .

11.4. Let  $W$  be the closure of

$$\{(s, x, \xi) \in \mathbf{C} \times T^*X; \xi = sd \log f(x), f(x) \neq 0\}$$

in  $\mathbf{C} \times T^*X$ . Set  $W_0 = W \cap \{s=0\} \subset T^*X$ . Then we can prove

**PROPOSITION 11.4.1** ([K1]).

- (i)  $N$  is a coherent  $\mathcal{D}_X$ -module and  $\text{Ch}(N) = p(W)$ , where  $p$  is the projection from  $\mathbf{C} \times T^*X$  to  $T^*X$ .
- (ii) For any  $\lambda \in \mathbf{C}$ ,  $\mathcal{M}_\lambda$  is a regular holonomic  $\mathcal{D}_X$ -module and  $\text{Ch}(\mathcal{M}_\lambda) = W_0$ .
- (iii)  $\mathcal{N}/t\mathcal{N}$  is a regular holonomic  $\mathcal{D}_X$ -module and  $\text{Ch}(\mathcal{N}/t\mathcal{N}) = W_0 \cap (\pi \circ f)^{-1}(0)$ .

11.5. In the sequel, for the sake of simplicity, we assume that there exists a vector field  $v$  such that  $v(f) = f$ . Therefore we have  $v^k(f^s) = s^k f^s$ . Hence  $\mathcal{N}$  is a  $\mathcal{D}$ -module generated by  $f^s$ . If we set  $\tilde{\mathcal{J}} = \mathcal{D} \cap \mathcal{J}$ , then  $\mathcal{N} \cong \mathcal{D}/\mathcal{J}$  and  $\mathcal{J} = \mathcal{D}[s](s-v) + \mathcal{D}[s]\tilde{\mathcal{J}}$ .

11.6. The following lemma is almost obvious but affords a fundamental tool to calculate the  $b$ -function.

LEMMA 11.6.1. Let  $\mathcal{L}$  be an  $\mathcal{E}_X$ -module and  $w$  a non-zero section of  $\mathcal{L}$ . For  $\lambda \in \mathbf{C}$ , we assume

- (i)  $v(w) = \lambda w$
- (ii)  $\tilde{\mathcal{J}}w = 0$
- (iii)  $fw = 0$ .

Then we have  $b_f(\lambda) = 0$ .

*Proof.* There is a  $P \in \mathcal{D}$  such that  $b_f(s)f^s = Pf^{s+1}$ . Hence  $(b_f(v) - Pf)f^s = 0$ , which implies  $b_f(v) - Pf \in \tilde{\mathcal{J}}$ . Since  $b_f(v)w = b_f(\lambda)w$  we have

$$0 = (b_f(v) - Pf)w = b_f(\lambda)w.$$

This implies  $b_f(\lambda) = 0$ .

11.7. Let  $\bar{\mathcal{J}}$  be the symbol ideal of  $\tilde{\mathcal{J}}$ . Then the zero set of  $\bar{\mathcal{J}}$  is  $W$ , and the zero of  $\bar{\mathcal{J}} + \mathcal{O}\sigma(v)$  is  $W_0$ . Let  $\Lambda$  be an irreducible component of  $W_0$ . If  $\bar{\mathcal{J}} + \mathcal{O}_{T^*X}\sigma(v)$  is a reduced ideal at a generic point  $p$  of  $\Lambda$  then we call  $\Lambda$  a *good Lagrangean*.

If  $\Lambda$  is a good Lagrangean, then  $W$  is non-singular on a neighborhood of a generic point  $p$  of  $\Lambda$  and  $\sigma = \sigma(s)|_W$  has non zero-differential. Let  $p: W \rightarrow X$  denote the projection. We define  $m(\Lambda) \in \mathbf{N}$  as the degree of zero of  $f \circ p$  along  $\Lambda$ , and set  $f_\Lambda = (f \circ p / \sigma^{m(\Lambda)})|_\Lambda$ . Let  $\omega$  be the non-vanishing  $n$ -form on  $X$ . Then  $(p^*\omega) \wedge d\sigma$  is an  $(n+1)$ -form on  $W$ . Let  $\mu(\Lambda)$  be the degree of zeros of  $(p^*\omega) \wedge d\sigma$  along  $\Lambda$ , and let  $\eta$  be the  $n$ -form on  $\Lambda$  given by

$$\frac{p^*\omega \wedge d\sigma}{\sigma^{\mu(\Lambda)}} \Big|_\Lambda = \eta \wedge d\sigma.$$

If we set  $\kappa_\Lambda = \eta \otimes \omega^{\otimes -1} \in \omega_\Lambda \otimes \omega_X^{\otimes -1}$ , then this is independent of the choice of  $\omega$ . We have

PROPOSITION 11.7.1 ([SKKO]). *If  $\Lambda$  is a good Lagrangean, then for any  $\lambda \in \mathbf{C}$ ,  $\mathcal{M}_\lambda$  is a simple holonomic system on a neighborhood of a generic point  $p$  of  $\Lambda$  and we have*

- (i)  $\sigma(f^\lambda) = f_\Lambda^\lambda \sqrt{\kappa_\Lambda}$ .

*In particular*

$$\text{ord } f^\lambda = -m(\Lambda)\lambda - \mu(\Lambda)/2.$$

- (ii) There exists a monic polynomial  $b_\Lambda(s)$  of degree  $m(\Lambda)$  and an invertible micro-differential operator  $P_\Lambda$  of order  $m(\Lambda)$  such that

$$b_\Lambda(s)f^s = P_\Lambda f \cdot f^s \quad \text{in} \quad \mathcal{E} \otimes_{\mathcal{D}} \mathcal{N}$$

and

$$\sigma(P_\Lambda)|_\Lambda = f_\Lambda^{-1}.$$

Remark that  $f_\Lambda$  and  $\omega_\Lambda$  are homogeneous of degree  $-m(\Lambda)$  and  $-\mu(\Lambda)$ , respectively.

Remark also that the minimal polynomial of  $s \in \operatorname{End}_{\mathcal{E}}(\mathcal{E} \otimes_{\mathcal{D}} \mathcal{N}/t\mathcal{N})|_\Lambda$  is  $b_\Lambda(s)$ . In fact, if  $Pf^{s+1} = b(s)f^s$  in  $\mathcal{E} \otimes \mathcal{N}$ , then  $(P \cdot P_\Lambda^{-1}b_\Lambda(s) - b(s))f^s = 0$ . This implies that  $P \cdot P_\Lambda^{-1}b_\Lambda(v) - b(v) \in \mathcal{E}\tilde{\mathcal{J}}$ . Hence

$$\sigma(P \cdot P_\Lambda^{-1}b_\Lambda(v) - b(v))|_W = 0.$$

If  $\operatorname{ord} P \cdot P_\Lambda^{-1}b_\Lambda(v) = \operatorname{ord} P > \deg b$ , then  $\sigma(P)|_W = 0$ . Therefore  $P = P' + P''$  with  $P'' \in \mathcal{E}\tilde{\mathcal{J}}$  and  $\sigma(P') < \sigma(P)$ . Hence  $P'f^{s+1} = b(s)f^s$ . Thus, we may assume  $\operatorname{ord} P \leq \deg b$ . Then

$$0 = \sigma(b(v) - P \cdot P_\Lambda^{-1}b_\Lambda(v))|_W = b(\sigma) - (\sigma(P)|_W f_\Lambda b_\Lambda(\sigma)).$$

This shows that  $b(s)$  is a multiple of  $b_\Lambda(s)$ .

**COROLLARY 11.7.2.** *If every irreducible component of  $W_0$  is good Lagrangean, then  $b_f(s)$  is the least common multiple of the  $b_\Lambda(s)$ .*

11.8. Let  $\Lambda_1$  and  $\Lambda_2$  be two good Lagrangeans. We assume the following conditions for a point  $p \in \Lambda_1 \cap \Lambda_2$ :

- (11.8.1)  $\dim_p \Lambda_1 \cap \Lambda_2 = n-1$  and  $\Lambda_1, \Lambda_2$  and  $\Lambda_1 \cap \Lambda_2$  are non singular on a neighborhood of  $p$ .
- (11.8.2) For any point  $p'$  on a neighborhood of  $p$  in  $\Lambda_1 \cap \Lambda_2$ , we have  $T_{p'}\Lambda_1 \cap T_{p'}\Lambda_2 = T_{p'}(\Lambda_1 \cap \Lambda_2)$ .
- (11.8.3)  $\tilde{\mathcal{J}} + \mathcal{O}\sigma(v)$  coincides with the defining ideal of  $\Lambda_1 \cup \Lambda_2$  with the reduced structure.

In this case we say that  $\Lambda_1$  and  $\Lambda_2$  have a *good intersection*.

We have the following theorem.

**THEOREM 11.7.3.** *Let  $\Lambda_1$  and  $\Lambda_2$  be good Lagrangeans with a good intersection. If  $m(\Lambda_1) \geq m(\Lambda_2)$ , then*

$$\prod_{k=0}^{m(\Lambda_1)-m(\Lambda_2)-1} \left( \text{ord}_{\Lambda_2} f^s - \text{ord}_{\Lambda_1} f^s + \frac{1}{2} + k \right) \mid b_f(s).$$

In order to prove this let us take  $\lambda \in \mathbf{C}$  such that

$$(11.8.4) \quad \begin{aligned} k &= \text{ord}_{\Lambda_1} f^\lambda - \text{ord}_{\Lambda_2} f^\lambda - 1/2 \in \mathbf{N} \quad \text{and} \\ k' &= \text{ord}_{\Lambda_1} f^{\lambda+1} - \text{ord}_{\Lambda_2} f^{\lambda+1} - 1/2 \in \mathbf{N}. \end{aligned}$$

Recall that

$$k = (m(\Lambda_2) - m(\Lambda_1))\lambda - \frac{1}{2}(\mu(\Lambda_2) - \mu(\Lambda_1) - 1/2)$$

and  $k' = k + (m(\Lambda_2) - m(\Lambda_1))$ . Then by Theorem 10.4.3,  $\mathcal{M}_\lambda$  has a non-zero quotient  $\mathcal{L}$  whose support is  $\Lambda_1$ . Let  $w \in \mathcal{L}$  be the image of  $f^\lambda \in \mathcal{M}_\lambda$ .

Let  $\alpha: \mathcal{M}_\lambda \rightarrow \mathcal{L}$  be the canonical homomorphism and  $\beta: \mathcal{M}_{\lambda+1} \rightarrow \mathcal{M}_\lambda$  be the homomorphism given by  $f^{\lambda+1} \mapsto f \cdot f^\lambda$ . Then, since  $k' \notin \mathbf{N}$ ,  $\mathcal{M}_{\lambda+1}$  has no non-zero quotient supported in  $\Lambda_1$ . Hence  $\alpha \circ \beta = 0$ . Therefore  $fw = \alpha\beta(f^{\lambda+1}) = 0$ . Thus we can apply Lemma 11.6.1 to conclude that  $b_f(\lambda) = 0$ . If  $k \in \mathbf{Z}$  with  $0 \leq k < m(\Lambda_1) - m(\Lambda_2)$  then

$$\lambda = \frac{1}{m(\Lambda_1) - m(\Lambda_2)} \left( k + \frac{1}{2}(\mu(\Lambda_1) - \mu(\Lambda_2) - 1) \right)$$

satisfies (11.8.4). This shows that  $b_f(s)$  is a multiple of

$$\begin{aligned} &\prod_{k=0}^{m(\Lambda_1)-m(\Lambda_2)-1} \left( (m(\Lambda_1) - m(\Lambda_2))s - \frac{1}{2}(\mu(\Lambda_1) - \mu(\Lambda_2) - 1) + k \right) \\ &= \text{const.} \prod_{k=0}^{m(\Lambda_1)-m(\Lambda_2)-1} \left( \text{ord}_{\Lambda_2} f^s - \text{ord}_{\Lambda_1} f^s + \frac{1}{2} + k \right). \end{aligned}$$


If we refine this argument, we can prove

**THEOREM 11.8.2 ([SKKO]).** *If  $\Lambda_1$  and  $\Lambda_2$  are good Lagrangeans with a good intersection and if  $m(\Lambda_1) \geq m(\Lambda_2)$  then*

$$\frac{b_{\Lambda_1}(s)}{b_{\Lambda_2}(s)} = \text{const.} \prod_{k=0}^{m(\Lambda_1)-m(\Lambda_2)-1} \left( \text{ord}_{\Lambda_2} f^s - \text{ord}_{\Lambda_1} f^s + \frac{1}{2} + k \right).$$

*Example 11.8.3.*


$$(i) \quad X = M_n(\mathbf{C}) = \mathbf{C}^{n^2} \quad \text{and} \quad f(x) = \det x.$$



Here  $\circlearrowleft$  means a good Lagrangean which is the conormal bundle to an  $a$ -codimensional submanifold.  $\circlearrowleft - \circlearrowleft$  means that the two corresponding good Lagrangeans have a good intersection.

The polynomial attached to the intersection is the ratio of the corresponding  $b_\Lambda$ -functions, calculated by Theorem 11.8.2. The polynomial attached to the circle is the order of  $f^\lambda$ .

$$(ii) \quad X = \mathbf{C}^n, f(x) = x_1^2 + \dots + x_n^2$$



$$(iii) \quad X = \mathbf{C}^3 \quad f = x^2y + z^2$$

