

TREES, TAIL WAGGING AND GROUP PRESENTATIONS

Autor(en): **Armstrong, M. A.**

Objekttyp: **Article**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **32 (1986)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **26.05.2024**

Persistenter Link: <https://doi.org/10.5169/seals-55090>

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

TREES, TAIL WAGGING AND GROUP PRESENTATIONS

by M. A. ARMSTRONG

The Bass-Serre theorem gives a presentation for a group of automorphisms of a tree. Like all good theorems it has attracted considerable attention and there are now several proofs available [4], [3], [1]. Our goal is a natural elementary proof which makes maximal use of the geometry of the tree.

1. GRAPHS

A graph X consists of two sets E (directed edges) and V (vertices) and two functions

$$\begin{aligned} E &\rightarrow E, & e &\mapsto \bar{e} \\ E &\rightarrow V \times V, & e &\mapsto (i(e), t(e)) \end{aligned}$$

which satisfy $\bar{\bar{e}} = e$, $\bar{e} \neq e$ and $i(\bar{e}) = t(e)$ for each $e \in E$. The vertices $i(e)$, $t(e)$ are the initial and terminal vertices of the directed edge e , and \bar{e} is the reverse of e . Henceforth we refer to directed edges simply as edges.

A path in X joining vertex u to vertex v is an ordered string of edges $e_1 e_2 \dots e_n$ such that $i(e_1) = u$, $i(e_{k+1}) = t(e_k)$ for $1 \leq k \leq n-1$, and $t(e_n) = v$. If $v = u$ we have a circuit. A path of the form $e\bar{e}$ is a *round trip* and a circuit which does not contain any round trips will be called a loop. If any two distinct vertices may be joined by a path then the graph is connected. A *tree* is a connected graph which does not contain any loops.

Let X be a tree. A path in X is a *geodesic* if it does not contain any round trips. Given distinct vertices u, v of X there is a *unique* geodesic \overrightarrow{uv} which joins u to v .

An action of a group G on a graph X is an action of G on E and on V such that $g\bar{e} = \overline{ge}$, $i(ge) = gi(e)$, $t(ge) = gt(e)$ and $ge \neq \bar{e}$ for each $e \in E$. Because group elements are not allowed to reverse edges we have a

quotient graph X/G . When G acts on X we shall often say that G is a group of automorphisms of X .

We adopt the usual notation whereby G_x denotes the stabilizer of a vertex x . If $g \in G$ happens to fix x we write g_x for the element g thought of as a member of G_x . Of course G_e denotes the stabilizer of the edge e . If x is a vertex of e then G_e is a subgroup of G_x .

Suppose G acts on a tree X . If $g \in G$ fixes the vertices u, v then it must fix the whole geodesic \overrightarrow{uv} , since otherwise the image of \overrightarrow{uv} under g would be a second geodesic from u to v .

2. LIFTING EDGES

Let G be a group of automorphisms of a tree X . Choose a maximal tree M in X/G and lift it [4, Proposition I.14] to a subtree T of X . The vertices of T form a set of representatives for the action of G on the vertices of X . For each pair of edges f, \bar{f} from $X/G - M$ select one, say f , and lift it to an edge e of X which has its initial vertex x in T . Exactly one vertex z of T lies in the same orbit as $t(e)$ and we choose an element γ_f from G that maps z onto $t(e)$. We can now lift \bar{f} to $(\gamma_f)^{-1}\bar{e}$. This has its initial vertex z in T and $\gamma_{\bar{f}} = (\gamma_f)^{-1}$ sends the vertex x of T to its terminal vertex (Figure 1). Finally we extend the correspondence $f \rightarrow \gamma_f$ over the edges of M by setting $\gamma_f = 1$ (the identity element of G) whenever $f \in M$.

The Bass-Serre theorem [4, Theorem I.13] gives the following presentation for G .

- (a) *Generators.* The elements of all the G_w where w is a vertex of T and the γ_f where f is an edge of X/G .
- (b) *Relations.* The internal relations of each stabilizer G_w together with

$$\gamma_f = 1 \text{ if } f \text{ is an edge of } M,$$

$$\gamma_{\bar{f}} = (\gamma_f)^{-1} \text{ and}$$

$$\gamma_{\bar{f}} g_x \gamma_f = (\gamma_{\bar{f}} g \gamma_f)_z \text{ where } e \text{ is the chosen lift of } f \text{ and } g \in G_e.$$

(If f is an edge of M then $z = t(e)$ and the final relation reduces to $g_x = g_z$ whenever $g \in G_e$).




FIGURE 1

3. TAIL WAGGING

With the notation established above let $*G_w$ denote the free product of the stabilizers of the vertices of T , and F the free group generated by symbols λ_f , one for each edge f of X/G . Let R be the normal consequence in $(*G_w)*F$ of the words

$$\begin{aligned} \lambda_f & \quad (f \text{ an edge of } M), \\ \lambda_{\bar{f}} \lambda_f & \quad \text{and} \\ \lambda_{\bar{f}} g_x \lambda_f (\gamma_{\bar{f}} g \gamma_f)_z^{-1} \end{aligned}$$

We shall produce an isomorphism

$$\psi: G \rightarrow [(*G_w)*F]/R.$$

Choose a vertex v of T as base point. If $g \in G$ fixes v set

$$\psi(g) = g_v R$$

where as usual g_v is the element g interpreted as a member of G_v . If g moves v then it sends it outside T because no two vertices of T lie in the same orbit. Let $e_1 e_2 \dots e_n$ be the geodesic which joins v to gv and suppose e_m is the first edge that is *not* in T . The path $e_m e_{m+1} \dots e_n$ will be called the *tail* of $\overrightarrow{v gv}$. Let x_1 be the initial vertex of e_m . Project e_m into X/G to give an edge f_1 . The canonical lift e^1 of f_1 into X has its initial vertex in T , so $i(e^1) = x_1$. Choose an element $a_{x_1} \in G_{x_1}$ which sends e^1 to e_m . Let

$$e_k^1 = (\gamma_{f_1} a_{x_1}^{-1}) e_k$$

for $m+1 \leq k \leq n$, and replace $e_1 e_2 \dots e_n$ by the new path $e_{m+1}^1 e_{m+2}^1 \dots e_n^1$. We call this process *tail wagging*. Our new path begins at

$$z_1 = t(\gamma_{f_1} e^1) = i(e_{m+1}^1)$$

which is a vertex of T and ends at $(\gamma_{f_1} a_{x_1}^{-1} g)v$, see Figure 2. We walk along it to the first point x_2 where it quits T and repeat the above




FIGURE 2

procedure. Since we shorten the tail at each step we eventually obtain a path which lies entirely in T and ends at say

$$(\gamma_{\bar{f}_r} a_{x_r}^{-1} \dots \gamma_{\bar{f}_2} a_{x_2}^{-1} \gamma_{\bar{f}_1} a_{x_1}^{-1} g)v.$$

Then $\gamma_{\bar{f}_r} a_{x_r}^{-1} \dots \gamma_{\bar{f}_1} a_{x_1}^{-1} g$ must fix v , say $\gamma_{\bar{f}_r} a_{x_r}^{-1} \dots \gamma_{\bar{f}_1} a_{x_1}^{-1} g = a_v \in G_v$.

We now have

$$g = a_{x_1} \gamma_{f_1} \dots a_{x_r} \gamma_{f_r} a_v$$

and we somewhat optimistically define

$$\psi(g) = a_{x_1} \lambda_{f_1} \dots a_{x_r} \lambda_{f_r} a_v R.$$

4. AN INEFFICIENT CHOICE

Is ψ well defined? The geodesic from v to gv is certainly unique, as is the first point x_1 where it leaves T and its first edge e_m outside T . Both the edge e^1 and the group element γ_{f_1} are now determined by our original construction. The only ambiguity at this stage is the choice of the element $a_{x_1} \in G_{x_1}$ which maps e^1 to e_m . A different choice b_{x_1} will give a path from z_1 to $(\gamma_{\bar{f}_1} b_{x_1}^{-1} g)v$ which leaves T for the first time at say y_2 . The first edge outside T will project to an edge f'_2 of X/G and so on until eventually we have g expressed as

$$g = b_{x_1} \gamma_{f_1} b_{y_2} \gamma_{f'_2} \dots b_{y_s} \gamma_{f'_s} b_v.$$

We must show that $a_{x_1} \lambda_{f_1} a_{x_2} \lambda_{f_2} \dots a_{x_r} \lambda_{f_r} a_v$ and $b_{x_1} \lambda_{f_1} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v$ determine the same left coset of R in $(*G_w)*F$.

Agree to select a_{x_1} from G_{x_1} so that the tail of the resulting path is as long as possible. Continue in this way selecting $a_{x_2}, a_{x_3} \dots$ so as to maximise the length of the tail at each stage. We shall compare any other set of choices with this rather inefficient selection.

Both a_{x_1} and b_{x_1} map e^1 to e_m , so $c = a_{x_1}^{-1} b_{x_1}$ must fix e^1 . Also, due to our particular selection of a_{x_1} , the geodesic from z_1 to x_2 is left fixed by $\gamma_{\bar{f}_1} c \gamma_{f_1}$. Therefore

$$\begin{aligned}
& b_{x_1} \lambda_{f_1} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R \\
&= a_{x_1} \lambda_{f_1} \lambda_{\bar{f}_1} a_{x_1}^{-1} b_{x_1} \lambda_{f_1} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R \\
&= a_{x_1} \lambda_{f_1} \lambda_{\bar{f}_1} c_{x_1} \lambda_{f_1} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R \\
&= a_{x_1} \lambda_{f_1} (\gamma_{\bar{f}_1} c \gamma_{f_1})_{x_1} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R \\
&= a_{x_1} \lambda_{f_1} (\gamma_{\bar{f}_1} c \gamma_{f_1})_{x_2} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R \\
&= a_{x_1} \lambda_{f_1} a'_{x_2} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R
\end{aligned}$$

where $a'_{x_2} = (\gamma_{\bar{f}_1} c \gamma_{f_1})_{x_2}$. If x_2 happens to equal y_2 then we simplify this further to

$$a_{x_1} \lambda_{f_1} a''_{x_2} \lambda_{f_2} b_{y_3} \lambda_{f'_3} \dots b_{y_s} \lambda_{f'_s} b_v R$$

where a''_{x_2} is the product $a'_{x_2} b_{y_2}$ in G_{x_2} . We now compare a_{x_2} with a'_{x_2} if $x_2 \neq y_2$, noting that $\gamma_{f_2} = 1$ in this case, or with a''_{x_2} if $x_2 = y_2$, and repeat the process. Eventually we obtain

$$b_{x_1} \lambda_{f_1} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R = a_{x_1} \lambda_{f_1} a_{x_2} \lambda_{f_2} \dots a_{x_r} \lambda_{f_r} a''_v R.$$

As $g = a_{x_1} \gamma_{f_1} \dots a_{x_r} \gamma_{f_r} a_v = a_{x_1} \gamma_{f_1} \dots a_{x_r} \gamma_{f_r} a''_v$ we see that $a''_v = a_v$. This completes the proof that ψ is well defined.

5. NEAREST FIXED POINTS

To show ψ is a homomorphism we shall verify

$$\psi(hg) = \psi(h)\psi(g)$$

under the assumption that h either leaves some vertex of T fixed or is one of the elements γ_f . This is sufficient because the elements of the G_w (w a vertex of T) together with the γ_f (f an edge of $X/G-M$) form a set of generators for G .

Suppose h fixes the vertex w of T . Walk along the geodesic \overrightarrow{vw} and let x be the first vertex we meet which is left fixed by h . Then \overrightarrow{vx} is contained in T , and \overrightarrow{vx} followed by $h(\overrightarrow{xv})$ is the geodesic from v to hv . This quits T for the first time at x and we see that

$$\psi(h) = h_x R.$$




FIGURE 3

Using the geodesic from v to gv we have $\psi(g) = a_{x_1} \lambda_{f_1} \dots a_{x_r} \lambda_{f_r} a_v R$ in the usual way. Therefore

$$\psi(h)\psi(g) = h_x a_{x_1} \lambda_{f_1} \dots a_{x_r} \lambda_{f_r} a_v R.$$

In order to compute $\psi(hg)$ we need the geodesic from v to $(hg)v$. We can construct this as follows, take $\overrightarrow{v hv}$ followed by the image of $\overrightarrow{v gv}$ under h and remove any round trips.

If $\overrightarrow{v gv}$ does not contain all of \overrightarrow{vx} (Figure 3) then $\overrightarrow{v(hg)v}$ leaves T for the first time at x . A tail wag of $\overrightarrow{v(hg)v}$ using h_x^{-1} leads us to a path which has the same tail as $\overrightarrow{v gv}$, then the process continues as for g . Thus

$$\psi(hg) = h_x a_{x_1} \lambda_{f_1} \dots a_{x_r} \lambda_{f_r} a_v R = \psi(h)\psi(g).$$

Otherwise $\overrightarrow{v gv}$ contains all of \overrightarrow{vx} (Figure 4) and we split the argument into three cases.




FIGURE 4

produces a path with the same tail as $\overrightarrow{v g v}$. The process then continues as for g and

$$\psi(\gamma_f g) = \lambda_f a_{x_1} \lambda_{f_1} \dots a_{x_r} \lambda_{f_r} a_v R = \psi(\gamma_f) \psi(g).$$

Otherwise $\overrightarrow{v g v}$ contains $\overrightarrow{v y}$. Then $x_1 = z$, $\gamma_{f_1} = \gamma_{\bar{f}}$ and we may as well take $a_{x_1} = 1$. A first tail wag of $\overrightarrow{v g v}$ using γ_f leaves a path with the same tail as $\overrightarrow{v(\gamma_f g)v}$. Thus

$$\begin{aligned} \psi(\gamma_f g) &= a_{x_2} \lambda_{f_2} \dots a_{x_r} \lambda_{f_r} a_v R \\ &= \lambda_f \lambda_{\bar{f}} a_{x_2} \lambda_{f_2} \dots a_{x_r} \lambda_{f_r} a_v R \\ &= \psi(\gamma_f) \psi(g). \end{aligned}$$

This completes the proof that ψ is a homomorphism.

Our construction of ψ ensures that if $\psi(g) = R$ then $g = 1$. So ψ is injective. The cosets $h_w R$ (w a vertex of T and $h(w) = w$) and $\lambda_f R$ (f an edge of X/G) together generate $[(\ast G_w) \ast F]/R$. Now $\psi(h) = h_x R$ where x is the nearest fixed point of h to v . But h fixes all of \overrightarrow{xw} so

$$\psi(h) = h_x R = h_w R.$$

Also

$$\psi(\gamma_f) = \lambda_f R.$$

Therefore the image of ψ is all of $[(\ast G_w) \ast F]/R$ and we have shown that ψ is an isomorphism.

The author would like to thank the members of the Mathematics Department of the University of Geneva for their hospitality during the preparation of this article.

REFERENCES

- [1] DICKS, W. *Groups Trees and Projective Modules*. Lect. Notes in Math. 790, Springer-Verlag 1980.
- [2] HAUSMANN, J.-C. Sur l'usage de critères pour reconnaître un groupe libre, un produit amalgamé ou une HNN-extension. *L'Enseignement Mathématique* 27 (1981), 221-242.

- [3] SCOTT, G. P. and C. T. C. WALL. Topological methods in group theory. *London Math. Soc. Lect. Notes* 36, Cambridge Univ. Press 1979, 137-203.
- [4] SERRE, J.-P. Arbres, Amalgames, SL_2 . Astérisque 46, Soc. Math. de France 1977.
- [5] STALLINGS, J. R. Topology of finite graphs. *Inventiones Math.* 71 (1983), 551-565.

(Reçu le 12 septembre 1985)

M. A. Armstrong

Mathematics Department

University of Durham

Durham DH1 3LE, England