

5. Nearest fixed points

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **32 (1986)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **26.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

$$\begin{aligned}
& b_{x_1} \lambda_{f_1} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R \\
&= a_{x_1} \lambda_{f_1} \lambda_{\bar{f}_1} a_{x_1}^{-1} b_{x_1} \lambda_{f_1} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R \\
&= a_{x_1} \lambda_{f_1} \lambda_{\bar{f}_1} c_{x_1} \lambda_{f_1} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R \\
&= a_{x_1} \lambda_{f_1} (\gamma_{\bar{f}_1} c \gamma_{f_1})_{z_1} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R \\
&= a_{x_1} \lambda_{f_1} (\gamma_{\bar{f}_1} c \gamma_{f_1})_{x_2} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R \\
&= a_{x_1} \lambda_{f_1} a'_{x_2} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R
\end{aligned}$$

where $a'_{x_2} = (\gamma_{\bar{f}_1} c \gamma_{f_1})_{x_2}$. If x_2 happens to equal y_2 then we simplify this further to

$$a_{x_1} \lambda_{f_1} a''_{x_2} \lambda_{f_2} b_{y_3} \lambda_{f'_3} \dots b_{y_s} \lambda_{f'_s} b_v R$$

where a''_{x_2} is the product $a'_{x_2} b_{y_2}$ in G_{x_2} . We now compare a_{x_2} with a'_{x_2} if $x_2 \neq y_2$, noting that $\gamma_{f_2} = 1$ in this case, or with a''_{x_2} if $x_2 = y_2$, and repeat the process. Eventually we obtain

$$b_{x_1} \lambda_{f_1} b_{y_2} \lambda_{f'_2} \dots b_{y_s} \lambda_{f'_s} b_v R = a_{x_1} \lambda_{f_1} a_{x_2} \lambda_{f_2} \dots a_{x_r} \lambda_{f_r} a''_v R.$$

As $g = a_{x_1} \gamma_{f_1} \dots a_{x_r} \gamma_{f_r} a_v = a_{x_1} \gamma_{f_1} \dots a_{x_r} \gamma_{f_r} a''_v$ we see that $a''_v = a_v$. This completes the proof that ψ is well defined.

5. NEAREST FIXED POINTS

To show ψ is a homomorphism we shall verify

$$\psi(hg) = \psi(h)\psi(g)$$

under the assumption that h either leaves some vertex of T fixed or is one of the elements γ_f . This is sufficient because the elements of the G_w (w a vertex of T) together with the γ_f (f an edge of $X/G - M$) form a set of generators for G .

Suppose h fixes the vertex w of T . Walk along the geodesic \overrightarrow{vw} and let x be the first vertex we meet which is left fixed by h . Then \overrightarrow{vx} is contained in T , and \overrightarrow{vx} followed by $h(\overrightarrow{vx})$ is the geodesic from v to hv . This quits T for the first time at x and we see that

$$\psi(h) = h_x R.$$




FIGURE 3

Using the geodesic from v to gv we have $\psi(g) = a_{x_1} \lambda_{f_1} \dots a_{x_r} \lambda_{f_r} a_v R$ in the usual way. Therefore

$$\psi(h)\psi(g) = h_x a_{x_1} \lambda_{f_1} \dots a_{x_r} \lambda_{f_r} a_v R.$$

In order to compute $\psi(hg)$ we need the geodesic from v to $(hg)v$. We can construct this as follows, take $\overrightarrow{v hv}$ followed by the image of \overrightarrow{vgv} under h and remove any round trips.

If \overrightarrow{vgv} does not contain all of \overrightarrow{vx} (Figure 3) then $\overrightarrow{v(hg)v}$ leaves T for the first time at x . A tail wag of $\overrightarrow{v(hg)v}$ using h_x^{-1} leads us to a path which has the same tail as \overrightarrow{vgv} , then the process continues as for g . Thus

$$\psi(hg) = h_x a_{x_1} \lambda_{f_1} \dots a_{x_r} \lambda_{f_r} a_v R = \psi(h)\psi(g).$$

Otherwise \overrightarrow{vgv} contains all of \overrightarrow{vx} (Figure 4) and we split the argument into three cases.




FIGURE 4

(a) \overrightarrow{vgv} stays in T for at least one more edge after x . Then $\overrightarrow{v(hg)v}$ must leave T at x . As above, a first choice of h_x^{-1} leads to a path with the same tail as \overrightarrow{vgv} .

(b) \overrightarrow{vgv} and $\overrightarrow{v(hg)v}$ both leave T at x . Then $x_1 = x$ and we write a_x instead of a_{x_1} . A first tail wag of $\overrightarrow{v(hg)v}$ using $\gamma_{\bar{f}_1}(h_x a_x)^{-1}$ produces the same path as a first tail wag of \overrightarrow{vgv} using $\gamma_{\bar{f}_1} a_x^{-1}$. Thus

$$\psi(hg) = h_x a_x \lambda_{f_1} a_{x_2} \lambda_{f_2} \dots a_{x_r} \lambda_{f_r} a_v R = \psi(h)\psi(g).$$

(c) \overrightarrow{vgv} leaves T at x , but $\overrightarrow{v(hg)v}$ stays in T for at least one more edge after x . Then $x_1 = x$, $\gamma_{f_1} = 1$ and we may as well equate a_{x_1} with h_x^{-1} . A first tail wag of \overrightarrow{vgv} using h_x gives a path with the same tail as $\overrightarrow{v(hg)v}$. Thus

$$\begin{aligned}\psi(hg) &= a_{x_2} \lambda_{f_2} \dots a_{x_r} \lambda_{f_r} a_v R \\ &= h_x h_x^{-1} a_{x_2} \lambda_{f_2} \dots a_{x_r} \lambda_{f_r} a_v R \\ &= \psi(h)\psi(g).\end{aligned}$$

Suppose finally that $h = \gamma_f$ for some edge f of $X/G - M$. As usual e is the chosen lift of f into X with $x = i(e) \in T$ and $z = t(\gamma_{\bar{f}} e)$. Let $y = i(\gamma_{\bar{f}} e)$. The geodesic from v to $\gamma_f v$ is made up of \overrightarrow{vx} followed by e followed by $\overrightarrow{\gamma_f(zv)}$. This leaves T for the first time at x and a single tail wag using $\gamma_{\bar{f}}$ produces \overrightarrow{zv} . Therefore

$$\psi(\gamma_f) = \lambda_f R.$$

To obtain the geodesic from v to $(\gamma_f g)v$ we follow $\overrightarrow{v\gamma_f v}$ by $\gamma_f \overrightarrow{(vgv)}$ and then remove any round trips (Figure 5). If \overrightarrow{vgv} does not contain \overrightarrow{vy} , then $\overrightarrow{v(\gamma_f g)v}$ leaves T for the first time at x and a single tail wag using $\gamma_{\bar{f}}$




FIGURE 5

produces a path with the same tail as \overrightarrow{vgv} . The process then continues as for g and

$$\psi(\gamma_f g) = \lambda_f a_{x_1} \lambda_{f_1} \dots a_{x_r} \lambda_{f_r} a_v R = \psi(\gamma_f) \psi(g).$$

Otherwise \overrightarrow{vgv} contains \overrightarrow{vy} . Then $x_1 = z$, $\gamma_{f_1} = \gamma_{\bar{f}}$ and we may as well take $a_{x_1} = 1$. A first tail wag of \overrightarrow{vgv} using γ_f leaves a path with the same tail as $\overrightarrow{v(\gamma_f g)v}$. Thus

$$\begin{aligned}\psi(\gamma_f g) &= a_{x_2} \lambda_{f_2} \dots a_{x_r} \lambda_{f_r} a_v R \\ &= \lambda_f \lambda_{\bar{f}} a_{x_2} \lambda_{f_2} \dots a_{x_r} \lambda_{f_r} a_v R \\ &= \psi(\gamma_f) \psi(g).\end{aligned}$$

This completes the proof that ψ is a homomorphism.

Our construction of ψ ensures that if $\psi(g) = R$ then $g = 1$. So ψ is injective. The cosets $h_w R$ (w a vertex of T and $h(w) = w$) and $\lambda_f R$ (f an edge of X/G) together generate $[(*G_w)*F]/R$. Now $\psi(h) = h_x R$ where x is the nearest fixed point of h to v . But h fixes all of \overrightarrow{xw} so

$$\psi(h) = h_x R = h_w R.$$

Also

$$\psi(\gamma_f) = \lambda_f R.$$

Therefore the image of ψ is all of $[(*G_w)*F]/R$ and we have shown that ψ is an isomorphism.

The author would like to thank the members of the Mathematics Department of the University of Geneva for their hospitality during the preparation of this article.

REFERENCES

- [1] DICKS, W. *Groups Trees and Projective Modules*. Lect. Notes in Math. 790, Springer-Verlag 1980.
- [2] HAUSMANN, J.-C. Sur l'usage de critères pour reconnaître un groupe libre, un produit amalgamé ou une HNN-extension. *L'Enseignement Mathématique* 27 (1981), 221-242.