

### **3. Hopf fibrations with fibre $S^3$**

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **32 (1986)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

#### **Haftungsausschluss**

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

- d) *There is no symmetry of  $H$  inducing the identity on the base space and reversing the orientations of the Hopf circles.*

Consider the symmetries  $z \cdot Id$ ,  $|z| = 1$ , which multiply each coordinate in  $C^n$  by the complex number  $z$  of unit length. They induce the identity on the base space, and can be selected to take a fibre to itself by a preassigned rotation, proving a).

The transformations in  $U(n)$  can take any complex line in  $C^n$  to any other by a preassigned orientation preserving rigid motion. Complex conjugation then adds the orientation reversing ones, proving b).

In particular, this implies c).

Suppose there were a symmetry of  $H: S^1 \hookrightarrow S^{2n-1} \rightarrow CP^{n-1}$  taking each Hopf circle to itself with reversal of orientation. Then, by restriction to  $C^2$ , such a symmetry would also exist for  $n = 2$ . Its reversal of orientation on the total space  $S^3$  would then contradict the remark following Proposition 2.1. This proves d). QED

*Remarks.* 1) Note that the existence of symmetries of  $H$  rotating each Hopf circle within itself shows again that these circles must be parallel.

2) Also note that a symmetry of  $H: S^1 \hookrightarrow S^{2n-1} \rightarrow CP^{n-1}$  induces an isometry of the base space  $CP^{n-1}$  in its canonical metric. We remark without proof that *all* isometries of  $CP^{n-1}$  can be produced this way.

### 3. HOPF FIBRATIONS WITH FIBRE $S^3$

Choose orthonormal coordinates in  $R^{4n}$  and identify this space with quaternionic  $n$ -space  $H^n$ . A little care is needed in dealing with  $H^n$  because the quaternions form a *non-commutative* division algebra:

- 1) Scalars  $v \in H$  will act on vectors  $(u_1, \dots, u_n) \in H^n$  from the *right*.

$$(u_1, \dots, u_n) v = (u_1 v, \dots, u_n v).$$

- 2)  $H$ -linear transformations of  $H^n$  will be expressed by matrices of quaternions acting from the *left* (so as to commute with scalar multiplication).

The quaternionic lines in  $H^n$ , each looking like a real 4-plane, form the quaternionic projective space  $HP^{n-1}$  and fill out  $H^n$ , with any two meeting only at the origin. The unit 3-spheres on these quaternionic lines give us the *Hopf fibration*

$$H: S^3 \hookrightarrow S^{4n-1} \rightarrow HP^{n-1}.$$

**PROPOSITION 3.1.** *The Hopf 3-spheres on  $S^{4n-1}$  are parallel to one another.*

The proof is similar to that of Proposition 1.1 for Hopf circles; it uses the fact that scalar multiplication by  $i, j$  and  $k$  are isometries of  $H^n$ . Alternatively, it will follow, as in Remark 1 above, from Proposition 4.2 a.

QED

The Riemannian metric on  $HP^{n-1}$  which makes the Hopf projection  $S^{4n-1} \rightarrow HP^{n-1}$  into a Riemannian submersion is known as the *canonical metric* on  $HP^{n-1}$ . The canonical metric on  $HP^1$  makes it into a round 4-sphere of radius  $1/2$ . This follows by the same argument given in Proposition 1.2 for the case  $H: S^1 \hookrightarrow S^3 \rightarrow CP^1$ .

#### 4. SYMMETRIES OF THE HOPF FIBRATIONS WITH FIBRE $S^3$

We now investigate the symmetries of the Hopf fibration

$$H: S^3 \hookrightarrow S^{4n-1} \rightarrow HP^{n-1}.$$

The symplectic group

$$Sp(n) = Gl(n, H) \cap O(4n)$$

consists of quaternionically linear maps which are also rigid. Since these maps take quaternionic lines to quaternionic lines, they must be symmetries of the above Hopf fibration.

There are other symmetries. For each unit quaternion  $v$ , consider the action of right scalar multiplication by  $v$  on  $H^n$ ,

$$R_v(u_1, \dots, u_n) = (u_1 v, \dots, u_n v).$$

This map is certainly *not*  $H$ -linear, since

$$R_v[(u_1, \dots, u_n)w] = (u_1 w v, \dots, u_n w v),$$

while

$$[R_v(u_1, \dots, u_n)]w = (u_1 v w, \dots, u_n v w).$$

Nevertheless,  $R_v$  takes each quaternionic line in  $H^n$  to itself. Thus the group  $S^3$  of unit quaternions, acting on  $H^n$  from the right, must also be counted among the symmetries of our Hopf fibration.

Since the symplectic group  $Sp(n)$  acts on  $S^{4n-1}$  from the left, while the group  $S^3$  of unit quaternions acts from the right, these two actions