

# 7. WINDING NUMBERS

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **35 (1989)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

## Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

## Haftungsausschluss


Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

## 7. WINDING NUMBERS

Let  $X$  be an  $n$ -dimensional oriented smooth manifold and  $s$  a point of  $X$ . Consider a compact  $n$ -dimensional submanifold with boundary  $B$  with  $s$  as an interior point and put

$$(7.1) \quad \text{Tr}(\omega; s) = \int_{\partial B} \omega, \quad \omega \in \Gamma(X - \{s\}, \Omega^{n-1}), d\omega = 0.$$

This symbol is independent of  $B$  as it follows by considering a small “ball”  $C$  around  $s$  contained in the interior of  $B$




Stokes formula for  $B - C^0$

$$\int_{\partial B} \omega - \int_{\partial C} \omega = \int_{B - C^0} d\omega.$$

Alternatively, choose a compactly supported smooth real function  $\rho_s$  on  $X$  which is constant 1 in a neighbourhood of  $s$ . Then

$$(7.2) \quad \text{Tr}(\omega; s) = (-1)^n \int_X \omega \wedge d\rho_s, \quad \omega \in \Gamma(X - \{s\}, \Omega^{n-1}), d\omega = 0.$$

*Proof.* Choose “balls”  $B$  and  $D$  with center  $s$  such that  $\rho_s$  is constant 1 on  $B$  while  $\text{Supp}(\rho_s)$  is contained in the interior of  $D$ . From Stokes formula we get that



$$\begin{aligned} \int_{\partial D} \rho_s \omega - \int_{\partial B} \rho_s \omega &= \int_{D - B} \rho_s \wedge \omega - \int_{D - B} \rho_s \omega \\ &- \int_{\partial B} \omega = -(-1)^n \int_X \omega \wedge d\rho_s + \int_{D - B} \rho_s d\omega. \end{aligned}$$

Notice that the last terms vanishes in case  $\omega$  is exact.

Q.E.D.

(7.3) *Example.* Let  $E$  denote an oriented  $n$ -dimensional Euclidian space. The distance  $r$  to the origin defines a 1-form  $dr^{2-n}$  on  $E - \{0\}$ . The dual form  $*dr^{2-n}$  in the sense of Hodge is closed with

$$\text{Tr}(*dr^{2-n}; 0) = (2-n)\sigma_{n-1}$$

where  $\sigma_{n-1}$  denotes the area of the unit sphere in  $E$ , compare [3] VII. 1.

Let us interprete (7.1) in terms of de Rham homology. Integration of  $n$ -forms on  $X$  over the manifold  $B$  determines a compact  $n$ -chain on  $X$  whose boundary, as written in (7.1), has support in  $X - \{s\}$ . The corresponding relative homology class

$$(7.4) \quad \theta_s \in H_n^c(X, X - \{s\}; \mathbf{C}), \quad s \in X,$$

is independent of  $B$ : with the notation above, the compact  $n$ -chain  $\int_B - \int_C$  has support in  $X - \{s\}$ . The relative homology class we have just constructed is often called *the local orientation class*.

(7.5) PROPOSITION. *Let  $s$  be a point of an oriented  $n$ -dimensional smooth manifold  $X$ . The local orientation class  $\theta_s$  generates  $H_n^c(X, X - \{s\}; \mathbf{C})$ .*

*Proof.* With the terminology from section 5 we may express formula (7.1) by means of the local orientation class

$$(7.6) \quad \text{Tr}(\omega; s) = \langle \theta_s, \partial\omega \rangle = \langle b\theta_s, \omega \rangle, \quad \omega \in H^{n-1}(X - \{s\}, \mathbf{C}).$$

In case  $n > 2$  we conclude from (7.3), that  $\theta_s \neq 0$ . The case  $n = 2$  is left with the reader.

Q.E.D.

Let us remark that formula (7.2) shows how to identify  $\theta_s$  under relative Poincaré duality (6.6).

(7.7) PROPOSITION. *Let  $S$  be a finite subset of the oriented  $n$ -dimensional compact manifold  $X$ . For any closed form  $\omega \in \Gamma(X - S, \Omega^n)$  we have that*

$$\sum_{s \in S} \text{Tr}(\omega; s) = 0.$$

*Proof.* Let the fundamental class  $\theta \in H_n(X, \mathbf{C})$  be given by

$$\langle \theta, \omega \rangle = \int_X \omega, \quad \omega \in \Gamma(X, \Omega^n).$$

Let us consider a point  $s \in S$  and use the notation from (7.1). The difference  $\int_X - \int_B$  has support in  $X - \{s\}$ , which shows that the image of  $\theta$  in  $H_n(X, X - \{s\}; \mathbf{C})$  is  $\theta_s$ . We have that

$$\sum_{s \in S} \text{Tr}(\omega; s) = \sum_{s \in S} \langle \theta_s, \partial\omega \rangle = \langle \theta_S, \partial\omega \rangle = \langle b\theta_S, \omega \rangle$$

where  $\theta_S$  denotes the restriction of  $\theta$  to  $H_n(X, X - S; \mathbf{C})$ . Conclusion by the fact that  $b\theta_S = 0$ . Q.E.D.

(7.8) *Definition.* Let  $\gamma$  be a compact  $n$ -chain on the oriented  $n$ -dimensional smooth manifold  $X$ . For a point  $s \in X$  outside  $\text{Supp}(b\gamma)$  the class of  $\gamma$  in  $H_n^c(X, X - \{s\}; \mathbf{C})$  can be written

$$[\gamma] = \text{Ind}(\gamma; s)\theta_s, \quad \text{Ind}(\gamma; s) \in \mathbf{C}.$$

The number  $\text{Ind}(\gamma; s)$  is called the *winding number* of  $\gamma$  with respect to  $s$ .

(7.9) *Example.* Let  $K$  denote an  $n$ -dimensional compact submanifold with boundary. Integration over  $K$  defines a compact  $n$ -chain  $\kappa$  with  $\text{Supp}(\partial\kappa) = \partial K$ . The winding number for  $\kappa$  is 1 in the interior of  $K$  and 0 outside  $K$ .

(7.10) *THEOREM.* Let  $\gamma$  be a compact  $n$ -chain on the oriented  $n$ -dimensional smooth manifold  $X$ . The winding number  $s \mapsto \text{Ind}(\gamma; s)$  is a locally constant function on the complement of  $\text{Supp}(b\gamma)$  in  $X$ . This function is zero outside some compact subset of  $X$  containing  $\text{Supp}(b\gamma)$ .

*Proof.* Let us consider an arbitrary open subset  $U$  of  $X$  containing  $\text{Supp}(b\gamma)$ . We shall now use relative Poincaré duality to describe the class of  $\gamma$  in  $H_n^c(X, U; \mathbf{C})$ . According to (6.6) and (6.7) we can represent  $\gamma$  by a relative  $n$ -chain of the form

$$\langle \gamma, v \rangle = \int_X \rho v, \quad v \in \Gamma(X, \Omega^n)$$

where  $\rho$  is a compactly supported smooth function on  $X$ , constant in a neighbourhood of any point  $s$  of  $Z = X - U$ . Let us notice that

$$\langle \partial\gamma, \omega \rangle = (-1)^n \int \omega \wedge d\rho, \quad \omega \in \Gamma(U, \Omega^{n-1}), \quad d\omega = 0.$$

In order to calculate  $\text{Ind}(\gamma; s)$  we replace  $U$  by a small pointed neighbourhood  $D^*$  of  $s$ . With the notation of (7.2) let us write  $\rho = \rho(s)\rho_s$  and deduce that

$$\langle \partial\gamma, \omega \rangle = \rho(s)\text{Tr}(\omega; s), \quad \omega \in \Gamma(D^*, \Omega^{n-1}), \quad d\omega = 0.$$

We can now conclude from (7.6) that

$$\text{Ind}(\gamma; s) = \rho(s), \quad s \in X - U.$$

This reveals that  $s \mapsto \text{Ind}(\gamma; s)$  is a compactly supported, locally constant function on  $X - U$ .

For a given fixed point  $s \notin \text{Supp}(b\gamma)$  choose  $U$  to be an open neighbourhood of  $\text{Supp}(b\gamma)$  with  $\bar{U}$  compact and  $s \notin U$ . We can apply the considerations above and conclude that the winding number is constant in a neighbourhood of  $s$  and zero outside some compact neighbourhood of  $\text{Supp}(b\gamma)$ . Q.E.D.

(7.11) COROLLARY. *Let  $\gamma$  be a compact  $n$ -chain on the oriented smooth manifold  $X$  and  $U$  an open subset of  $X$  containing  $\text{Supp}(b\gamma)$ . The relative de Rham homology class*

$$[\gamma] \in H_n^c(X, U; \mathbf{C})$$

*is zero if and only if  $\text{Ind}(\gamma; s) = 0$  for all  $s \in X - U$ .*

*Proof.* This is a corollary to the proof of (7.10) rather than the statement (7.10). Anyway, the basic point is Poincaré duality (6.6). Q.E.D.

## 8. CAUCHY'S RESIDUE THEOREM

We shall consider a smooth map  $\gamma: S^{n-1} \rightarrow E$  from the oriented  $n-1$  sphere into an oriented  $n$ -dimensional real vector space  $E$ . For a point  $s$  outside  $\gamma(S^{n-1})$  pick a closed  $(n-1)$ -form  $\omega_s$  on  $E - \{s\}$  with  $\text{Tr}(\omega_s; s) = 1$  and define the *winding number* of  $\gamma$  with respect to  $s$  to be

$$(8.1) \quad \text{Ind}(\gamma; s) = \int_{S^{n-1}} \gamma^* \omega_s.$$

(8.2) CAUCHY'S RESIDUE THEOREM. *Let  $\gamma: S^{n-1} \rightarrow X$  denote a smooth map into an open subset  $X$  of  $E$  with  $\text{Ind}(\gamma; z) = 0$  for all  $z \in E - X$ .*