

Introduction

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **35 (1989)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **25.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

KNOTTING CODIMENSION 2 SUBMANIFOLDS LOCALLY

by Mikiya MASUDA and Makoto SAKUMA

INTRODUCTION

Let L be a connected oriented n -dimensional closed manifold smoothly embedded in a connected oriented $(n+2)$ -dimensional closed manifold M , and let K be an oriented n -dimensional smooth knot in the oriented S^{n+2} . Then we consider the connected sum $(M, L) \# (S^{n+2}, K)$. In other words, we knot L locally using K . It yields another embedding of L in M ; however, it does not always give a new embedding. In fact, the lightbulb theorem says that the connected sum of $(S^2 \times S^1, \{*\} \times S^1)$ with any knot in S^3 is always equivalent to the original embedding. Moreover, by the prime decomposition theorem for knots in 3-manifolds [My], $(S^2 \times S^1, \{*\} \times S^1)$ is essentially the only embedding of a circle with the above property. Litherland [Li] has generalized the lightbulb theorem to the higher dimensional cases. In the appendix of [V], Viro exhibits an example of a 2-knot whose connected sum with the standard projective plane in S^4 does not change the isotopy type of the projective plane. (See also [La].)

The purpose of this paper is to study under what conditions this phenomenon occurs (or does not occur). The first named author [Ms] studied this problem when the codimension is greater than 2.

Put it in another way. Let \mathcal{K}_n be the set of isotopy classes of oriented n -knots diffeomorphic to S^n in the oriented S^{n+2} . The set forms an abelian monoid under connected sum for pairs. Analogously to the inertia group of a manifold, we define

$$I(M, L) = \{(S^{n+2}, K) \in \mathcal{K}_n \mid (M, L) \# (S^{n+2}, K) = (M, L)\}$$

where $=$ in the parenthesis indicates that there is an orientation preserving diffeomorphism of pairs. The set forms a submonoid of \mathcal{K}_n and describes the effect of knotting L locally. We are also concerned with the following intermediate submonoid

$$I_0(M, L) = \{(S^{n+2}, K) \in I(M, L) \mid (M, L) \# (S^{n+2}, K) \equiv (M, L)\}$$

where \equiv indicates that there is an orientation preserving diffeomorphism of pairs which is concordant to the identity map as a diffeomorphism of the ambient space M .

Our results suggest that $I(M, L)$ and $I_0(M, L)$ depend only on the order of a meridian of L in $\pi_1(M - L)$ or $H_1(M - L; \mathbf{Z})$. Roughly speaking, according as the order is infinite, 1, or p ($1 < p < \infty$), they can be distinguished by (at least) these three types:

Type 1 $I(M, L) = \{0N\}$,

Type 2 $I(M, L) = \mathcal{K}_n$, $I_0(M, L) = \ker \sigma$,

Type 3 $\{0\} \subsetneq I(M, L) \subsetneq \mathcal{K}_n$, $\{0\} \subsetneq I_0(M, L) \subsetneq \ker \sigma$,

(see section 4 for $\sigma(S^{n+2}, K)$).

We refer the reader to 1.1, 2.6, 3.4, 5.1, 5.2, and 5.8 for the precise statement.

This paper consists of five sections. In Section 1, we deduce a necessary condition for $I_0(M, L)$, which is valid for any (M, L) . We treat type 1 in Section 2. Type 2 is discussed in Sections 3, 4 and type 3 is discussed in Section 5. We will find that type 3 is closely related to the generalized Smith conjecture.

The authors would like to express their hearty thanks to Professors A. Kawauchi and T. Maeda for helpful conversations and suggestions.

§ 1. GENERAL REMARKS ON $I_0(M, L)$

It is known (and it is easily verified) that the signature of a Seifert surface of an oriented n -knot K in S^{n+2} is independent of the choice of a Seifert surface; so it is an invariant of the oriented knot K . The invariant is called the signature of the knot K and denoted by $\text{Sign}(S^{n+2}, K)$. We note that $\text{Sign}(S^{n+2}, K)$ is trivially zero unless $n + 1 \equiv 0 \pmod{4}$.

As is seen in Section 3, there is a pair (M^{n+2}, L^n) such that $I(M, L) = \mathcal{K}_n$ for any $n \geq 3$. In contrast, we can deduce a necessary condition for $I_0(M, L)$ which holds for any pair (M, L) .

THEOREM 1.1. *If $(S^{n+2}, K) \in I_0(M, L)$, then $\text{Sign}(S^{n+2}, K) = 0$.*

Proof. Let V be a Seifert surface of K . Since $S^{n+2} = \partial D^{n+3}$, we can push the interior of V into the interior of D^{n+3} so that V is transverse to S^{n+2} . This yields an oriented pair (D^{n+3}, V) having (S^{n+2}, K) as the boundary.