

§2. Type 1 case

Objekttyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **35 (1989)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **26.05.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern.

Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden.

Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

§ 2. TYPE 1 CASE

In this section we consider the case where a meridian of L^n in M^{n+2} has infinite order in $H_1(M-L; \mathbf{Z})$. We shall denote by $[m]$ the homology class in $H_1(M-L; \mathbf{Z})$ represented by a meridian m of L in M . For a manifold pair (X, Y) of codimension 2 and an epimorphism γ from $\pi_1(X-Y)$ to a finite group, let $(X, Y)_\gamma$ be the branched covering of (X, Y) corresponding to γ . Each knot group $\pi_1(S^{n+2}-K)$ has a natural epimorphism to \mathbf{Z}_p for any positive integer p , and the corresponding p -fold branched cyclic covering of (S^{n+2}, K) is denoted by $(S^{n+2}, K)_p$.

LEMMA 2.1. *Suppose $[m]$ is of infinite order. Then if $(S^{n+2}, K) \in I(M, L)$ then $(S^{n+2}, K)_p$ is a homotopy $(n+2)$ -sphere for any positive integer p .*

Proof. Since $[m]$ represents a nontrivial element in the finitely generated free abelian group $B_1(M-L) \equiv H_1(M-L; \mathbf{Z})/\text{Tor } H_1(M-L; \mathbf{Z})$, there is a positive integer r and a primitive element x in $B_1(M-L)$ such that $[m] = rx$ in $B_1(M-L)$. For each positive integer p , let γ_p be the canonical epimorphism $\pi_1(M-L) \rightarrow B_1(M-L) \otimes \mathbf{Z}_{pr}$. Noting the naturality of the homomorphism γ_p , we can see the following:

$$\begin{aligned} (M, L)_{\gamma_p} &= ((M, L) \# (S^{n+2}, K))_{\gamma_p \circ f_*} \\ &= (M, L)_{\gamma_p} \# d_p(S^{n+2}, K)_p \end{aligned}$$

Here f is a diffeomorphism $(M, L) \# (S^{n+2}, K) \rightarrow (M, L)$ and d_p is the order of $B_1(M-L) \otimes \mathbf{Z}_{pr}$ divided by p . Hence $H_*(S^{n+2}, K)_p; \mathbf{Z} \simeq H_*(S^{n+2}; \mathbf{Z})$ and $\pi_1((S^{n+2}, K)_p) \simeq 1$ by the existence of prime decompositions of finitely generated groups into free products [Wg]. Q.E.D.

It is conjectured that those knots which satisfy the conclusion of the above lemma are trivial. In fact, for $n = 1$, it follows from the Smith conjecture [MB]. As a supporting evidence for higher dimensional cases, we have

LEMMA. *Suppose that $(S^{n+2}, K)_p$ is a homotopy $(n+2)$ -sphere for every positive integer p . Then the Alexander modules of K are trivial.*

Proof. Let $\tilde{E}(K)$ be the infinite cyclic cover of the exterior $E(K)$ of K in S^{n+2} , and let t denote the automorphism of the homology group of $\tilde{E}(K)$ induced by the action of a meridian. Then, by the arguments of [Sm1],

we can see that $t^p - 1 : H_q(\tilde{E}(K); \mathbf{Z}_r) \rightarrow H_q(\tilde{E}(K); \mathbf{Z}_r)$ is an isomorphism for any positive integers p , q , and r . Assume r is prime. Then $H_q(\tilde{E}(K); \mathbf{Z}_r)$ is a finite abelian group, since it is a finitely generated torsion module over the principal ideal domain $\mathbf{Z}_r\langle t \rangle$ (see [Le3, p. 8]). So the automorphism t on $H_q(\tilde{E}(K); \mathbf{Z}_r)$ has a finite order, say d , and we have $t^d - 1 = 0$. Hence $H_q(\tilde{E}(K); \mathbf{Z}_r) = 0$, and by the universal coefficient theorem, the following holds for any prime r and any positive integer q :

$$(2.3) \quad H_q(\tilde{E}(K); \mathbf{Z}) \otimes \mathbf{Z}_r = 0$$

$$(2.4) \quad \text{Tor}(H_q(\tilde{E}(K); \mathbf{Z}), \mathbf{Z}_r) = 0$$

By (2.4), $H_q(\tilde{E}(K); \mathbf{Z})$ has no nontrivial elements of finite order; so it has a square presentation matrix $M(t)$ as a $\mathbf{Z}\langle t \rangle$ -module by [Le3, Proposition 3.5]. By (2.3) the q -th Alexander polynomial $\det M_q(t)$ ($\in \mathbf{Z}\langle t \rangle$) is a unit mod. r for any prime r . Hence it is a unit in $\mathbf{Z}\langle t \rangle$, and we have $H_q(\tilde{E}(K); \mathbf{Z}) = 0$ for any positive integer q . Q.E.D.

Thus, as a consequence of Lemmas 2.1 and 2.2 and the results of [Le2] and [T], we have the following:

PROPOSITION 2.5. *Suppose $[m]$ is of infinite order. Then any knot in $I(M, L)$ has trivial Alexander modules and is null cobordant.*

Hence the only obstruction for a knot (S^{n+2}, K) in $I(M, L)$ to be trivial lies in the knot group $\pi_1(S^{n+2} - K)$. For the special case where $[m]$ generates $H_1(M - L)$, we can apply the result of Maeda [Ma] (cf. [DF]), and obtain the following:

THEOREM 2.6. *Suppose $n \geq 3$ and $H_1(M - L)$ is the infinite cyclic group generated by $[m]$. Then $I(M, L)$ is trivial.*

Proof. Let (S^{n+2}, K) be a knot in $I(M, L)$. Note that $\pi_1(M - L)$ is isomorphic to the amalgamated free product $\pi_1(M - L) *_{\langle m \rangle} \pi_1(S^{n+2} - K)$.

Then we can conclude $\pi_1(S^{n+2} - K) \simeq \mathbf{Z}$ by the result of [Ma] (cf. [DF]) which asserts the existence of a prime decomposition of a finitely presented group G with $G/[G, G] \simeq \mathbf{Z}$ with respect to such amalgamated free products. Combined with Proposition 2.5, we see $S^{n+2} - K$ is homotopy equivalent to a circle. Hence (S^{n+2}, K) is trivial by [Le1].